Evidence for Rehabilitation Interventions After Acute Lateral Ankle Sprains in Athletes: A Scoping Review

Tee, Eugene; Melbourne, Jack; Sattler, Larissa; Hing, Wayne

Published in:
Journal of Sport Rehabilitation

DOI:
[10.1123/jsr.2021-0244](https://doi.org/10.1123/jsr.2021-0244)

Licence:
Other

[Link to output in Bond University research repository.](#)

Recommended citation(APA):
Tee, E., Melbourne, J., Sattler, L., & Hing, W. (2022). Evidence for Rehabilitation Interventions After Acute Lateral Ankle Sprains in Athletes: A Scoping Review. *Journal of Sport Rehabilitation*, 31(4), 457-464.
<https://doi.org/10.1123/jsr.2021-0244>

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

For more information, or if you believe that this document breaches copyright, please contact the Bond University research repository coordinator.

Title:

Evidence for rehabilitation interventions after acute lateral ankle sprains in athletes: A scoping review

Running title:

Rehabilitation after acute lateral ankle sprain

Authors

Eugene Tee¹ (DPT), Jack Melbourne¹ (DPT), Larissa Sattler¹ (PhD), Wayne Hing¹ (PhD)

Institutions:

¹ Bond University Institute of Health and Sport, 2 Promethean way, Robina, QLD, Australia 4226

Author details:

Eugene Tee (Corresponding Author)

Position: Physiotherapist

Institution: Bond Institute of Health and Sport, Bond University

Postal address: 6/30 Crump Street, Holland Park West, 4121, QLD, Australia

Email: eugenetee93@gmail.com

Jack Melbourne

Position: Physiotherapist

Institution: Bond Institute of Health and Sport, Bond University

Email: jack.melbourne12@outlook.com

Dr Larissa Sattler

Assistant Professor – Doctor of Physiotherapy program, Bond University

Email: lsattler@bond.edu.au

Dr Wayne Hing

Professor – Doctor of Physiotherapy program, Bond University

Email: whing@bond.edu.au

All authors (ET, JM, LS, WH) certify that he or she has no commercial associations (e.g., consultancies, stock ownership, equity interest, patent/licensing arrangements, etc.) that might pose a conflict of interest in connection with the submitted article.

Title: Evidence for rehabilitation interventions after acute lateral ankle sprains in athletes: A scoping review

ABSTRACT

Context: Acute lateral ankle sprain (LAS) is a common injury in athletes and is often associated with decreased athletic performance, and if treated poorly, chronic ankle issues such as instability. The physical performance demands such as cutting, hopping and landing involved with certain sport participation suggests that the rehabilitation needs of an athlete after LAS may differ from the those of the general population.

Objective: To review the literature to determine the most effective rehabilitation interventions reported for athletes returning to sport after acute LAS.

Evidence Acquisition:

Data Sources: Databases Pubmed, Embase, Cinahl, SportDiscus and PEDro were searched to July 2020.

Study Selection: A scoping review protocol was developed and followed in accordance with the PRISMA-ScR guidelines and prospectively registered (<https://osf.io/bgek3/>). Study selection included published articles on rehabilitation for ankle sprain in an athletic population.

Data Extraction: Parameters included athlete and sport type, age, sex, intervention investigated, outcome measures, measurement tool and follow up period.

Data Synthesis: A qualitative synthesis for all articles was undertaken, and a quantitative sub-analysis of randomized controlled trials (RCT) and critical methodological appraisal was also conducted.

Evidence Synthesis: 37 articles were included in this review consisting of 5 systematic and 20 narrative reviews, 7 RCTs, a single case series, case report, position statement, critically appraised topic, and descriptive study were also included. RCT interventions included early dynamic training, electrotherapy, and hydrotherapy.

Conclusions: Early dynamic training, after acute LAS in athletes, results in a shorter time to return to sport, increased functional performance, and decreased self-reported re-injury. The results of this scoping review support an early functional and dynamic rehabilitation approach when compared to passive interventions for athletes returning to sport after LAS. Despite existing research on rehabilitation of LAS in the general population, a lack of evidence exists related to athletes seeking to return to sport.

Keywords: foot, rehabilitation, physiotherapy, physical therapy, return to sport.

78

79 **Introduction**

80 Lateral ligament injuries to the ankle are a common musculoskeletal complaint, with an
81 estimated rate of one ankle injury per 10,000 people per day ¹⁻³. In a sporting population, the
82 incidence is even greater, constituting 15% to 45% of all sport related injuries ³. When an
83 athlete sustains a lateral ankle sprain (LAS) this can often result in lost playing time, decreased
84 athletic performance, and if treated poorly, chronic ankle issues such as instability ⁴.
85 Additionally, LAS presents a significant economic burden when medical costs and leave from
86 work are taken into consideration ⁵. The high frequency of LAS in athletes, alongside the costs
87 of injury management, demonstrate the importance for rehabilitation interventions to be
88 evidence-based.

89

90 Research to date, supports non-operative management for excellent long-term outcomes when
91 compared with surgical intervention in cases not involving a complete ligament rupture ⁶⁻⁹. The
92 increased costs, intra and post-operative complication risks and prolonged absence that are
93 associated with surgical interventions are also reasons that make conservative management
94 more favorable ⁹⁻¹¹. Clinical practice guidelines have been developed with respect to
95 rehabilitation after LAS, however, these do not appear to inform the efficacy of rehabilitation
96 interventions specific to athletes wishing to return to sport (RTS)^{10,12}.

97

Considering the greater physical requirements such as cutting, jumping, hopping and landing associated with certain sport participation it would be reasonable to suggest that an athlete's rehabilitation plan should differ from that of the general population. Therefore, this review aims to scope the literature to determine the most effective rehabilitation interventions reported for athletes returning to sport after sustaining a LAS.

Materials and Methods

Study design

A scoping review study design was utilised to identify key concepts, knowledge gaps and types of evidence currently available and the protocol was prospectively registered with Open Science Framework (<https://osf.io/bgek3/>)¹³.

Data Sources and Searches

Two researchers (ET & JM) conducted the literature search to identify, screen and select studies to be included in accordance with the Preferred Reporting Items for Systematic Reviews and Meta-Analysis Extension for Scoping Reviews (PRISMA-ScR) ¹⁴. The search strategy was developed through the methodological procedure proposed by Arksey and O'Malley ¹³ and Johanna Briggs Institute ¹⁵, and followed a 3-step approach:

1. A pilot search of the database PubMed using the subject headings: (acute) AND (ankle) AND 'sprain OR injury' AND 'treat* OR manage* OR rehab* OR therap*' AND 'sport* OR athlet* OR 'funct* OR exercis*'.
 2. Identified key words and terms relating to acute lateral ankle sprains and rehabilitation interventions and return to sport.
 3. Execution of the final search strategy and further searching of referenced lists of the selected articles, including systematic and narrative reviews.

A search was formulated and conducted in five databases (Pubmed, Embase, Cinahl, SportDiscus and PEDro) from inception to 27 July 2020. Articles were downloaded to the EndNote reference management software (<http://www.endnote.com/>) for selection by ET & JM according to the PRISMA-ScR statement ¹⁴.

Study Selection

The eligibility criteria were defined by the *population* (**athletes** clinically diagnosed with acute lateral ankle sprain), *concept* (any study reporting on the effects of rehabilitation interventions for acute lateral ankle sprain) and *context* (all periods of time, outcomes, comparators, follow up, return to sport setting and during and type of intervention). The following types of publications were eligible for inclusion: intervention studies, narrative reviews, scoping reviews, systematic reviews, meta-analysis, case series and clinical commentaries. Exclusion criteria were (1) non-English language, **(2) non athlete only population**, (3) studies involving animals, (4) chronic ankle instability population, (5) fractures and/or dislocations, (6) deltoid ligament,

peroneal, syndesmosis injuries, (7) preventative interventions, (8) surgical interventions, and (9) pharmacological interventions. The following were also excluded from our review: conference abstracts/proceedings, magazines and newspaper articles, grey literature and in those articles where the full text was not available.

Data Extraction and Quality Assessment

Data extraction, categorization and mapping was performed by ET & JM using methodology adapted from the Joanna Briggs Institute (JBI) guidelines for scoping reviews ¹⁵. The data were extracted into the following categories: (1) author, (2) year, (3) aim/purpose, (4) study population (including demographic details) and size, (5) study design, (6) intervention (type, comparator, duration and details), (7) outcomes, and (8) key findings.

Data synthesis and Methodological quality appraisal

A qualitative synthesis for all articles was undertaken, and a further quantitative synthesis of randomized controlled trials (RCT) and methodological quality appraisal was included. For critical appraisal two reviewers (ET & JM) independently scored studies using the PEDro scale, where disagreement existed a consensus decision was made through discussion. The Kappa coefficient statistic was used to measure the interrater reliability and agreement between the two reviewers ¹⁶. The PEDro scale is a methodological measure of quality used to appraise RCTs, consisting of an 11-item checklist with a maximum possible score of 10 ¹⁷. Studies of good

quality received a score of greater than 6, fair quality scores were between 4 and 6, and poor studies received scores of less than 4^{18,19}.

Results

The search strategy yielded a total 2600 articles, of which 2525 records were removed as duplicates or not meeting the pre-defined inclusion criteria. Upon completion of the title and abstract screening, 75 articles were retrieved and assessed for eligibility. Subsequently, 38 full text articles fulfilled the eligibility criteria and were included (Figure 1).

The number of published studies relating to rehabilitation interventions following LAS to RTS has been consistent over the last 20 years. Of the 37 articles included in this scoping review, 25 were review articles, comprised of 20 narrative reviews and 5 systematic reviews. Also included were a critically appraised topic, position statement and a descriptive study were also identified (Supplementary Figure 1). The five systematic reviews investigated a range of interventions including bracing, functional treatment, cryotherapy, manual mobilisation and exercise²⁰. There were also 7 RCTs included, along with a case series, and a case report.

Characteristics of the included reviews

There were five systematic reviews reported in this review containing 82 original studies (Table 1). The methodological quality of the included articles within the systematic reviews were

classified as poor or limited quality. Additionally, methodological quality was not reported in all of the included reviews²¹. The five systematic reviews contained literature from 1990-2012 and there were several outcomes investigated for the rehabilitation of LAS. Two of the reviews investigated early dynamic training^{9,22}. The remaining three articles explored bracing vs functional treatment²¹, cryotherapy²³ and exercise and manual therapy²⁴. A critically appraised topic was included containing 3 studies exploring deep oscillation therapy in athletes with acute LAS ²⁵. A position statement from 2012 proposing an evidence-based clinical guideline to treat LAS was also included in this review¹⁰. This position statement explored the use of early dynamic training, ice and compression, immobilization, electrotherapy and manual mobilizations. Additionally, a descriptive study relating to the clinical application of cryotherapy was also included in this review. The modalities of cryotherapy surveyed included an ice pack, ice pack and cryokinetics, ice immersion, and ice immersion and cryokinetics ²⁶.

Characteristics of the included intervention studies

Participants

There was a total of 830 participants across the 7 RCTs and the overall sex ratio was 57% male and 43% female. All RCTs contained both male and female participants apart from one ²⁷ which included 22 female participants only and the average mean age was 27 years. Two studies involved professional athletes (29%), with the remainder of participants in the five studies (71%) recreationally or competitively active. The type of sport was unspecified in five studies (62.5%). The remainder included the following sports: soccer, football, basketball, mixed martial arts, volleyball, handball and running. Table 2 provides a summary of

participant characteristics across each intervention study including population, type of sport, sample size, age (mean) and gender. The case series contained 3 intercollegiate athletes comprised of 2 males aged 18 and 21 and single female aged 18. The sports of the participants were women's lacrosse and men's basketball and soccer²⁸. The case report examined a 29-year-old male Caucasian professional football athlete ²⁹.

Intervention type and effectiveness

The most investigated intervention in the RCTs was early dynamic training (5 studies) ³⁰⁻³⁴ , while one study examined the electro modality, non-invasive interactive neurostimulation ³⁵, and one study aquatic therapy ²⁷. Early dynamic training included a combination of early weightbearing, range of motion (ROM) exercises, proprioceptive or balance training, running and plyometrics, resistance strength training, or functional rehabilitation. Hydrotherapy showed more effective improvements in pain than standard land-based therapy ²⁷. Non-invasive interactive neurostimulation was effective in improving short-term outcomes, however showed no significant difference in long-term outcomes to RTS ³⁵.

Early dynamic training was effective in increasing functional performance, strength, postural control, decreasing recess fluid, and a shortened time to RTS ³⁰⁻³⁴. All reported key findings and intervention details can be found in Tables 3 and 4, respectively. The case series explored the use of the Mulligan Concept Mobilization with Movement and all three participants from this study showed an increase in pain-free function at the one month follow up ²⁸. The application of acute hydrotherapy with super saturated hydrogen-rich water for acute LAS was supported

in the case report. This study indicated multi session hydrotherapy within the first 24 hours may provide benefits in terms of pain, swelling reduction, and ROM ²⁹.

Outcome Measures

Although a range of outcome measures were utilized across the included RCTs (Table 4), the most common were those related to pain and swelling. The visual analogue scale (VAS) was frequently used to assess pain while the measurement tool to assess swelling was inconsistent across the studies. Only two RCT's listed time until RTS as an outcome measure ^{30,34}. The follow up time for an outcome measure varied between studies with the shortest being three days and longest 18 months ^{34,36}. Not all studies included a follow up assessment ^{27,30,32}. The case series utilized outcome measures examining pain, functional limitations, balance, and self-reported ability to weight bear ²⁸. The case report focused on outcome measures relating to pain, swelling and weight bearing during lunge ²⁹.

Methodological Quality

The methodological quality of the seven RCT's was assessed using the PEDro scale (Supplementary Table 1). The methodological quality in the included articles varied, ranging from 3 to 8 points out of a possible 10 (Table 3). The mean PEDro score was 5.42, meaning overall, the studies are classified as fair quality. One study ³¹ demonstrated poor methodological quality, scoring 3/10. Four studies (57%) were classified as fair quality and two studies (28%) classified as good quality. None of the studies blinded subjects or therapists. All studies had a between group comparison.

The interrater agreeability between the two reviewers was 81% (6 of 7 items). The Kappa score for the measure of agreement between the two reviewers on the individual validity was 0.86 indicating an almost perfect agreement ¹⁶. The case series noted a limited number of participants, lack of control group, and randomization of selected interventions as a limitation of the study ²⁸. Similarly, the case report also cited the small number of participants along with a relatively short period of intervention and no medium- or long-term efficacy of topical hydrogen as limitations ²⁹.

Discussion

Main findings

The purpose of this scoping review was to systematically identify the scientific literature reporting on rehabilitation interventions for athletes RTS after acute LAS, and to (1) provide the type and efficacy of interventions utilised (2) summarize the studies completed to date, and (3) identify any knowledge gaps to inform future research. This review demonstrated that while there was supporting evidence for the use of early dynamic training to rehabilitate athletes post-acute LAS, the evidence was of low-moderate quality ³⁰⁻³⁴. These findings are consistent with existing literature that recommends a combination of early weight bearing exercises such as proprioception, resistance, plyometric and functional training³⁷⁻⁴¹.

263 An ankle sprain can result in reduced strength and postural control which can increase risk of
 264 re-injury and delay RTS. Given the high level manoeuvres associated with sport, rehabilitation
 265 interventions targeting strength, force production and functional endurance have shown to
 266 have favourable outcomes^{30 34 31 32 33}. In the early stages of a rehabilitation program following
 267 LAS, hydrotherapy was shown to provide advantages over standard land-based therapy for a
 268 sooner RTS²⁷. **Following the acute stage, progressive controlled rehabilitation involving jump**
 269 **stretch flex bands and postural sway showed positive outcomes following LAS and should be**
 270 **considered for inclusion to RTS training protocols. Additionally, rehabilitation programs**
 271 **involving plyometric exercises such as cone hops with 180 degree turn and diagonal cone**
 272 **hops improved functional performance after LAS with the combined effect of strength and**
 273 **speed enhancement**^{30 31}. While passive treatments such as non-invasive interactive
 274 neurostimulation may improve short-term outcomes such as pain, however long-term
 275 outcomes show no significant difference on an athlete's ability to RTS³⁵.
 276 The results of one systematic review favoured of early mobilisation, early exercise and external
 277 ankle support, which may allow a safer and quicker RTS²¹. Although this review provided
 278 support for functional treatment, the population of interest was not athletes only. Moreover,
 279 none of the included systematic reviews had a sole focus on athletes RTS, instead athletes were
 280 combined with the general population in all analyses^{9,21-23,42}. Most sports require movement
 281 patterns that involve a combination of forces in multiple directions, whilst maintaining the need
 282 for optimal technique⁴³. Given the increased multidirectional forces associated with sports
 283 when compared with a typical walking pattern, the increased demands of an athletic
 284 population when developing a rehabilitation program should be considered. However, the

current body of evidence available is largely focused on non-athletic populations or recreational athletes which utilise RTS as a secondary outcome.

Limitations

Limitations of this review include the lack of RCTs available on LAS with respect athletes and RTS and the relatively poor methodological quality of those studies that were available. Although the inclusion criteria for this scoping review was broad, articles not available in English or full text were excluded. Due to the considerable heterogeneity of the interventions and outcomes reported, it remains a challenge for clinicians working with a RTS population to base rehabilitation protocols after LAS on evidence-based practice.

Clinical and Research Implications

Clinicians should utilize the best available evidence within clinical practice to maximize positive outcomes. Through **sport specific** rehabilitation, athletes may report less weakness, and achieve greater return to baseline biomechanics and motor patterns of sporting movements. **Clinicians should monitor the progression through a multidisciplinary approach involving the medical team, coaches and athlete using sport specific outcomes measures, on field performance and the athlete's confidence.** Our findings are consistent with other recommendations which suggest that an active rehabilitation protocol is superior to passive treatment techniques for RTS ¹². A combination of short-term immobilization, external ankle support, and early dynamic training such as proprioception, strength and balance exercises,

should be utilized to produce optimal outcomes for athletes wishing to RTS ^{9,22,39}. **As previously noted, a limitation of this review were the lack of high quality RCTs investigating these interventions in an athlete only, RTS population. A consideration for the direction of future research would be for clinical trial design on this topic to be of high quality and consistent use of relevant outcome measures, such as athlete re-injury utilized.**

Conclusion

The results of this scoping review provide moderate quality evidence to support for an early weightbearing and dynamic exercise rehabilitation approach, in contrast to passive or restrictive interventions in rehabilitation protocols for athletes after LAS. Given the higher physical requirements in certain sports like cutting, jumping and hopping, rehabilitation should be specific to these demands.

Disclosures

Funding sources: none to declare.

Conflict of interests: none to declare.

References

1. Kerkhoffs GM, Van Dijk CN. Acute lateral ankle ligament ruptures in the athlete: the role of surgery. *Foot Ankle Clin.* 2013;18(2):215-218.
2. Kerkhoffs GM, Struijs PA, Marti RK, Assendelft WJ, Blankevoort L, van Dijk CN. WITHDRAWN: Different functional treatment strategies for acute lateral ankle ligament injuries in adults. *Cochrane Database Syst Rev.* 2013(3):Cd002938.
3. Malliaropoulos N, Papacostas E, Papalada A, Maffulli N. Acute lateral ankle sprains in track and field athletes: an expanded classification. *Foot Ankle Clin.* 2006;11(3):497-507.
4. Hertel J, Corbett RO. An Updated Model of Chronic Ankle Instability. *J Athl Train.* 2019;54(6):572-588.
5. Shah S, Thomas AC, Noone JM, Blanchette CM, Wikstrom EA. Incidence and Cost of Ankle Sprains in United States Emergency Departments. *Sports Health.* 2016;8(6):547-552.
6. Dubin JC, Comeau D, McClelland RI, Dubin RA, Ferrel E. Lateral and syndesmotic ankle sprain injuries: a narrative literature review. *J Chiropr Med.* 2011;10(3):204-219.
7. Kemler E, van de Port I, Backx F, van Dijk CN. A Systematic Review on the Treatment of Acute Ankle Sprain. *Sports Medicine.* 2011;41(3):185-197.
8. Polzer H, Kanz KG, Prall WC, et al. Diagnosis and treatment of acute ankle injuries: development of an evidence-based algorithm. *Orthop Rev (Pavia).* 2012;4(1):e5.
9. Seah R, Mani-Babu S. Managing ankle sprains in primary care: what is best practice? A systematic review of the last 10 years of evidence. *Br Med Bull.* 2011;97:105-135.
10. Kerkhoffs GM, van den Bekerom M, Elders LAM, et al. Diagnosis, treatment and prevention of ankle sprains: an evidence-based clinical guideline. *British Journal of Sports Medicine.* 2012;46(12):854-860.

- 347 11. Kyaw SL, Moore, I. S., & Oo, M. L. . A systematic review on the effectiveness of different
 348 functional treatments for acute ankle sprains. *Journal of sports medicine & Doping studies*
 349 2019;9 (1).
- 350 12. Vuurberg G, Hoorntje A, Wink LM, et al. Diagnosis, treatment and prevention of ankle sprains:
 351 update of an evidence-based clinical guideline. *Br J Sports Med.* 2018;52(15):956.
- 352 13. Arksey H, O'Malley L. Scoping studies: towards a methodological framework. *International*
 353 *Journal of Social Research Methodology.* 2005;8(1):19-32.
- 354 14. Tricco AC, Lillie E, Zarin W, et al. PRISMA Extension for Scoping Reviews (PRISMA-ScR): Checklist
 355 and Explanation. *Ann Intern Med.* 2018;169(7):467-473.
- 356 15. Peters M, Godfrey, C., McInerney, P., Soares, C. B., Khalil, H., & Parker, D. . Reviewers Manual
 357 Methodology for JBI Scoping Reviews 2015. 2015;2:3-24.
- 358 16. McHugh ML. Interrater reliability: the kappa statistic. *Biochemia medica.* 2012;22(3):276-282.
- 359 17. Hariom K, Prakash V, Saravankumar J. Quantity and quality of randomized controlled trials
 360 published by Indian physiotherapists. *Perspect Clin Res.* 2015;6(2):91-97.
- 361 18. Kennelly J. Reducing Racial/Ethnic Disparities in Reproductive and Perinatal Outcomes. In:
 362 Springer: New York, NY, USA; 2011.
- 363 19. Lyons K, Radburn C, Orr R, Pope R. A Profile of Injuries Sustained by Law Enforcement Officers: A
 364 Critical Review. *Int J Environ Res Public Health.* 2017;14(2).
- 365 20. **Lynch SA, Renstrom PA. Treatment of acute lateral ankle ligament rupture in the athlete.**
 366 **Conservative versus surgical treatment. *Sports Med.* 1999;27(1):61-71.**
- 367 21. Kemler E, van de Port I, Backx F, van Dijk CN. A systematic review on the treatment of acute
 368 ankle sprain: brace versus other functional treatment types. *Sports Med.* 2011;41(3):185-197.
- 369 22. Petersen W, Rembitzki IV, Koppenburg AG, et al. Treatment of acute ankle ligament injuries: a
 370 systematic review. *Arch Orthop Trauma Surg.* 2013;133(8):1129-1141.

- 371 23. Hubbard TJ, Aronson SL, Denegar CR. Does Cryotherapy Hasten Return to Participation? A
372 Systematic Review. *J Athl Train.* 2004;39(1):88-94.
- 373 24. van der Wees PJ, Lenssen AF, Hendriks EJ, Stomp DJ, Dekker J, de Bie RA. Effectiveness of
374 exercise therapy and manual mobilisation in ankle sprain and functional instability: a systematic
375 review. *Aust J Physiother.* 2006;52(1):27-37.
- 376 25. Hausmann M, Ober J, Lepley AS. The Effectiveness of Deep Oscillation Therapy on Reducing
377 Swelling and Pain in Athletes With Acute Lateral Ankle Sprains. *J Sport Rehabil.* 2019:1-4.
- 378 26. Hawkins SW, Hawkins JR. CLINICAL APPLICATIONS OF CRYOTHERAPY AMONG SPORTS PHYSICAL
379 THERAPISTS. *International Journal of Sports Physical Therapy.* 2016;11(1):141-148.
- 380 27. Kim E, Kim T, Kang H, Lee J, Childers MK. Aquatic versus land-based exercises as early functional
381 rehabilitation for elite athletes with acute lower extremity ligament injury: a pilot study. *Pm r.*
382 2010;2(8):703-712.
- 383 28. May JM, Nasypany A, Paolino J, Baker R, Seegmiller J. Patient Outcomes Utilizing the Mulligan
384 Concept of Mobilization With Movement to Treat Intercollegiate Patients Diagnosed With
385 Lateral Ankle Sprain: An a Priori Case Series. *J Sport Rehabil.* 2017;26(6):486-496.
- 386 29. Javorac D, Stajer V, Ostojic S. Case Report: Acute hydrotherapy with super-saturated hydrogen-
387 rich water for ankle sprain in a professional athlete. *F1000Res.* 2020;9:245.
- 388 30. Van Der Linde E, Oschman Z. Comparative ultrasound study of acute lateral ankle ligament
389 injuries rehabilitated with conventional and jump stretch flex band programmes. *African Journal*
390 *for Physical, Health Education, Recreation & Dance.* 2011;17(4):591-602.
- 391 31. Holme E, Magnusson SP, Becher K, Bieler T, Aagaard P, Kjaer M. The effect of supervised
392 rehabilitation on strength, postural sway, position sense and re-injury risk after acute ankle
393 ligament sprain. *Scand J Med Sci Sports.* 1999;9(2):104-109.

32. Ismail MM, Ibrahim MM, Youssef EF, El Shorbagy KM. Plyometric training versus resistive exercises after acute lateral ankle sprain. *Foot Ankle Int.* 2010;31(6):523-530.
33. Hupperets MD, Verhagen EA, van Mechelen W. Effect of unsupervised home based proprioceptive training on recurrences of ankle sprain: randomised controlled trial. *Bmj.* 2009;339:b2684.
34. Karlsson J, Eriksson BI, Swärd L. Early functional treatment for acute ligament injuries of the ankle joint. *Scand J Med Sci Sports.* 1996;6(6):341-345.
35. Razzano C, Izzo R, Savastano R, Colantuoni C, Carbone S. Noninvasive Interactive Neurostimulation Therapy for the Treatment of Low-Grade Lateral Ankle Sprain in the Professional Contact Sport Athlete Improves the Short-Term Recovery and Return to Sport: A Randomized Controlled Trial. *J Foot Ankle Surg.* 2019;58(3):441-446.
36. Nunes GS, Vargas VZ, Wageck B, Haupenthal DP, da Luz CM, de Noronha M. Kinesio Taping does not decrease swelling in acute, lateral ankle sprain of athletes: a randomised trial. *J Physiother.* 2015;61(1):28-33.
37. Anderson SJ. Acute ankle sprains: keys to diagnosis and return to play. *Phys Sportsmed.* 2002;30(12):29-35.
38. Kennedy JG, Hodgkins CW, Sculco P, Carter T, Robinson SP. Sports injuries of the foot and ankle in the adolescent athlete. *International SportMed Journal.* 2006;7(2):85-97.
39. McKeon PO, Donovan L. A Perceptual Framework for Conservative Treatment and Rehabilitation of Ankle Sprains: An Evidence-Based Paradigm Shift. *J Athl Train.* 2019;54(6):628-638.
40. Hudson Z. Rehabilitation and return to play after foot and ankle injuries in athletes. *Sports Med Arthrosc Rev.* 2009;17(3):203-207.

41. van den Bekerom MP, Kerkhoffs GM, McCollum GA, Calder JD, van Dijk CN. Management of acute lateral ankle ligament injury in the athlete. *Knee Surg Sports Traumatol Arthrosc.* 2013;21(6):1390-1395.
42. van der Wees PJ, Lenssen AF, Hendriks EJM, Stomp DJ, Dekker J, de Bie RA. Effectiveness of exercise therapy and manual mobilisation in acute ankle sprain and functional instability: a systematic review. *Australian Journal of Physiotherapy.* 2006;52(1):27-37.
43. Kolt GS. Research on the ankle in sport. *J Sci Med Sport.* 2013;16(5):387.

Tables

Table 1. Summary of included systematic reviews						
Author (year)	Title	Aim	Dates	Methodological quality	Number of studies	Key findings
Kemler et al.2011	A Systematic Review on the Treatment of Acute Ankle Sprain Brace versus Other Functional Treatment Types	To evaluate the effectiveness of braces as a treatment compared with functional treatments such as ankle tape and elastic bandages	1990-2009	Not assessed	8	A few individual studies reported positive functional outcomes after treatment with an ankle brace compared with other functional methods
Hubbard et al. 2004	Does Cryotherapy Hasten Return to Participation? A systematic review	To search original research addressing the effect of cryotherapy on return to participation after injury	1967-2003	Poor quality evidence	4	Cryotherapy may have a positive effect on return-to-participation measures with the relatively poor quality of the studies reviewed
Van Der Wees et al. 2006	Effectiveness of exercise therapy and manual mobilisation in acute ankle sprain and functional instability: A systematic review	Critically review the effectiveness of exercise therapy and manual mobilisation in acute ankle sprains and functional instability by conducting a systematic review of RCTs	1966-2005	Poor quality evidence	17	Manual mobilisation has an initial positive effect on dorsiflexion ROM
Petersen et al. 2013	Treatment of acute ankle ligament injuries: A systematic review	To perform a systematic literature review of the last 10 years regarding evidence for the treatment and prevention of LAS	2002-2012	Poor quality evidence	20	Majority of LAS can be managed without surgery. The results support a phase adapted non-surgical treatment of acute LAS with a short-term immobilization for grade III injuries followed by a semi-rigid brace

Seah et al. 2011	Managing ankle sprains in primary care: what is best practice? A systematic review of the last 10 years of evidence	To summarize the best available evidence in the last decade for managing ankle sprains in the community	2000-2009	Limited	33	Management is not readily agreed. In mild-to-moderate sprains, functional treatment was better than immobilization. In severe sprains, short period of immobilization in a cast/brace results in a quicker recovery than compression bandage alone. Supervised rehabilitation with conventional treatment can be beneficial
-------------------------	---	---	-----------	---------	----	---

Methodological quality was determined by the individual reviewers not the authors of this study. The conclusion stated is that of the included review in reference to interventions for rehabilitation following LAS. The number of original studies is only those included in each review.

Table 2. RCT Study Characteristics

Author (year)	Title	Population	Sport	Number of participants	Age (Mean)	Sex	
						Male	Female
Razzano et al. (2018)	Non-invasive Interactive Neurostimulation Therapy for the Treatment of Low-Grade Lateral Ankle Sprain in the Professional Contact Sport Athlete Improves the Short-Term Recovery and Return to Sport	Professional athletes with a diagnosis of grade I or II LAS that occurred during a contact sport competition	Soccer, football, basketball, mixed martial arts	61	23	27	10
Kim et al. (2010)	Aquatic Versus Land-based Exercises as Early Functional Rehabilitation for Elite Athletes with Acute Lower Extremity Ligament Injury	Professional athletes with isolated grade I or II ligament injury in ankle	Unspecified	22	26	0	22

Karlsson et al. (1996)	Early functional treatment for acute ligament injuries of the ankle joint	Recreational or competitive athletes with acute (<24 h) grade I or II lateral ligament ruptures	Unspecified	86	22	57	29
Van Der Linde et al. (2011)	Comparative ultrasound study of acute lateral ankle ligament injuries rehabilitated with conventional, and jump stretch flex band programs	Active athletes who sustained grade I or grade II ankle ligamentous injuries	Unspecified	25	21.6	19	6
Holme et al. (1999)	The effect of supervised rehabilitation on strength, postural sway, position sense and re-injury risk after acute ankle ligament sprain	Recreational athletes	Unspecified	92	26	44	27
Ismail et al. (2010)	Plyometric Training Versus Resistive Exercises After Acute Lateral Ankle Sprain	Athletes with grade I or II LAS	Basketball, handball, football, running	22	27	11	11
Hupperts et al. (2007)	Effect of unsupervised home-based proprioceptive training on recurrences of ankle sprain	Recreational athletes with a LAS < 2 months	Unspecified	522	48	136	120

Table 3. Study aims and key findings

Author (year)	Aim	Key Findings/Conclusion	PEDro score /10
Razzano et al. (2018)	To compare the results of improvement of a foot functional score, lower level of reported pain, and return to sports in 2 groups of contact sport athletes affected by a grade I or II LAS	Non-invasive interactive neurostimulation can improve short-term outcomes that can hasten RTS in athletes with acute grade I or II ankle sprain	6
Kim et al. (2010)	To compare outcomes between aquatic and land-based exercises during early-phase recovery from acute lower extremity ligament injuries in elite athletes	Aquatic therapy showed more rapid improvement in pain compared with land-based interventions	6

Karlsson et al. (1996)	To compare two different non-surgical treatment modalities in patients with grade I and II LAS	Early dynamic training resulted in an earlier RTS	4
Van Der Linde et al. (2011)	To establish the difference in rehabilitation outcomes between the Jump Stretch Flex Band (JSFB) programme and conventional ankle rehabilitation programmes of acute LAS	The JSFB group showed a decrease in anterior/posterior recess fluid, increase in ligament size and shortened time to RTS	5
Holme et al. (1999)	The effect of an early rehabilitation program, including postural training, on ankle joint function after LAS	Supervised early dynamic training showed an increase in isometric ankle strength and postural control	3
Ismail et al. (2010)	To determine the effects of plyometric training versus resistive exercises on muscle strength and function following acute LAS	Plyometric training improved functional performance after LAS with the combined effect of strength and speed enhancement	6
Hupperets et al. (2007)	To evaluate the effectiveness of an unsupervised proprioceptive training programme on recurrences of ankle sprain after usual care in athletes who had sustained an acute sports related LAS	Proprioceptive training program is effective for the prevention of self-reported recurrences	8

Table 4. Intervention and outcome details

Author (year)	Intervention details	Outcome	Measurement tool: Unit	Follow up
Razzano et al. (2018)	Electro-modalities: Non-invasive interactive neurostimulation vs sham device	1. Pain 2. Self-Reported Functional Impairment Walking Ability	1. VAS: 0-10. 2. Inability Walking Scale (IWS): 0-10	2 & 4 months
Kim et al. (2010)	Aquatic therapy: Aquatic Therapy vs Land Based exercise	1. Pain 2. Static/Dynamic Stability (Biodex Balance System) 3. Percentage single-limb support time	1. VAS: 100mm line. 2. Biodex Medical System, Electric scale level 1-3. 3. Instrumented walkway system, GAITRite:	N/A

		(GAITrite)	seconds	
Karlsson et al. (1996)	Early dynamic training: Compression pads, early weight bearing, ROM training vs compression wrapping, partial weight bearing (PWB) and crutches	1. Stability 2. Pain 3. Swelling 4. Stiffness 5. Work/sport/activities of daily living 6. Running 7. Stair climbing 8. Support	1-8. Orthopaedic surgeon scoring scale: 0-100	18 months
Van Der Linde et al. (2011)	Early dynamic training : Jump stretch flex band program (JSFB) with ultrasound vs conventional ankle rehabilitation with ultrasound	1. Change in ligament size 2. Decrease of fluid in anterior/posterior recess 3. Time until RTS	1-2. Ultrasound machine (7.5 - 14 MHz linear probe): To the closest 0.01mm. 3. Calendar: days	N/A
Holme et al. (1999)	Early dynamic training: Early ankle mobilization, strength, mobility vs supervised physiotherapy	1. Position sense 2. Isometric strength 3. Postural sway	1. Torsiometer: Degrees 2. Dynamometer: Nm 3. Force Platform: Cm	12 months
Ismail et al. (2010)	Early dynamic training: Plyometric training vs Resistive exercises	1. Climbing down-stairs 2. Raising on heel 3. Raising on toes 4. Single limb stance 5. Eversion/inversion strength	1. Stopwatch: seconds 2-3. Number repetitions: Numerical 4. Stopwatch: Seconds 5. Dynamometer: Nm/kg	N/A
Hupperets et al. (2007)	Early dynamic training: Usual care (any form of rehabilitation used by an athlete) vs usual care and proprioceptive training	1. Self- reported Ankle Sprain	1. Number of Incidences: Numerical	12 months

121x127mm (96 x 96 DPI)

Supplementary Material

Supplementary Table 1. Results of the methodological quality appraisal (PEDro Scale)												
Author (year)	Eligibility	Randomised	Concealed	Baseline	Blinding Subjects	Blinding Therapists	Blind Assessors	Adequate Follow up	Intention-to- Treat Analysis	Between- Group Comparisons	Point estimates and variability	Score
Razzano et al. 2018	Yes	Yes	No	Yes	No	No	Yes	Yes	No	Yes	Yes	6
Hupperts et al. 2007	Yes	Yes	Yes	Yes	No	No	Yes	Yes	Yes	Yes	Yes	8
Eriksson et al. 1996	No	Yes	No	No	No	No	No	No	Yes	Yes	Yes	4
Van Der Linde et al. 2011	Yes	Yes	No	No	No	No	No	Yes	Yes	Yes	Yes	5
Ismail et al. 2010	Yes	Yes	Yes	Yes	No	No	Yes	No	No	Yes	Yes	6
Holme et al. 1999	Yes	Yes	No	No	No	No	No	No	No	Yes	Yes	3
Kim et al. 2010	Yes	Yes	No	Yes	No	No	Yes	Yes	No	Yes	Yes	6

Supplementary Figure 1. Number of included studies by study design type

199x132mm (120 x 120 DPI)