Mapping the drivers of overdiagnosis to potential solutions

Pathirana, Thanya; Clark, Justin; Moynihan, Ray

Published in:
BMJ (Clinical research ed.)

DOI:
[10.1136/bmj.j3879](https://doi.org/10.1136/bmj.j3879)

Licence:
CC BY-NC

[Link to output in Bond University research repository.](#)

Recommended citation(APA):
Pathirana, T., Clark, J., & Moynihan, R. (2017). Mapping the drivers of overdiagnosis to potential solutions. *BMJ (Clinical research ed.)*, 358, 1-9. [j3879]. <https://doi.org/10.1136/bmj.j3879>

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

For more information, or if you believe that this document breaches copyright, please contact the Bond University research repository coordinator.

ANALYSIS

TOO MUCH MEDICINE

Mapping the drivers of overdiagnosis to potential solutions

Thanya Pathirana and colleagues explore strategies to tackle the problem of too much medicine

Thanya Pathirana *PhD scholar*, Justin Clark *senior information specialist*, Ray Moynihan *senior research fellow*

Center for Research in Evidence Based Practice, Bond University, Australia

In our collective enthusiasm to diagnose and treat disease, a growing body of evidence indicates that we may often be doing too much of a good thing.¹⁻⁵ “Overdiagnosis” is now widely recognised to occur when people are labelled with or treated for a disease that would never cause them harm—often as a result of undergoing screening—and it can lead to the overuse of further tests and treatments.^{2,6} One example is thyroid cancer, with estimates that over 500 000 people may have received overdiagnoses across 12 countries in the past two decades, leading to unnecessary surgery and lifelong medication for many.⁷

Overdiagnosis is a challenge to the sustainability of human health and health systems. Its causes—including the best of intentions—are as complex and multifaceted as the potential solutions.⁸⁻¹³ As part of the preparation for a possible national action plan in Australia, we searched the literature for causes of and responses to overdiagnosis. Here we provide the first comprehensive analysis of the possible drivers of overdiagnosis and related overuse, mapped to potential solutions.

Searching the literature

Our approach to the initial and updated PubMed searches of the literature is outlined in the data supplement (see bmj.com) and yielded a total of 36 articles, to which we added a further five (fig 1⇓). We included articles that explicitly discussed possible drivers and potential responses or solutions to the problem of overdiagnosis. We included original research as well as opinion, commentary, and analysis articles. Reflecting the relatively recent and growing interest in overdiagnosis, the vast majority were published since 2013, with generally increasing numbers each year (rising from three before 2013 to nine in 2016 and four in 2017 up until July).

Given the limitations of the literature to date, we couldn't assess the quality of evidence behind each claim in each article, so this is not a systematic review. The vast majority of included articles are analyses or commentaries, with a small number of important

exceptions, including a systematic review. Although discussion of drivers and solutions in many of the included analysis pieces were informed directly by empirical evidence, including systematic reviews the original studies are not included here, as they did not explicitly discuss drivers or solutions in ways captured by our search strategy. The body of empirical evidence highlighting the problem of overdiagnosis is growing, but a systematic review is beyond the scope of this article. Similarly, although analysing the complex inter-relation between overdiagnosis and the overuse it drives¹⁴ is vitally important, it is outside the reach of this article.

The map arising from our analysis is broad but not definitive—potential causes or solutions might not yet have been identified in the literature, and breadth might come at the cost of depth. In addition, our search was based in medicine, and a wider analysis might identify important sociological investigations of medicalisation¹⁵ resulting in different conceptions of the problem, drivers, and solutions. Importantly no strict or established criteria for what defines a driver or a solution exist, so our decisions about inclusion and mapping are open to discussion. Moreover, individual items could in some cases map to more than one domain. We have made no attempt to integrate this map with the extensive science of behaviour change¹⁶ or to specify potential actors to work on solutions, but hope that others will pursue this work.

What's driving overdiagnosis?

Possible drivers of overdiagnosis span five domains: culture, the health system, industry, professionals, and patients and the public (table 1⇓; fig 2⇓). In this section, we offer a narrative summary of some important drivers that appear most commonly in the literature.

Key messages

- Interest is growing in tackling the problems of overdiagnosis and overtreatment
- Possible drivers and potential solutions arise across five inter-related domains: culture, the health system, industry and technology, healthcare professionals, and patients and the public
- More work is needed to develop and evaluate interventions aimed at preventing overdiagnosis
- Raising public awareness of overdiagnosis is a priority

Culture drivers

Popular deep seated beliefs that in healthcare “more is better” and “new is better” are often cited as drivers of unnecessary testing and overdiagnosis.³⁻²⁴ Related to this is a strong collective faith in the benefits of screening the healthy and making an early diagnosis, arising in part from our fears of a serious disease being missed or a diagnosis made too late.³⁻²⁹ As Welch, Schwartz, and Woloshin argue in their 2011 book *Overdiagnosed*, which draws on a wealth of empirical evidence, “early diagnosis is a double edged sword,” with the potential to help but also hidden danger: “the detection of abnormalities that are not destined to ever bother us.”³² Fears of uncertainty, ageing, death, and disease also collectively contribute to this culture of too much medicine.⁸⁻⁴⁷

Health system drivers

Expanding disease definitions, which identify more previously healthy people as “sick,” are commonly cited as a driver of overdiagnosis.³⁻⁴⁷ Health professionals and hospitals frequently have financial incentives to perform more investigations or treatments for their patients, favouring increased and sometimes unnecessary care. Moreover, a system based on fee for service may lead to time restraints during consultations with inadequate time available for shared decision making or the complex explanation of the counterintuitive problem of overdiagnosis.⁴⁻³² As Malvinder Parmar said, “the current fee-for-service system does not compensate for a comprehensive explanation to the patient why a test is not required.”³⁹ Current quality measures in health systems may lack emphasis on preventing overdiagnosis or overuse and instead may indirectly promote these problems.³⁻²¹

Industry and technology drivers

The most important driver in this domain is the use and promotion (to clinicians and the public) of increasingly sensitive tests, leading to detection—often incidentally—of minor “abnormalities,” which may be of uncertain clinical significance and can cause overdiagnosis.³⁻⁴⁷ Industry promotion can also include the funding of patient and advocacy groups.⁸⁻²⁸ As Eric Coon and colleagues point out in their well reasoned and evidence based exploration of potential drivers of overdiagnosis among children, “Advertisements capitalize on our fear of undiagnosed disease and urge us to see our doctor for testing . . . Once considered unbiased, third party advocacy groups are often used to deliver the same message.”¹⁷ Commercial imperatives and conflicts of interest, including financial or reputational conflicts of interests of those involved in guideline panels that expand disease definition, are also cited as a concern.³⁻⁴⁷

Professional drivers

Many authors argue that health professionals are driven to practise defensive medicine owing to their fear of litigation arising from a purported omission.³⁻⁴¹ Closely related is the doctor’s fear of missing a diagnosis, also commonly cited as a

potential driver of overdiagnosis.³⁻²⁹ Health professionals’ unease with dealing with an uncertain diagnosis may lead them towards overtesting and overdiagnosis. This lack of professional confidence and knowledge of harms,⁸⁻³⁰ as well as the tendency to routinely diagnose or “do something” may arise from flaws in medical training,⁹⁻²³ with underemphasis on patient preferences and overemphasis on diagnosis.²⁰⁻⁴⁷

Patient and public drivers

While important, the results of our analysis indicate that this domain has received less attention in the literature, although it clearly overlaps with the culture domain. A number of authors point to a perception that many people have a lack of knowledge about the limits to, and harms of, medicine⁸⁻⁴⁷ and suggest that patients tend to over-rely on tests, including as a means of reassurance.⁸⁻²⁸ Others identify patient expectations that clinicians will “do something” as a potential driver.⁹⁻²¹

What are the potential solutions?

Many of the potential solutions commonly identified in the literature map closely to explicit drivers, with some important exceptions (table 1). For some drivers, such as the increasing complexity and fragmentation of care, specific relevant solutions were not identified. Other drivers showed considerable overlap, both within and across domains. The health system domain, for example, overlaps with the industry and technology domain, where enhanced government regulation of commercial promotion or health technology evaluation clearly falls primarily to policy makers. We made every attempt to link drivers to potential solutions.

Culture solutions

Public awareness and education campaigns are needed to challenge beliefs that in healthcare “more is better”⁹⁻²⁶ and to promote a more healthy scepticism about the benefits and potential harms of early diagnosis.²⁻²² Arguably, initiatives like *The BMJ’s Too Much Medicine* campaign,⁴⁸ Choosing Wisely,⁴⁹ *JAMA Internal Medicine’s* “Less is more,”⁵⁰ and *Health News Review*⁵¹ are moving in this direction. Given the powerful role that media can play in shaping public beliefs, strategies to improve media reporting on overdiagnosis are needed.²⁻²⁸

Health system solutions

Reforming incentives for professionals and healthcare organisations to reward the quality rather than quantity of care is commonly cited as a key way to tackle the problem of too much medicine.³⁻³⁰ Some authors also cite the need for new evidence informed frameworks to be used when disease definitions are changed,⁵² with calls for changes to disease terminology and new expert panels that are more widely representative and have reduced or minimal conflicts of interests.³⁻³⁷ An influential group convened by the US National Cancer Institute is among those advocating changes to disease terminology for indolent lesions.²⁶ Quality indicators and guidelines are also targeted for reform, to tackle any incentives

for medical excess, as well as include new measures of overdiagnosis and overuse.⁴⁻³⁸ More targeted screening programmes that might, for example, limit some screening to well defined high risk populations^{9,39} and mandated strategies to inform patients of the benefits and harms of screening³⁻³⁶ are among potential solutions for minimising the risks of overdiagnosis associated with screening. The 2016 systematic review of studies aimed at reducing low value care and underuse across different parts of the health system found that interventions using multiple strategies and targeting the roles of both clinicians and consumers had the greatest potential.¹⁰

Industry and technology solutions

More rigorous evaluation of the effects of both new and existing diagnostic technology on health outcomes is commonly recommended³⁻³⁹ as a key solution to the problem of increasingly sensitive tests that detect “abnormalities” of uncertain clinical significance. Drawing from the field of ecological economics to frame overdiagnosis as overconsumption, Hensher and colleagues call for “a more rigorous application of the precautionary principle” in technology assessment to avoid giving “potentially harmful overuse the benefit of the doubt.”¹² Other potential solutions include stronger regulation of the advertising of new tests and treatments to the public and health professionals³⁰ and paying greater attention to managing and reducing conflicts of interest with industry.⁴⁷

Professional solutions

The need to tackle the medicolegal concerns regarding missing or delaying a diagnosis was one of the key solutions discussed in the literature.³⁻⁴⁷ Another recommended solution is updating current medical curriculums and continuing medical education to include overdiagnosis and overuse, for both students and practitioners.³⁻⁴⁷ As future practitioners, students must be taught to “look always for the possibility of harm alongside that of benefit.”²⁵ This is vital because, according to Eric Coon and colleagues, “if physicians are not aware of the potential harms of overdiagnosis, patients and families cannot be expected to appreciate them either.”¹⁷

Patients and the public solutions

Widespread awareness campaigns to inform and educate patients and the public on harms as well as benefits of screening and treatment options are commonly cited as essential to tackling overdiagnosis,³⁻⁴⁷ echoing and overlapping with solutions we have classified in the cultural domain. Another frequently recommended solution was promoting shared decision making as a response to several key drivers in this domain.¹³⁻⁴⁷ In addition, several authors proposed the need for prioritising treatment options such as watchful waiting or active surveillance, where appropriate.³⁻³⁵

Where to from here?

We have attempted to retrieve, analyse, and summarise the existing literature on drivers and responses to overdiagnosis and related overuse. Many authors have identified a wide range of potential solutions across five inter-related domains of culture, the health system, industry and technology, professionals, and patients, with the ultimate shared aims to prevent harm, reduce waste, and redirect resources to treating and preventing unmet need in healthcare. The results of this analysis emphasise the need for more evidence about the problem, increased evaluation of potential solutions, and enhanced education across all sectors,

to help wind back the harms of too much medicine effectively, safely, and fairly.

As part of multiple level strategies, in our view the most urgent need is to generate accessible evidence based information and educational materials about overdiagnosis for the public, professionals, and decision makers—both general information and condition specific. Tackling the gamut of financial incentives that drive unnecessary diagnoses and strengthening regulatory processes to enhance evaluation of new and existing diagnostic technology are two more solutions, as difficult as they are desirable. Reforming inappropriately widened disease definitions is arguably the most challenging but most important solution.

We strongly encourage critical responses to this analysis, offering alternative interpretations or missed drivers or solutions. Indeed optimising societies’ responses to medical excess will require broader thinking and analysis outside medicine—from such places as economics and sociology.

As the evidence base around this problem continues to grow, so do attempts to translate that evidence into action. In Norway, a position paper from the College of General Practice calls for action⁵³; in Australia, major influential professional and consumer organisations are launching a national plan; and in Canada, the Quebec Medical Association is already implementing a province-wide strategy. Building on existing initiatives, we hope this analysis will help offer a suite of possible solutions to those seeking to reduce iatrogenic harm and enhance health system sustainability.

Contributors and sources: TP is a medical doctor and PhD candidate, undertaking her doctorate about overdiagnosis of prostate cancer at the Centre for Research in Evidence-Based Practice (CREBP) at Bond University’s Faculty of Health Sciences and Medicine, Australia. JC is a senior research information specialist with extensive experience working on systematic searches including for systematic reviews, based at CREBP. JC is also information specialist with the acute respiratory infections group within the Cochrane Collaboration. RM is senior research fellow, and National Health and Medical Research Council early career fellow, based at CREBP, researching overdiagnosis. RM has a background in medical reporting, including as a columnist with *The BMJ*, and gained his PhD titled Preventing Overdiagnosis in 2015. The sources of information for the paper, and the accompanying tables, are all listed as references and the methods used are explicitly listed in the supplementary file. RM is guarantor for the paper.

Competing interests: RM is co-chair of the scientific committee for the Preventing Overdiagnosis conference, which is supported by the BMJ. TP and JC declare no competing interests.

Provenance and peer review: Commissioned; externally peer reviewed.

- Hoffman JR, Cooper RJ. Overdiagnosis of disease: a modern epidemic. *Arch Intern Med* 2012;358:1123-4. doi:10.1001/archinternmed.2012.3319 pmid:22733387.
- Welch HG, Schwartz L, Woloshin S. *Overdiagnosed: Making People Sick in the Pursuit of Health*. Beacon Press, 2011.
- Moynihan R, Doust J, Henry D. Preventing overdiagnosis: how to stop harming the healthy. *BMJ* 2012;358:e3502. doi:10.1136/bmj.e3502 pmid:22645185.
- Hicks LK. Reframing overuse in health care: time to focus on the harms. *J Oncol Pract* 2015;358:168-70. doi:10.1200/JOP.2015.004283 pmid:25804988.
- Berwick DM, Hackbarth AD. Eliminating waste in US health care. *JAMA* 2012;358:1513-6. doi:10.1001/jama.2012.362 pmid:22419800.
- Bhatt JR, Klotz L. Overtreatment in cancer—is it a problem? *Expert Opin Pharmacother* 2016;358:1-5. doi:10.1517/14656566.2016.1115481 pmid:26789721.
- Vaccarella S, Franceschi S, Bray F, Wild CP, Plummer M, Dal Maso L. Worldwide thyroid-cancer epidemic? the increasing impact of overdiagnosis. *N Engl J Med* 2016;358:614-7. doi:10.1056/NEJMp1604412 pmid:27532827.
- Saini V, Garcia-Armesto S, Klempner D, et al. Drivers of poor medical care. *Lancet* 2017;358:178-90. doi:10.1016/S0140-6736(16)30947-3 pmid:28077235.
- Parmar MS. A systematic evaluation of factors contributing to overdiagnosis and overtreatment. *South Med J* 2016;358:272-6. doi:10.14423/SMJ.0000000000000409 pmid:27043814.

- 10 Colla CH, Mainor AJ, Hargreaves C, Sequist T, Morden N. Interventions aimed at reducing use of low-value health services: a systematic review. *Med Care Res Rev* 2016;1077558716656970. doi:10.1177/1077558716656970. PMID:27402662.
- 11 Chiolerio A, Paccaud F, Aujesky D, Santschi V, Rodondi N. How to prevent overdiagnosis. *Swiss Med Wkly* 2015;358:w14060. PMID:25612105.
- 12 Hensher M, Tisdell J, Zimitat C. "Too much medicine": Insights and explanations from economic theory and research. *Soc Sci Med* 2017;358:77-84. doi:10.1016/j.socscimed.2017.01.020 PMID:28131024.
- 13 Québec Medical Association. Overdiagnosis: Findings and Action plan. Québec; 2014.
- 14 Brownlee S, Chalkidou K, Doust J, et al. Evidence for overuse of medical services around the world. *Lancet* 2017;358:156-68. doi:10.1016/S0140-6736(16)32585-5 PMID:28077234.
- 15 Conrad P. The shifting engines of medicalization. *J Health Soc Behav* 2005;358:3-14. doi:10.1177/002214650504600102 PMID:15869117.
- 16 National Institute for Health and Care Excellence (NICE). Behaviour change: general approaches: Public health guideline 2007 [Available from: <https://www.nice.org.uk/Guidance/PH6>].
- 17 Coon ER, Quinonez RA, Moyer VA, Schroeder AR. Overdiagnosis: how our compulsion for diagnosis may be harming children. *Pediatrics* 2014;358:1013-23. doi:10.1542/peds.2014-1778 PMID:25287462.
- 18 Boddington J. Powerful societal discourses and vested interests need to be tackled by too much medicine campaign. *BMJ* 2013;358:f1724. doi:10.1136/bmj.f1724 PMID:23512462.
- 19 Hofmann B. Medicalization and overdiagnosis: different but alike. *Med Health Care Philos* 2016;358:253-64. doi:10.1007/s11019-016-9693-6 PMID:26912187.
- 20 Hofmann B. Diagnosing overdiagnosis: conceptual challenges and suggested solutions. *Eur J Epidemiol* 2014;358:599-604. doi:10.1007/s10654-014-9920-5 PMID:24880635.
- 21 Malhotra A, Maughan D, Ansell J, et al. Choosing Wisely in the UK: the Academy of Medical Royal Colleges' initiative to reduce the harms of too much medicine. *BMJ* 2015;358:h2308. doi:10.1136/bmj.h2308 PMID:25985331.
- 22 Hoffman JR, Kanzaia HK. Intolerance of error and culture of blame drive medical excess. *BMJ* 2014;358:g5702. doi:10.1136/bmj.g5702 PMID:25315302.
- 23 Morgan DJ, Brownlee S, Leppin AL, et al. Setting a research agenda for medical overuse. *BMJ* 2015;358:h4534. doi:10.1136/bmj.h4534 PMID:26306661.
- 24 Hofmann BM. Too much technology. *BMJ* 2015;358:h705. doi:10.1136/bmj.h705 PMID:25687230.
- 25 Heath I. Overdiagnosis: when good intentions meet vested interests—an essay by Iona Heath. *BMJ* 2013;358:f6361. doi:10.1136/bmj.f6361 PMID:24162944.
- 26 Esserman LJ, Thompson IM Jr, Reid B. Overdiagnosis and overtreatment in cancer: an opportunity for improvement. *JAMA* 2013;358:797-8. doi:10.1001/jama.2013.108415 PMID:23896967.
- 27 Van den Bruel A. The triumph of medicine: how overdiagnosis is turning healthy people into patients. *Fam Pract* 2015;358:127-8. doi:10.1093/fampra/cmv008 PMID:25805809.
- 28 McCaffery KJ, Jansen J, Scherer LD, et al. Walking the tightrope: communicating overdiagnosis in modern healthcare. *BMJ* 2016;358:i348. doi:10.1136/bmj.i348 PMID:26850726.
- 29 Heath I. Role of fear in overdiagnosis and overtreatment—an essay by Iona Heath. *BMJ* 2014;358:g6123. doi:10.1136/bmj.g6123 PMID:25954986.
- 30 Carpenter CR, Raja AS, Brown MD. Overtesting and the downstream consequences of overtreatment: implications of "preventing overdiagnosis" for emergency medicine. *Acad Emerg Med* 2015;358:1484-92. doi:10.1111/acem.12820 PMID:26568269.
- 31 Moynihan R, Smith R. Too much medicine? *BMJ* 2002;358:859-60. doi:10.1136/bmj.324.7342.859 PMID:11950716.
- 32 Treadwell J, McCartney M. Overdiagnosis and overtreatment: generalists—it's time for a grassroots revolution. *Br J Gen Pract* 2016;358:116-7. doi:10.3399/bjgp16X683881 PMID:26917633.
- 33 Glasziou P, Moynihan R, Richards T, Godlee F. Too much medicine; too little care. *BMJ* 2013;358:14247. doi:10.1136/bmj.14247 PMID:23820022.
- 34 Moynihan R, Henry D, Moons KGM. Using evidence to combat overdiagnosis and overtreatment: evaluating treatments, tests, and disease definitions in the time of too much. *PLoS Med* 2014;358:e1001655. doi:10.1371/journal.pmed.1001655 PMID:24983872.
- 35 Esserman LJ, Thompson IM, Reid B, et al. Addressing overdiagnosis and overtreatment in cancer: a prescription for change. *Lancet Oncol* 2014;358:e234-42. doi:10.1016/S1470-2045(13)70598-9 PMID:24807866.
- 36 Parker LM, Rychetnik L, Carter S. Framing overdiagnosis in breast screening: a qualitative study with Australian experts. *BMC Cancer* 2015;358:606. doi:10.1186/s12885-015-1603-4 PMID:26314748.
- 37 Kamerow D. How to decrease overtreatment in cancer. *BMJ* 2013;358:f5071. doi:10.1136/bmj.f5071 PMID:23945466.
- 38 Morden NE, Colla CH, Sequist TD, Rosenthal MB. Choosing wisely—the politics and economics of labeling low-value services. *N Engl J Med* 2014;358:589-92. doi:10.1056/NEJMp1314965 PMID:24450859.
- 39 Mortani Barbosa EJ Jr. Overdiagnosis in lung cancer screening can be reduced to a low, manageable level via a multilayered strategy involving perfecting reporting systems, restricting screening to high-risk groups, developing better risk stratification models, and improving management algorithms. *Acad Radiol* 2016;358:113-5. doi:10.1016/j.acra.2015.09.006 PMID:26514433.
- 40 Cervera Deval J, Sentis Crivillé M, Zulueta JJ. [Overdiagnosis in cancer screening]. *Radiologia* 2015;358:188-92. doi:10.1016/j.rx.2014.06.007 PMID:25174786.
- 41 Samadian S, Farhat A. Problem based learning, litigation, and EWTD contribute to too much medicine. *BMJ* 2013;358:f4790. doi:10.1136/bmj.f4790 PMID:23900824.
- 42 Moynihan R, Heneghan C, Godlee F. Too much medicine: from evidence to action. *BMJ* 2013;358:f7141. doi:10.1136/bmj.f7141 PMID:24307723.
- 43 Nicholson BD. Detecting cancer in primary care: Where does early diagnosis stop and overdiagnosis begin? *Eur J Cancer Care* 2017;26:e12692.39.
- 44 Jansen J, Naganathan V, Carter SM, et al. Too much medicine in older people? Deprescribing through shared decision making. *BMJ* 2016;358:i2893. doi:10.1136/bmj.i2893 PMID:27260319.
- 45 Brindle LA. GP-patient communication about possible cancer in primary care: Re-evaluating GP as gatekeeper. *Eur J Cancer Care (Engl)* 2017;358:e12699. doi:10.1111/ecc.12699 PMID:28489299.
- 46 Hodgson H. Choosing wisely—choosing tactics. *Clin Med (Lond)* 2015;358:315-6. PMID:26407376.
- 47 Hanslik T, Flahault A. [Overmedicalization: When too much medicine harms]. *Rev Med Intern* 2016;358:201-5. doi:10.1016/j.revmed.2015.10.009 PMID:26586148.
- 48 Too Much Medicine. BMJ campaign. <http://www.bmj.com/too-much-medicine>.
- 49 Choosing Wisely 2017 <http://www.choosingwisely.org/>.
- 50 Grady D, Redberg RF. Less is more: how less health care can result in better health. *Arch Intern Med* 2010;358:749-50. doi:10.1001/archinternmed.2010.90 PMID:20458080.
- 51 Schwitzer G. Pollution of health news. *BMJ* 2017;358:j1262. doi:10.1136/bmj.j1262 PMID:28298320.
- 52 Doust J, Vandvik PO, Qaseem A, et al. Guidelines International Network (G-I-N) Preventing Overdiagnosis Working Group. Guidance for modifying the definition of diseases - a checklist. 2017. *JAMA Intern Med* 2017;358:1020-5. doi:10.1001/jamainternmed.2017.1302 PMID:28505266.
- 53 Brelvi P. Position paper. Overdiagnosis and related medical excess: The Norwegian College of General Practice, The Norwegian Medical Association; 2016 <http://legeforeningen.no/Fagmed/Norsk-forening-for-allmennmedisin/Styret/Sentrale-fagpolitiske-dokumenter/2016-Position-paper-on-overdiagnosis-and-medical-excess/>

Published by the BMJ Publishing Group Limited. For permission to use (where not already granted under a licence) please go to <http://group.bmj.com/group/rights-licensing/permissions>

Table

Table 1 | Possible drivers of and potential solutions to overdiagnosis

Possible drivers	Potential solutions
Culture	
Beliefs: "More is better," "new is better," "early is better," "wants to know/screen"—even if currently healthy, imperative of possibility (we have to test because we can) ^{3-17,24}	Public awareness, information, and education campaigns challenging beliefs ^{9-24,26}
Faith in early diagnosis or detection and prevention is better than cure, fear of diagnosis missed or made too late ^{3-27,29}	More healthy scepticism on the benefits of early diagnosis especially in populations less likely to benefit ^{2,22}
Intolerance of uncertainty and error ^{9-28,30}	Address intolerance of uncertainty ²²
Medicalisation of life; fear of ageing, sickness, or death ^{9-22,25-31}	NA
Biased reporting by media—eg, reporting lack of care more than underdiagnosis and overdiagnosis ⁸⁻²³	Improve media reporting, journalist training, campaigns about overdiagnosis and overuse ^{2,28}
Health system	
Financial incentives for more tests, treatments, and diagnoses, including fee for service and not enough time to talk ^{4,8,11-21,23-32}	Reform incentives from quantity to quality ³⁻³⁰
Unique market system within healthcare; third party payer can shield patient from cost ^{4,23}	
Corporate health insurance packages for healthy adults ¹⁸	
Supplier induced demand ¹²	
Expanding disease definitions ³⁻³³	Reform process of disease definition and constitution of guideline panels—fewer conflicts of interest, more broadly representative ³⁻³⁴ Change terminology for indolent lesions—eg, labelling innocuous cancers as "indolent lesion of epithelial origin" rather than cancer ^{11-35,37} Avoid or rename certain disease labels ²⁷
Quality measures that encourage doing more ³⁻²¹	Reform performance and quality measures and develop new measures of overdiagnosis and overuse ⁴⁻³⁸
Lack of focus of quality and performance indicators on overuse and overmedicalisation ¹⁷	
Guidelines⁴⁻³²	Reform guidelines on tests and treatments: Include what's not recommended ⁴⁻²³ Review and include harms and risk of overdiagnosis ⁶⁻³⁰ Avoid conflict of interests in guideline panels, use evidence ^{11,13}
Screening³⁻³³	Reform or limit screening: Avoid certain tests or screening tests ^{11,35} Less frequent screening tests ¹¹⁻³⁵ Targeted screening tests (eg, limit screening to well defined high risk population) ^{9-35,37-40} Inform patients of benefits and harms of screening ³⁻³⁶ Raise threshold for "positive" screen or recall and biopsy ²⁶⁻³⁹ Change radiological advice ³⁶
Increasing complexity† and fragmentation‡ of care and loss of continuity of care ⁴⁻⁴¹	NA
NA	More research on: drivers of overdiagnosis and overuse ^{9,23} overdiagnosis and overuse frequency and best methods to quantify the problem ¹⁷⁻⁴² undiagnosing and deprescribing ³ deliberative and qualitative methods (including community juries) to inform policy and other decisions ²³⁻³⁶ evaluating responses to overuse, including communication ²³
NA	Multicomponent interventions tackling patient and provider roles, to reduce overuse and low value care ¹⁰
Industry§ and technology	

Table 1 (continued)

Possible drivers	Potential solutions
Industry promotion of tests and treatments to professionals and public: Advertising (including direct to consumer marketing), promotion, influence using the media to generate demand for diagnosis ³⁻²⁹ Industry funding of patient and advocacy groups ⁸⁻²⁸ Industry wants to develop better diagnostic technology—technological arms race ¹⁹⁻²⁴	Better regulate advertising ³⁰
Diagnostic tests with increased sensitivity: New technology captures ever smaller “abnormalities” or incidental findings ³⁻³³ Changes nature and prevalence of disease ²⁰⁻²⁴ Feedback loop: diagnosing milder cases, high prevalence, better outcomes, more tests and treatments—illusion of progress ³⁻²⁵	Better evaluation of diagnostic technology: Better evaluation of new and existing diagnostic technology ²³⁻⁴³ Evaluate effects of test-treatment strategies on outcomes, not just test performance alone, more caution in treating incidentalomas ³⁻³⁴ Better identify harmful v benign abnormalities ³⁻²⁷ Increasing specificity of tests at the cost of slightly decreased sensitivity ³⁹ Include patient voice in development, assessment, implementation, and use of technology ²⁴ Strengthen health technology assessments and cost effectiveness analyses ¹²
Medicine as a business and conflicts of interest Commercial and corporate clinics—self referral, vertical integration ³⁻⁸ Commercial and other conflicts of interest ³⁻²⁷ Professional conflicts of interest, specialist societies protecting or expanding turf ³⁻⁹ Toxic combination of vested interests and good intentions ²⁵ Supply-sensitive care ¹⁷	Declare, reduce, exclude conflicts of interest ¹⁹
Industry interest in expanding markets, including financial ties to those setting diagnostic thresholds ³⁻³²	Better evaluation of changes to (expansion of) disease definitions ³⁴
Professionals	
Fear of litigation or defensive medicine ^{3-19, 23, 27, 30-41}	Tackle and reform drivers of litigation and defensive medicine ³⁻²²
Fear of missing disease or decision regret ³⁻²⁹	Critical thinking and comfort with uncertainty (eg, with a period of active surveillance, watchful waiting) ¹²⁻³⁵
Flaws in training: Hard to do nothing rather than something ⁹⁻²³ Overemphasis on diagnosis ²⁰ Problem based learning approach leads to shotgun approach to diagnosis, rewards unusual or rare diagnoses, and contributes to overtesting and overdiagnosis, with an emphasis on avoiding omission errors ⁴⁻⁴¹ Poor knowledge of patient preference and shared decision making ²³ Improper weighting of absolute v relative risk ⁸⁻⁴⁴ Lack of professional self confidence or knowledge of harms ⁸⁻³⁰	Educational curricula and information materials on overdiagnosis and overuse for students and professionals ³⁻⁴⁵
NA	Interventions to reduce overuse and overdiagnosis: Multicomponent interventions tackling patient and provider roles (eg, clinician education with decision support or feedback) to reduce overuse and low value care ¹⁰ Using minimally invasive techniques to eventually replace surgery for low risk tumours (eg, low staged lung tumours) ³⁹ Clinical decision support and performance feedback ¹⁰⁻¹³ Compare practices with peers (eg, prescribing, referrals, tests) ¹³ Deprescribing ⁴⁴
Overusing or over-reliance on tests ⁹⁻¹³ : Using tests to close consultation ¹³ Chasing incidental findings and test outcomes of no or doubtful significance ⁹ Chart overload syndrome (order test to avoid reading thick chart) ⁹ Specialists ordering tests before clinical appointment assessments ⁹ Specialists suggesting unnecessary tests to primary provider, especially during short consultation or in corridor ⁹ Incomplete information of test ordering requests ⁹	Reduce over-reliance on tests Anticipate consequences of discovering abnormalities ¹¹ More time with patient to order less tests ³⁰⁻⁴⁶ Tools to promote discussion with patients on why test might not be needed ¹³ Don't use diagnostic labels unless needed ³³ Consider stepped or slower diagnosis ¹⁷⁻³³

Table 1 (continued)

Possible drivers	Potential solutions
Radiologists suggesting further imaging studies (self referral) when they don't have complete clinical information about patient ⁹	
Tests express "care" to patients and families, whereas watchful waiting deemed uncaring ¹⁷	
"Better to know" bias that may not be warranted ⁸	
Too much focus on surrogate markers ²⁰	
Patients and public	
Over-reliance on tests, including for reassurance ^{8,28}	NA
NA	Shared decision making ^{13-21,23-46}
Lack of knowledge or confidence about limits and harms of medicine ^{9,23}	Information and education campaigns, with accessible evidence, on the benefits and harms of interventions, including screening, from unconflicted sources ^{3,8,10-46}
Expectation that clinicians will "do something" ^{19,21}	Research and promote doing nothing, watchful waiting, or active surveillance as legitimate options, where appropriate ³⁻³⁵

Some responses may map to more than one driver. NA=not applicable. *In most markets, costs are constrained by consumers' ability to pay; in health care, services are ordered by physicians, who are unaffected by cost, and received by patients, who often bear only a portion of the cost. [†]Leading to test duplication, polypharmacy, and redundant opinions owing to one person seeking advice from several physicians. [‡]When patients are forced or indirectly encouraged to seek care through an inappropriate or inefficient avenue. For instance, all too frequently, the most expeditious route to urgent, but not emergent, care is through an emergency department. §Industry includes for-profit drug and device companies and commercial or corporate medicine providers. ¶Where higher capacity in the form of hospital beds and imaging modalities drives medical use and inevitably uncovers patient abnormalities

Figures

Fig 1 Flow diagram

Fig 2 Overdiagnosis and related overuse. Mapping possible drivers to potential solutions. COI=conflict of interest; OD=overdiagnosis; OU=overuse.