

Quantification of Competitive Game Demands of NCAA Division I College Football Players Using Global Positioning Systems

Wellman, Aaron D.; Coad, Sam C.; Goulet, Grant C.; McLellan, Christopher P.

Published in:
Journal of Strength and Conditioning Research

DOI:
[10.1519/JSC.0000000000001206](https://doi.org/10.1519/JSC.0000000000001206)

Licence:
Other

[Link to output in Bond University research repository.](#)

Recommended citation(APA):
Wellman, A. D., Coad, S. C., Goulet, G. C., & McLellan, C. P. (2016). Quantification of Competitive Game Demands of NCAA Division I College Football Players Using Global Positioning Systems. *Journal of Strength and Conditioning Research*, 30(1), 11-19. <https://doi.org/10.1519/JSC.0000000000001206>

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

For more information, or if you believe that this document breaches copyright, please contact the Bond University research repository coordinator.

QUANTIFICATION OF COMPETITIVE GAME DEMANDS OF NCAA DIVISION I COLLEGE FOOTBALL PLAYERS USING GLOBAL POSITIONING SYSTEMS

Aaron D. Wellman¹, Sam C. Coad¹, Grant C. Goulet², Christopher P. McLellan¹.

¹ Faculty of Health Sciences and Medicine, Bond University, Queensland, Australia.

² University of Michigan, School of Kinesiology, Ann Arbor, MI.

ABSTRACT

The aim of the present study was to examine the competitive physiological movement demands of NCAA Division I college football players using portable global positioning system (GPS) technology during games, and to examine positional groups within offensive and defensive teams, to determine if a player's physiological requirements during games are influenced by playing position. Thirty-three National Collegiate Athletic Association (NCAA) Division I Football Bowl Subdivision football players were monitored using GPS receivers with integrated accelerometers (GPSports, Canberra, Australia) during 12 regular season games throughout the 2014 season. Individual datasets (n = 295) from players were divided into offensive and defensive teams, and

subsequent position groups. Movement profile characteristics including total, low-, moderate-, high-intensity and sprint running distances (m), sprint counts, and acceleration and deceleration efforts, were assessed during games. A one-way ANOVA and post-hoc Bonferroni statistical analysis were used to determine differences in movement profiles between each position group within offensive and defensive teams. For both offensive and defensive teams, significant ($p < 0.05$) differences exist between positional groups for game physical performance requirements. The results of the present study identified that wide receivers (WR) and defensive backs (DB) completed significantly ($p < 0.05$) greater total distance, high-intensity running, sprint distance, and high-intensity acceleration and deceleration efforts compared to their respective offensive and defensive positional groups. Data from the present study provide novel quantification of position specific physical demands of college football games and support the use of position-specific training in the preparation of NCAA Division I college football players for competition.

Key Words: GPS, monitoring, American football

INTRODUCTION

American football is a field-based team sport requiring high levels of muscular strength, power, speed and agility, and is characterized by intense collisions and repeated high-intensity movements (27). American football games are intermittent in nature involving short-duration high-intensity bouts of exercise which incorporate movements such as

sprinting, backpedaling, accelerating, decelerating, and physical collisions, separated by transient periods of low-intensity recovery between plays (12). During the in-season period of competition, players competing in NCAA Division I college football are required to participate in twelve regular season games on a consecutive weekly basis. Few studies have investigated (12,26) the demands of NCAA Division I football games and as such, the movement characteristics of competition in college football players remain ambiguous. While research (12,26) has provided a rudimentary description of exercise to rest ratios encountered during NCAA Division I college football games, a more detailed assessment of position-specific movement demands during competition provides novel insight to improve our understanding of the demands of competition and enable increased scope for position-specific training and conditioning programs to optimize on-field performance.

The development of global positioning system (GPS) technology with integrated tri-axial accelerometers have allowed the physiological demands of training and competition in contact team sport to be quantified by tracking the movement of players (2,10,32). Improvements in GPS technology have subsequently resulted in enhancements in accuracy (13), and the validity and reliability of GPS to determine the movement demands of team sports is well established (6,14,15,30). The quantification of team-sport competition demands using GPS technology has been reported in sports similar in nature to American football, including rugby league (2,10,24), rugby sevens (11), Australian football league (AFL) (18,29,31), and rugby union (7,21). Further substantiating the use of GPS technology to accurately determine position-specific

demands of team sport, Boyd et. al. (4) demonstrated the capacity of GPS units with integrated accelerometry to differentiate between training drills and competitive games, and discriminate between players competing in elite and sub-elite team-sport competitions. Although GPS technology is widely used in team sports for analysis of game and training movement demands, current literature on the movement profile characteristics of American football players is limited (8).

DeMartini et. al. (8) reported movement profile characteristics associated with pre-season training sessions in NCAA Division I college football by examining the physical demands of Division I college football players during nine pre-season practices over the course of eight days, utilizing GPS to evaluate total distance covered and running velocity characteristics. The main findings reported by DeMartini et. al. (8) were that non-linemen covered greater total distance and sprint distance than linemen, who covered greater distance at slower speeds. To date, ambiguity remains regarding the demands of in-season NCAA Division I college football games and team training activities (8).

In American football each position group has distinct physiologic and biomechanical demands associated with specific technical and tactical requirements (16), however uncertainty exists regarding the position-specific movement demands of NCAA football competition. Given the widespread inclusion of GPS technology in collegiate American football programs, a detailed assessment of competitive movement profile

characteristics will provide sports performance specialists with quantified information on game demands. A more comprehensive understanding of the demands of NCAA football competition will augment our understanding of the position-specific movement demands of NCAA college football players, and allow sport coaches to individualize training programs that replicate the demands of American football games.

The aim of the present study was to 1) examine the competitive physiological movement demands of NCAA Division I college football players using portable global positioning system (GPS) technology during games, and 2) to examine positional groups within offensive and defensive teams, to determine if a player's physiological requirements during games are influenced by playing position. We hypothesized that there will be substantial positional differences in movement demands of NCAA Division I college football players during games. Data obtained will provide scope for performance coaches seeking to optimize position-specific training regimens.

METHODS

EXPERIMENTAL APPROACH TO THE PROBLEM

Portable GPS and integrated tri-axial accelerometry technology was used in the present study to quantify the position-specific movement characteristics of NCAA Division I

college football games. The GPS movement profile data was collected during twelve regular season NCAA Division I college football games. All games were 60-minutes in duration, comprised of four 15-minute quarters, each followed by a brief recovery period, and played outdoors between the hours of 12:00 and 21:00 over a period of twelve to thirteen weeks from September to November. All participants were required to participate in a minimum of 75% of the total offensive or defensive plays for the GPS datasets to be included in the present study. Each individual GPS dataset was characterized as constituting either offensive or defensive team performance, and subsequently divided into specific positional groups for the offense that included wide receivers (WR), quarterbacks (QB), running backs (RB), tight ends (TE), offensive linemen (OL), and for the defense that included defensive backs (DB), linebackers (LB), defensive ends (DE) and defensive tackles (DL).

SUBJECTS

Thirty-three National Collegiate Athletic Association (NCAA) Division I Football Bowl Subdivision (FBS) football players (age 20.7 ± 1.0 years; height 188.6 ± 7.2 cm; and mass 106.7 ± 19.6 kg) participated in the present study. The heights and weights for each position group are expressed as means \pm standard deviation and presented in Table 1. All subjects were collegiate athletes whom had been selected to participate in the football program eight months prior to the commencement of the study. All participants in the present study took part in the teams' off-season physical

development training program that included a full-body strength and power training program and specific skills and conditioning sessions designed to simulate the demands of NCAA Division I college football competition. The present study comprises statistical analysis of data collected as part of the day to day student athlete monitoring and testing procedures within the university's football program. Researchers were provided with de-identified GPS datasets from twelve regular season games for analysis. De-identified data included participant playing position for the purposes of position-specific data analysis. Ethical approval was obtained from the the university's human research ethics committee.

Insert Table 1 Here

PROCEDURES

Global Positioning System Units. The present study used commercially available GPS receivers (SPI HPU, GPSports, Canberra, Australia) which operated in a non-differential mode at a sampling frequency of 15 Hz. The GPS receivers also contain integrated tri-axial accelerometers (IA), which operated at 100 Hz and assessed the frequency and magnitude of full-body acceleration ($\text{m}\cdot\text{second}^{-2}$) in three dimensions, namely, anterior-posterior, mediolateral, and vertical (17,22). Subjects had previously worn GPS receivers in outdoor training sessions that included football-specific running, and skill-related and game-simulated contact activities during a three week pre-season training

period. Prior to the commencement of each game, GPS receivers were placed outside for 15 minutes to acquire a satellite signal, after which, receivers were placed in a custom designed pocket attached to the shoulder pads of the subjects. Shoulder pads were custom-fit for each individual, thereby minimizing movement of the pads during competition. The GPS receivers used in the present study (66 g; 74 mm x 42 mm x 16 mm) were positioned in the center of the upper back, slightly superior to the scapulae. Subjects were outfitted with the same GPS receiver for each of the twelve games. Following the completion of games, GPS receivers were removed from the shoulder pads, and subsequently downloaded to a computer for analysis utilizing commercially available software (Team AMS, GPSports, Canberra, Australia). The validity and reliability of GPS to measure distance and velocity during high-intensity exercise that characterizes contact and noncontact team sports have been reported (3,9,14,25). Johnston et. al. (14) have demonstrated GPS receivers utilized in the present study to be valid for measuring total distance and average peak speed in a team sport simulation circuit, with intraclass correlation values of interunit reliability reported to be 0.94 for high speed running (14.00 – 19.99 km·h⁻¹) distance, 0.81 for very high speed running (> 20.00 km·h⁻¹) distance, - 0.20 for total distance, and - 0.14 for peak speed.

Data provided from GPS receivers were assessed as movement profile variables including total, low-intensity, moderate-intensity, high-intensity and sprint distances (m), max velocity achieved (km/h), and counts of sprint, acceleration and deceleration efforts. Classifications of parameters of movement profile variables are described below and presented in Table 2. Each of the GPS variables measured in the present

study was calculated using commercially available software (Team AMS, GPSports, Canberra, Australia).

Movement Profile Classification. Movement profile classifications have been described for game analysis in similar contact team sports (19,20,23,24), however the classification profile utilized in the present study was devised for American football players. Each movement classification was coded as one of four speeds of locomotion (Table 2). Low-intensity movements, such as standing, walking and light jogging, were considered to be 0 - 10 km·h⁻¹, moderate-intensity movements, such as a cruising jog, were considered to be 10.1 – 16.0 km·h⁻¹, high-intensity movements, such as fast jog or striding, were classified as 16.1 – 23.0 km·h⁻¹, and sprinting or maximal effort movements were classified as exceeding 23.0 km·h⁻¹. Short duration high-intensity movement efforts, or measures of acceleration and deceleration, were classified as three groups, specifically, moderate (1.5 – 2.5 m·s⁻²), high (2.6 – 3.5 m·s⁻²) and maximal (> 3.5 m·s⁻²) and presented as a count of how many efforts an athlete undertook per game.

Insert Table 2 Here

STATISTICAL ANALYSES

All movement and variables from the present study were presented as descriptive statistics, mean \pm standard deviation (SD). Hypothesis testing was conducted to determine any main effects for movement profile data between position groups on the offensive and defensive teams. A one-way ANOVA was used to determine positional group main effects. In the event homogeneity of variance assumption was violated, a Welch Robust Test of Equality was used to determine main effects between position groups. For all main effects detected by a one-way ANOVA, post-hoc Bonferroni tests were utilized. Alpha intervals for all hypothesis testing were set at $p < 0.05$ as the level of significance for statistical tests. All statistical analyses were performed using the Statistical Package for the Social Sciences (SPSS for Windows, version 14.0; SPSS, Inc., Chicago, IL. USA).

RESULTS

Offense: Significant ($p < 0.001$) main effects from ANOVA testing were reported for all movement profile variables measured in the present study for the offensive position groups (Table 3). From post-hoc analysis of movement profile variables, total distance, moderate-intensity distance, high-intensity distance and sprinting distance covered by the WR position was significantly ($p < 0.001$) greater in comparison to all other offensive position groups, including RB, QB, TE, and OL. Low-intensity distance covered by the WR position was also significantly ($p < 0.001$) greater for all offensive position groups apart from QB. The QB position group covered significantly ($p < 0.001$) more low-

intensity distance than RB, TE, and OL positions. Moderate-intensity distances were significantly ($p < 0.05$) greater for RB and QB position groups compared to TE and OL position groups. High-intensity distances were significantly ($p < 0.01$) greater for the RB and TE positions compared to QB and OL positions. Sprinting distances were significantly ($p < 0.001$) greater for RB compared to OL. The average max speed achieved by WR, RB and QB positions was significantly ($p < 0.05$) greater than TE and OL positions, while the average max speed achieved by WR position group was significantly ($p < 0.05$) greater than the RB position group.

For all high-intensity movement profile variables, including sprint efforts, moderate-, high-intensity, maximal-intensity acceleration and deceleration efforts, the WR position was involved in significantly ($p < 0.01$) more efforts than any other offensive position group. The QB and RB positions were involved in significantly ($p < 0.01$) more sprint efforts per game compared to TE and OL positions. The TE and OL groups were involved in significantly ($p < 0.001$) more moderate acceleration efforts than the RB and QB positions; however, the OL position group had significantly ($p < 0.001$) less maximal acceleration efforts compared to QB and RB positions. The OL position was also involved in significantly ($p < 0.001$) more moderate deceleration efforts compared to the RB position, while for maximal deceleration efforts the OL position was involved in significantly ($p < 0.05$) less than the RB and QB position groups.

Insert Table 3 Here

241

242 *Defense:* Significant ($p < 0.001$) main effects from ANOVA testing were reported for all
243 movement profile variables measured in the present study for defensive position groups
244 team (Table 4). Post-hoc analysis of movement profile variables including total distance,
245 moderate-intensity distance, high-intensity distance and sprinting distance covered,
246 revealed that both the DB and LB positions covered significantly ($p < 0.05$) greater
247 distances in all zones than the DE and DT positions during games. The only main effect
248 reported for distance covered between the DB and LB position groups was for low-
249 intensity distance covered, with the DB position covering significantly ($p < 0.05$) more
250 than the LB position group. The DB position had the highest average max speed which
251 was significantly ($p < 0.05$) greater than all other defensive positions. The average max
252 speed of the LB position group was significantly ($p < 0.05$) greater than DE and DT
253 positions, although significantly ($p < 0.05$) less than DB. The DE position average max
254 speed was significantly ($p < 0.05$) greater than the DT position, and significantly ($p <$
255 0.05) less than DB and LB positions.

256

257 The DB position group was involved in significantly ($p < 0.05$) more sprint efforts,
258 moderate-, high-, and maximal-intensity acceleration and deceleration efforts, than the
259 DE and DT positions groups. Apart from moderate acceleration and deceleration efforts
260 and high-intensity deceleration efforts, the DB position group was involved in
261 significantly ($p < 0.05$) more high-intensity movements than the LB position group. The
262 LB position group was involved in significantly ($p < 0.05$) more sprint efforts, high- and
263 maximal-intensity acceleration and deceleration efforts than the DE and DT positions.

Lastly, the DE position group was involved in significantly ($p < 0.05$) more high-intensity acceleration efforts than the DT position group.

Insert Table 4 Here

DISCUSSION

The present study examined the competitive physiological movement demands of NCAA Division I college football players using portable GPS technology during games, and assessed positional groups within offensive and defensive teams, to determine if a player's physiological requirements during games are influenced by playing position.

The results of the present study provide novel insight into the competitive demands experienced by NCAA Division I college football players, and provide scope for the design of position-specific and game-specific physical conditioning strategies for coaches seeking to optimize training for the demands of competition. The results confirm our hypothesis that significant differences in movement profiles accompanying NCAA Division I college football games exist between playing positions. The most notable finding for physical characteristics of games in both offensive and defensive teams were the movement profiles of the WR, DB, and LB positions, with athletes in these three position groups covering more total distance at higher intensities compared to all other positions on their respective offensive and defensive teams.

285

286 The total distance covered by athletes in team-sport competition such as American
287 football, may be considered an overall reflection of running volume. The present study
288 found a significant ($p < 0.001$) difference in total distance traveled between position
289 groups within both the offensive and defensive teams. The WR position group covered
290 more total distance per game than all other offensive groups. Similarly on defense, the
291 DB and LB position groups covered greater total distance than the DT and DE position
292 groups. The finding of the present study that the WR, DB, and LB position groups
293 covered greater total distance, is consistent with the work of DeMartini et. al. (8) that
294 found significant differences in distance traveled between linemen (2573 ± 489 m) and
295 non-lineman (3532 ± 943 m) during pre-season training. However, the present study
296 evaluated game data over the course of twelve games compared to DeMartini et. al. (8)
297 who evaluated data obtained during pre-season training in the heat. The absence of
298 published research in relation to the demands of NCAA Division I football games make
299 comparisons with others problematic. Despite the absence of comparable studies, the
300 present results indicate that the total distance covered for both linemen (3314.0 m) and
301 non-linemen (4141.3 m) during games are greater than those data reported by
302 DeMartini et. al. (8). From an observational perspective, results from the present study
303 are potentially due to the distance away from the line of scrimmage where the WR, DB
304 and LB position groups started plays. Beginning play further from the line of scrimmage
305 gives athletes a larger area for movement, providing an increased movement
306 requirements during plays and further distances to travel between plays to huddle for
307 brief tactical discussion related to subsequent play. Given WR, DB and LB covered

greater total running distance throughout games than their offensive and defensive teammates, it is reasonable to suggest athletes in these positions may require modified running volumes in training to support recovery and adequately prepare them for the physical demands of subsequent competition.

In addition to differences in total distance covered by WR, DB, and LB, the present study found significant ($p < 0.05$) differences in moderate-intensity, high-intensity, and sprint distances covered by WR, DB, and LB compared to all other positions on their respective teams. The RB and TE covered significantly ($p < 0.05$) more high-intensity distance than OL. Similar observations in American football training were made by Demartini et. al. (8) who reported non-linemen covering significantly ($p < 0.001$) more high-intensity ($> 16.0 \text{ km}\cdot\text{h}^{-1}$) distance for position drills, team drills, and total practice time than linemen in pre-season training. Positional differences observed in the present study may be attributed to the position-specific requirements of games. Tactically, the primary responsibility of OL is to block defensive players, preventing opponents from tackling their own team's ball carrier. These movements are associated with short bursts of acceleration, deceleration, and change of direction, which most frequently occur within a few yards of the line of scrimmage, thereby limiting the distance traveled per play. Players in the DT and DE position groups characteristically accelerate short distances and perform rapid change of direction movements before engaging the opposing OL, followed by pursuing the ball carrier. The position-specific requirements of the OL, DT, and DE positions, requiring a static play initiation posture at or near the line of scrimmage at commencement of each play followed by contact with an opponent

positioned approximately one meter apart, likely influences subsequent running distances. These distances are less than that covered by other positions on the offensive and defensive teams that require players to travel greater distances prior to engaging an opponent. The differences in high-intensity distance covered by TE and RB, compared to OL, may be attributed to the more diverse requirements of these position groups, including blocking, running with the ball, and releasing on pass routes. The WR position group is required to repeatedly run passing routes at high velocities throughout the course of games, consequently accounting for significantly greater high-intensity distance and significantly more sprint efforts when compared to all other offensive positions. The DB position group is primarily responsible for defending WR on passing routes, however they also provide secondary support on running plays. As the last line of defense, the DB position is often responsible to make tackles on long running or passing plays, which is indicated in the current study with greater high-intensity distance and more sprint efforts of DB when compared to all other defensive positions.

In addition to the distance covered during play, the WR and DB cover more distance between plays as they are required to jog back to the line of scrimmage at the conclusion of plays, which may be a distance 20-30 m to either huddle or re-assume their alignment for subsequent play, whereas OL, DT, and DE characteristically walk short distances during recovery between plays (26). The LB position is required to defend running plays in addition to covering WR, RB and TE on passing plays which may account for similar movement characteristics to the DB position. The results of the present study highlight the unique movement demands of WR, DB and LB position

groups in comparison to other positions on their respective offensive and defensive teams, and is potentially related to their proximity to the line of scrimmage at the initiation of play. Young et. al. (32) reported greater running distance covered at high speed, along with moderate and high accelerations and decelerations to be associated with markers of muscle damage in collision team-sport players, and consequently, the monitoring and prudent adjustment of weekly training loads of the WR, DB and LB position groups specifically, may reduce the likelihood of subsequent performance decrements associated with fatigue.

Research (1,21,24) in team-sports utilizing portable GPS technology indicate positional differences in movement characteristics during competition. No previous studies have reported the movement demands of NCAA Division I football competition, consequently a lack of understanding exists regarding the demands of American football games. Investigations in team sports similar to American football, including rugby league, rugby union, and Australian rules football, indicate significant differences exist in high-intensity movements including acceleration and deceleration efforts (28,32), and maximal speed (5,24) between position groups. The present study found significant differences in maximal running speeds and maximal acceleration and deceleration efforts recorded from offensive position groups. The average max speed of WR position was significantly ($p < 0.05$) greater than all other offensive positions except QB. The RB and QB position groups average max speed was significantly ($p < 0.05$) greater than that of both the TE and OL position groups. The WR group had significantly ($p < 0.05$) more sprint, maximal acceleration, and maximal deceleration efforts than all other offensive

position groups, presumably do to repeated route running requiring sprinting and frequent changes of direction.

Defensively, there were no significant differences between total, moderate-, or high-intensity distance covered between DB and LB position groups, however, significant ($p < 0.05$) differences were indicated for average max speed, sprint, maximal acceleration, and maximal deceleration efforts. The DB group had significantly ($p < 0.05$) more sprint, maximal acceleration, and maximal deceleration efforts than all other defensive positions, highlighting the specific high-intensity running requirements of this position during defensive play. The LB position group demonstrated significantly ($p < 0.05$) greater average max speeds, sprint, maximal acceleration, and maximal deceleration efforts than the DE and DT groups. Similar research (8) has not quantified high-intensity movement characteristics of individual position groups, making comparisons with the present study difficult.

The significant differences between the DB group when compared to the defense as a whole, and the LB compared to DT and DE, highlight three distinct running profiles for the defensive team, requiring different forms of training to achieve optimal development. The starting positions upon commencement of each play for the DB and LB groups afford larger areas to achieve higher max speeds, while the positional requirements of defending pass routes and pursuing ball carriers result in greater changes of direction for the DB and LB groups. The WR and DB position groups achieved significantly

greater max speeds, sprint efforts, and maximal acceleration and deceleration efforts than their respective offensive and defensive counterparts throughout the course of games, indicating the need for positional specificity in speed training for NCAA Division I football players.

The results of the present study provide novel insight into position-specific physical demands of NCAA Division I football games and provide physical performance staff with quantified information, which can potentially be used to replicate the physical demands of games in training. The present study demonstrated appreciable differences in the positional movement demands of NCAA Division I college football games, emphasizing the need for position-specific training to adequately prepare players for the rigors of competition.

PRACTICAL APPLICATIONS

The present study provided a novel analysis of the movement demands associated with NCAA Division I college football games. The results indicated significant differences in total running volume and high-intensity movement demands, most notably for the WR, DB, and LB position groups. Higher overall running loads were experienced for these three position groups, while greater high-intensity movement demands were required of the WR and DB groups. Data from the present study augments our understanding of the competitive demands experienced by NCAA Division I college football players, and

provides scope for the design of position-specific and game-specific physical conditioning strategies for coaches seeking to optimize training for the demands of competition.

Data from the present study support the use of position-specific training in the preparation of NCAA Division I college football players for competitive games. Maximizing performance and limiting the effects of fatigue are critical challenges for performance coaches, and as such, accounting for the physical demands associated with weekly training and games is imperative. Modifying weekly training loads of individuals within position groups involved in greater high-speed running volumes and a higher number of acceleration and deceleration efforts may mitigate fatigue, accelerate recovery, and improve subsequent performance. The WR, DB, and LB position groups are exposed to greater running volumes, faster running velocities, and a higher number of acceleration and deceleration efforts in games compared to their offensive and defensive counterparts, and may benefit from carefully monitored and individualized training load prescriptions throughout the week. Additionally, while RB and TE groups do not accrue the total distances during games of the WR group, they are exposed to greater running volumes than the OL, warranting individualized training load prescriptions governed by the physical demands of competition. Clearly, performance coaches seeking to optimize physical performance characteristics associated with competition must differentiate training programs based upon position-specific movement demands.

Data obtained from the present study provide a better understanding of the demands of NCAA Division I football and provide a foundation from which to implement a systematic approach to the development of individual and position-specific training programs. Future studies should examine how coaches seeking to enhance competitive performance, can manipulate individual and position-specific training programs to mitigate fatigue, enhance recovery, and optimize game-day performance.

ACKNOWLEDGEMENTS

No grant aid or manufacturer's aid was received in conjunction with the present study, and no conflicts of interest are declared. The results of this study do not constitute endorsement of the product by the authors of the National Strength and Conditioning Association.

REFERENCES

1. Austin, DJ and Kelly, SJ. Positional differences in professional Rugby League match play through the use of global positioning systems. *J Strength Cond Res* 27: 14-19, 2013.
2. Austin, DJ, and Kelly, SJ. Professional Rugby League positional match-play analysis through the use of global positioning system. *J Strength Cond Res* 28: 187-193, 2013.

3. Barbero-Álvarez, JC, Coutts, A, Granda, J, Barbero-Álvarez, V, and Castagna, C. The validity and reliability of a global positioning satellite system device to assess speed and repeated sprint ability (rsa) in athletes. *J Sci Med Sport* 13: 232-235, 2010.
4. Boyd, LJ, Ball, K, and Aughey, RJ. Quantifying external load in Australian Football matches and training using accelerometers. *Int J Sports Physiol Perform* 8: 44-51, 2013.
5. Cahill, N, Lamb, K, Worsfold, P, Headey, R, and Murray, S. The movement characteristics of English Premiership rugby union players. *J Sports Sci* 31: 229-237, 2013.
6. Coutts, AJ, and Duffield, R. Validity and reliability of GPS devices for measuring movement demands of team sports. *J Sci Med Sport* 13: 133-135, 2010.
7. Cunniffe, B, Pector, W, Baker, JS, and Davies, B. An evaluation of the physiological demands of Elite Rugby Union using global positioning system tracking software. *J Strength Cond Res* 23: 1195-1203, 2009.
8. DeMartini, JK, Martschinske, JL, Casa, DJ, Lopez, RM, Ganio, MS, Walz, SM, and Coris, EE. Physical demands of National Collegiate Athletic Association division I football players during preseason training in the heat. *J Strength Cond Res* 25: 2935-2943, 2011.
9. Edgecomb, SJ and Norton, KI. Comparison of global positioning and computer-based tracking systems for measuring player movement distance during Australian football. *J Sci Med Sport* 9: 25-32, 2006.

- 487 10. Gabbett, TJ, Jenkins, DG, and Abernethy, B. Physical demands of professional
488 rugby league training and competition using microtechnology. *J Sci Med Sport* 15:
489 80-86, 2011.
- 490 11. Granatelli, G, Gabbett, TJ, Briotti, G, Padulo, J, Buglione, A, D'Ottavio, S, and
491 Ruscello, BM. Match analysis and temporal patterns of fatigue in Rugby Sevens. *J*
492 *Strength Cond Res* 28: 728-734, 2014.
- 493 12. Iosia, MF, and Bishop, PA. Analysis of exercise-to-rest ratios during division I
494 televised football competition. *J Strength Cond Res* 22: 332-340, 2008.
- 495 13. Johnston, RD, Gabbett, TJ, and Jenkins, DG. Applied sports science of Rugby
496 League. *Sports Med* 44: 1087-1100, 2014.
- 497 14. Johnston, RD, Watsford, ML, Kelly, SJ, Pine, MJ, and Spurrs, RW. Validity and
498 interunit reliability of 10 Hz and 15 Hz units for assessing athlete movement
499 demands. *J Strength Cond Res* 28: 1649-1655, 2014.
- 500 15. Johnston, RD, Watsford, ML, Pine, MJ, Spurrs, RW, Murphy, AJ, and Pruyn, EC. The
501 validity and reliability of 5-Hz global positioning system units to measure team sport
502 movement demands. *J Strength Cond Res* 26: 758-765, 2012.
- 503 16. Kraemer, WJ and Gotschalk, LA. Physiology of American football. In: Exercise and
504 Sport Science. W.E. Garrett and D.T. Kirkendall, eds. Philadelphia: Lippincott,
505 Williams and Wilkins, 2000. pp. 795-813.
- 506 17. Krasnoff, JB, Kohn, MA, Choy, FKK, Doyle, J, Johansen, K, and Painter, PL.
507 Interunit and intraunit reliability of the RT3 triaxial accelerometer. *J Phys Act Health*
508 5: 527-538, 2008.

- 509 18. Loader, J, Montgomery, P, Williams, MD, Lorenzen, C, and Kemp, JG. Classifying
510 training drills based on movement demands in Australian Football. *Int J of Sports Sci*
511 *Coach 7*: 57-67, 2012.
- 512 19. McLellan, CP and Lovell, DI. Neuromuscular responses to impact and collision
513 during elite Rugby League match play. *J Strength Cond Res* 26: 1431-1440, 2012.
- 514 20. McLellan, CP and Lovell, DI. Performance analysis of professional,
515 semiprofessional, and junior elite Rugby League match-play using global positioning
516 systems. *J Strength Cond Res* 27: 3266-3274, 2013.
- 517 21. McLellan, CP, Coad, S, Marsh, D, and Lieschke, M. Performance analysis of Super-
518 15 Rugby match-play using portable micro-technology. *J Athl Enhanc* 2:5, 2013.
- 519 22. McLellan, CP, Lovell, DI, and Gass, GC. Biochemical and endocrine responses to
520 impact and collision during elite Rugby League match play. *J Strength Cond Res* 25:
521 1553-1562, 2011.
- 522 23. McLellan, CP, Lovell, DI, and Gass, GC. Creatine kinase and endocrine responses
523 of elite players pre, during, and post Rugby League match play. *J Strength Cond*
524 *Res* 24: 2908-2919, 2010.
- 525 24. McLellan, CP, Lovell, DI, and Gass, GC. Performance analysis of elite Rugby
526 League match play using global positioning systems. *J Strength Cond Res* 25: 1703-
527 1710, 2011.
- 528 25. Petersen, C, Pyne, D, Portus, M, and Dawson, B. Validity and reliability of gps units
529 to monitor cricket-specific movement patterns. *Int J Sports Physiol Perform* 2: 381-
530 393, 2009.

- 531 26. Rhea, MR, Hunter, RL, and Hunter, TJ. Competition modeling of American football:
532 observational data and implications for high school, collegiate, and professional
533 player conditioning. *J Strength Cond Res* 20: 58-61, 2006.
- 534 27. Robbins, DW, and Young, WB. Positional relationships between various sprint and
535 jump abilities in elite American football players. *J Strength Cond Res* 26: 388-397,
536 2012.
- 537 28. Sirotic, AC, Knowles, H, Catterick, C, and Coutts, AJ. Positional match demands of
538 professional Rugby League competition. *J Strength Cond Res* 25: 3076-3087, 2011.
- 539 29. Sullivan, C, Bilsborough, JC, Cianciosi, M, Hocking, J, Cordy, J, and Coutts, AJ.
540 Match score affects activity profile and skill performance in professional Australian
541 Football Players. *J Sci Med Sport* 17: 326-331, 2014.
- 542 30. Varley, MC, Fairweather, IH, and Aughey, RJ. Validity and reliability of GPS for
543 measuring instantaneous velocity during acceleration, deceleration, and constant
544 motion. *J Sports Sci* 30: 121-127, 2012.
- 545 31. Wisbey, B, Montgomery, PG, Pyne, DB, and Rattray, B. Quantifying movement
546 demands of AFL football using GPS tracking. *J Sci Med Sport* 13: 531-536, 2010.
- 547 32. Young, WB, Hepner, J, and Robbins, DW. Movement demands in Australian Rules
548 football as indicators of muscle damage. *J Strength Cond Res* 26: 492-496, 2012.