Truth disclosure on prognosis: Is it ethical not to communicate personalised risk of death?

Cardona, Magnolia; Kellett, John; Lewis, Ebony; Brabrand, Mikkel; Ní Chróinín, Danielle

Published in:
International Journal of Clinical Practice

DOI:
[10.1111/ijcp.13222](https://doi.org/10.1111/ijcp.13222)

Licence:
Other

[Link to output in Bond University research repository.](#)

Recommended citation(APA):

Cardona, M., Kellett, J., Lewis, E., Brabrand, M., & Ní Chróinín, D. (2018). Truth disclosure on prognosis: Is it ethical not to communicate personalised risk of death? *International Journal of Clinical Practice*, 72(8), [e13222]. <https://doi.org/10.1111/ijcp.13222>

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

For more information, or if you believe that this document breaches copyright, please contact the Bond University research repository coordinator.

Truth disclosure on prognosis: is it ethical not to communicate personalized risk of death?

Journal:	<i>International Journal of Clinical Practice</i>
Manuscript ID	IJCP-03-18-0099.R2
Wiley - Manuscript type:	Perspective
Date Submitted by the Author:	n/a
Complete List of Authors:	Cardona, Magnolia; Bond University and Gold Coast Hospital and Health Service , Centre for Research in Evidence-Based Practice; The University of New South Wales, The Simpson Centre for Health Services Research, South Western Sydney Clinical School Kellett, John; University of Southern Denmark Lewis, Ebony; The University of New South Wales, The Simpson Centre for Health Services Research, South Western Sydney Clinical School Brabrand, Mikkel; Odense University Hospital Ní Chróinín, Danielle; Liverpool Hospital, Department of Geriatric Medicine
Specialty area:	

SCHOLARONE™
Manuscripts

Truth disclosure on prognosis: is it ethical not to communicate personalized risk of death?

Summary

Predicting risk of death based on personalised and objective clinical indicators is an improvement over intuition and clinical judgment. Risk assessment can benefit clinicians by improving prognostic certainty, and truth disclosure helps patients and families by preventing futile management. Some argue that consent should be obtained before a patient is given an estimate of their prognosis as disclosure of bad news can overburden patients. In this article we argue that it is unethical not to use existing person-specific information to guide diagnosis and shared decision-making on management in partnership with well-informed patients. Disclosure of a poor prognosis should be normalized in personalized medicine, performed incrementally and with sensitivity so that it is acceptable to patients, and only occur if patients want to know it. However, a requirement of consent for truth disclosure should not be mandatory. Despite some level of imprecision, personalized risk estimations can be used to tailor management to the patient's informed wishes, and ensure that healthcare providers and families are acting ethically in the patient's best interest.

Keywords: risk assessment, prognosis, truth disclosure, clinical ethics, personalized medicine

Prognostication is a challenging art, and science. Through advanced age, increasing frailty and chronic illness, treatment becomes increasingly complex and the clinician's ability to prognosticate is tested further. Prognosis does not solely rely on the natural history of a major disease but on the interaction with other comorbidities, the choice of treatment and response to it, and other patient-specific socio-demographic influences, all of which may impact on the accuracy of the prediction. Yet prognostication has become a lost skill: at the end of the nineteenth century *prognostication* took up about ten percent of medical textbooks, but by 1970 it was hardly addressed.¹ As a result, although modern physicians commonly encounter situations that require prognostication, they feel "poorly trained to do it, and find the whole process difficult and stressful" and, therefore, may favor partial or non-disclosure altogether.²

Personalized medicine has been broadly defined as "*the tailoring of medical treatment to the individual characteristics, needs and preferences of a patient during all stages of care, including prevention, diagnosis, treatment and follow-up*".³ Data to assist personalized decisions tailored to individuals, thus departing from the "one size fits all" paradigm⁴ is valuable as disease trajectories vary, and individuals respond differently to treatment options and value different states of health. The development of prediction tools for all ages, and in particular the very old,⁵ provides information on likely outcomes, including death, and is but one essential component of truthful communication.

Delivering a prognosis, especially a bad one, is complex, takes time, and requires sensitive communication filled with good judgment, skill, empathy and compassion. Some patients may hold beliefs that can distort their understanding or expectations of cure; others may choose to know less as disease progresses or ask more about the risk of treatments; and families may want to know more about what to expect of the course of illness and the time to death. So proficiency in tactful disclosure becomes necessary when dealing with older patients suffering complex co-morbidities.⁶ However, many patients and families may find confronting such an explicit prognosis challenging. This is largely because popular culture continues to promote the myth that there is an effective treatment for every illness and that medical technology, no matter how painful or expensive, should be used at all costs. All deaths are, therefore perceived as "failures" of medicine.⁷ Even today, some medical and nursing schools' curricula do not emphasise sensitive communication training; and some health systems are yet to abandon paternalistic attitudes and therefore are still not

systematically encouraging patient autonomy, either through voluntary or mandatory approaches. The solution is not to avoid the difficult discussion using unpredictability of disease progression as an excuse and thus deny the patient their right to know, but to improve expertise in providing a personalized prognosis in order to empower patients to take control of their immediate or future care.

Although giving information on the progression and recurrence of illness, along with advice and shared decision-making, is now a routine- and accepted- part of cancer care, there is concern that unwanted information may overburden patients and families and take away hope, so that in geriatric and critical care populations providing a prognosis is sometimes viewed as overriding patient autonomy and potentially causing harm.⁸ It has therefore been suggested by others that consent for disclosure should be obtained before a patient is told their risk of death. Yet, in routine practice with non-terminal patients anyone who consults a physician implies consent to being given a professional opinion on their probable diagnosis, prognosis and treatment. While some prognostically important tests require explicit informed consent, as for HIV testing, neither evidence nor law provide clear guidance on the need for or benefit of detailed, written consent for discussions about other diagnoses. At present, the judgment about whether or not to disclose prognosis is often left up to the physician. The appropriateness of disclosing prognosis to elderly patients who have their lost decision-making capacity and do not have a surrogate decision-maker is particularly concerning, because such patients would not be able to make subsequent decision. It is said that “*consent was developed for research ethics, not for clinical care.*”⁹ While consent to perform invasive procedures is the norm and a service requirement, we believe consent to communicate prognosis is both a moral obligation and an ethical requirement that fosters patient autonomy.

Although it is important to respect the right of patients not to know their prognosis, telling a patient their risk of death can help them wisely direct their subsequent treatment. There are many benefits provided by an accurate estimation of a patient’s time to death and their clinical course prior to death: for patients and their relatives being clearer that cure is unlikely and burdensome interventions can be harmful, provides opportunities to discuss preferences, consider comfort and palliative care if indicated, and coordinate services accordingly.¹⁰ In fact, increased certainty regarding disease trajectory can be liberating for some, as it puts a timeframe to the end of suffering, assists in the planning of their final days with good quality, and helps them to embrace the inevitable: death from old age, frailty, chronic illness, and

multi-organ failure,¹¹ concepts that modern society often attempts to deny. For clinicians, an accurate prognosis can help avoid futile and potentially harmful therapies, assist end-of-life care that best responds to patient and family values and preferences, and ensure better stewardship of scarce health system resources.¹² But delivery of patient-centred care would of course include adherence to the patient's disclosure preferences.

One of the pillars of personalized medicine is the sharing of prognostic data between patient and clinician so that patients engage in a proactive and dynamic decision-making partnership.¹³ Not only is outcome estimation a necessary step in routine practice, many clinicians find it a rewarding experience, as patients often welcome their guidance at a seminal time in their lives. Factors such as education, age, disease type, availability of social or medical support, religion or cultural background, family dynamics and the way in which prognosis is delivered impact on people's openness to the news of an impending death. Certainly, no one wants to receive or digest bad news, and there will always be those who prefer little or no disclosure of poor prognosis in terminal illness; this is their right. Nevertheless, we believe it is unethical and unjust not to formulate an explicit person-specific prognosis when the information and technology that can do so are available. Withholding that information undermines justice and autonomy, except if the concept of autonomy extends to the right to not know.¹⁴ The estimated risk of death and the benefits of less aggressive interventions should be disclosed to all patients, unless they choose otherwise, and it is not in the patient's best interest to withhold information or give false hope with overoptimistic outcome predictions, as this may lead to loss of patient trust. In the case of chronic or progressive disease, ideally, patients and healthcare professionals will have the opportunity to revisit disclosure preferences, as with other goals of care, over time, as these may change. The timing of re-discussion will need to be individualised to the patient.

A more accurate, personalized prognosis also allows patients to more confidently formulate advanced care directives, reflecting their values and goals. The absence of health care directives, based on accurately estimated likely outcomes, hampers wise decision-making and often leads to regret when clinicians or families have to make decisions on behalf of patient without knowing their wishes. Without them clinicians can also experience job dissatisfaction and moral distress if they feel urged to comply with family requests to administer futile treatment. While patients may change their mind in either direction about treatment preferences as their age increases, their disease progresses and their ability to partake in

decision-making varies , accurate prediction of their individual risks may make this change of mind less likely. An individualised prognosis that is based on the unique health determinants of an individual’s profile will improve patient’s understanding of their own disease, and clarify their own goals of care. Since all involved in delivering care will have a realistic perception of the patient’s life expectancy, prevention of unnecessary, potentially harmful treatments would be prevented, distributive justice ¹⁵ may be achieved, and the family’s anticipatory grief may also be reduced.

In summary, despite the ever-present uncertainty in healthcare, personalised risk prediction near the end of life is possible, and a professional responsibility. Prognostic disclosure should be normalised and offered regulatory support without the barriers of formal consent procedures. Timely communication of a poor prognosis and risk of death needs a sensitive, incremental approach and should be in a format that patients and families find acceptable and can understand. Precise predictions of the future are never possible, but well validated actuarial science and recent advances in information technology, genetics and other areas of medical biology now make it possible to formulate much improved estimates of the likely outcomes for individual patients. Even if some level of imprecision remains, personalized risk estimations should inform care pathways, as they make it possible to tailor management to the patient’s informed wishes, and ensure that physicians, family and other healthcare providers are acting in the patient’s best interest.

References

1. Kellett J. Prognostication- The lost skill of medicine *European Journal of Internal Medicine*. 2008;19(3):155-64.
2. Lamont EB, Christakis NA. Prognostic disclosure to patients with cancer near the end of life. *Annals of internal medicine*. 2001;134(12):1096-105.
3. US Food and Drug Administration. *Paving the Way for Personalized Medicine. FDA’s Role in a New Era of Medical Product Development*. . October 2013 ed: FDA, US Department of Health and Human Services; 2013. p. 61.
4. Tutton R. Personalizing medicine: futures present and past. *Social science & medicine* (1982). 2012;75(10):1721-8.
5. Cardona-Morrell M, Hillman K. Development of a tool for defining and identifying the dying patient in hospital: Criteria for Screening and Triaging to Appropriate aLternative care (CriSTAL). *BMJ supportive & palliative care*. 2015;5(1):78-90.
6. Woo JA, Maytal G, Stern TA. Clinical Challenges to the Delivery of End-of-Life Care. *Primary Care Companion to The Journal of Clinical Psychiatry*. 2006;8(6):367-72.
7. Thomas L. Dying as Failure. *The Annals of the American Academy of Political and Social Science*. 1980;447:1-4.
8. Rudnick A. Informed consent to breaking bad news. *Nursing ethics*. 2002;9(1):61-6.

9. Smith AK, Glare P. Ethical Issues in Prognosis and Prognostication Oxford Handbooks Online; 2014 [cited 2018 January]. Available from: <http://www.oxfordhandbooks.com/view/10.1093/oxfordhb/9780199974412.001.0001/oxfordhb-9780199974412-e-9>.
10. Finlay E, Casarett D. Making difficult discussions easier: using prognosis to facilitate transitions to hospice. *CA: a cancer journal for clinicians*. 2009;59(4):250-63.
11. Howard M, Bernard C, Tan A, Slaven M, Klein D, Heyland DK. Advance care planning: Let's start sooner. *Canadian Family Physician*. 2015;61(8):663-5.
12. Zapata JA, Moriates C. The High-Value Care Considerations of Inpatient versus Outpatient Testing. *American Medical Association Journal of Ethics*. 2015;17(11):1022-7.
13. Corcoran G, Lyons C, O'Donoghue M, Donnelly U, Lawlor L, Crinion G, et al. 026A Plan for Every Patient: Achieving Excellence in Communication and Interdisciplinary Working on a Specialist Geriatric Ward. *Age and Ageing*. 2017;46(Suppl_3):iii13-iii59.
14. Helgesson G. Autonomy, the Right Not to Know, and the Right to Know Personal Research Results: What Rights Are There, and Who Should Decide about Exceptions? *The Journal of law, medicine & ethics : a journal of the American Society of Law, Medicine & Ethics*. 2014;42(1):28-37.
15. Gefenas E, Cekanauskaite A, Tuzaitė E, Dranseika V, Characiejus D. Does the "new philosophy" in predictive, preventive and personalised medicine require new ethics? *EPMA Journal*. 2011;2(2):141-7.