

Bond University
Research Repository



Higher- Versus Lower-Intensity Strength-Training Taper: Effects on Neuromuscular Performance

Pritchard, Hayden J; Barnes, Matthew J; Stewart, Robin J; Keogh, Justin W; McGuigan, Michael R

Published in:
International Journal of Sports Physiology and Performance

DOI:
[10.1123/ijsp.2018-0489](https://doi.org/10.1123/ijsp.2018-0489)

[Link to output in Bond University research repository.](#)

Recommended citation(APA):
Pritchard, H. J., Barnes, M. J., Stewart, R. J., Keogh, J. W., & McGuigan, M. R. (2019). Higher- Versus Lower-Intensity Strength-Training Taper: Effects on Neuromuscular Performance. *International Journal of Sports Physiology and Performance*, 14(4), 458-463. <https://doi.org/10.1123/ijsp.2018-0489>

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

For more information, or if you believe that this document breaches copyright, please contact the Bond University research repository coordinator.

Note. This article will be published in a forthcoming issue of the *International Journal of Sports Physiology and Performance*. The article appears here in its accepted, peer-reviewed form, as it was provided by the submitting author. It has not been copyedited, proofread, or formatted by the publisher.

Section: Original Investigation

Article Title: Higher vs. Lower Intensity Strength Training Taper: Effects on Neuromuscular Performance

Authors: Hayden J. Pritchard^{1,2}, Matthew J. Barnes³, Robin J. Stewart¹, Justin W. Keogh^{4,2,5}, and Michael R. McGuigan^{2,6}

Affiliations: ¹Faculty of Health Sciences, Universal College of Learning, Palmerston North, New Zealand; ²Sports Performance Research Institute New Zealand, Auckland University of Technology, Auckland, New Zealand; ³School of Sport and Exercise, Massey University, Palmerston North, New Zealand; ⁴Faculty of Health Sciences and Medicine, Bond University, Queensland, Australia; ⁵Faculty of Science, Health, Education and Engineering, University of the Sunshine Coast, Queensland, Australia; ⁶School of Medical and Health Sciences, Edith Cowan University, Western Australia, Australia.

Journal: *International Journal of Sports Physiology and Performance*

Acceptance Date: September 3, 2018

©2018 Human Kinetics, Inc.

DOI: <https://doi.org/10.1123/ijspp.2018-0489>

sessions, at the same time in the morning (± 30 min). Testing occurred prior to each four-week training period (T1), 36-60 hours after the final training session of the four week period (T2), and 36-60 hours after their final taper training session (T3). The time between final training sessions and testing times was kept consistent for each participant. For example, if a participant was tested 36 hours following training in Condition A (i.e. trained late afternoon on Thursday, had one full day’s rest Friday, tested on Saturday morning) this was kept consistent for Condition B.

Gym-based Testing

In order to establish training loads, participants performed 1RM testing, according to National Strength & Conditioning Association (NSCA) guidelines¹⁵ for the three powerlifts (squat, bench press and deadlift) within one week of the first testing session. Additionally, participants were also tested for a 2-8RM on all other programmed lifts. 1RM was estimated from these results using the following formula:

$$1RM = \text{Load} / (1.0278 - 0.0278 * \text{Repetitions Performed})^{16}$$

1RM testing was repeated during the week between conditions. All strength testing was performed at the same time of day (\pm one hour).

Resting Measures

Saliva was collected and CK measured using previously described techniques¹⁷. Saliva samples were analysed in duplicate according to manufacturer’s instructions using an enzyme-linked immunosorbent assay (ELISA; DRG International, USA). The coefficient of variation (CV) of duplicate ELISA samples was 3.9% for testosterone and 8.5% for cortisol. The typical CV of CK measurements using the Reflotron® systems spectrophotometer (Roche Diagnostics, Switzerland) is 3.5%.

The Daily Analysis of Life Demands in Athletes (DALDA) questionnaire was then completed by participants (as well as prior to each training session, as part of the training records kept by participants). The frequency of ‘worse than normal’ results was recorded at each testing session and was analysed to determine any changes across testing sessions. This questionnaire has been used previously to monitor athletes^{18,19}.

Performance Measures

Participants then undertook performance tests consisting of the countermovement jump (CMJ), IMTP, and isometric bench press (IBP) on a force plate (400 Series, Fitness Technology, Australia). Performance test procedures occurred as previously described¹⁷. Analysis occurred using the Ballistic Measurement Systems software (BMS, Innervations, Australia). Three maximal efforts were undertaken for each performance test, with the effort that had the highest jump (CMJ) or greatest peak force (IMTP and IBP) utilised for analysis. For the CMJ, the CV was 3.0% for jump height and 7.0% for flight-time: contraction-time. For the IMTP, CV was 3.3% for IMTP peak force. For IBP, CV was 3.9% for IBP peak force.

Training protocol

After the first testing session, participants commenced their training program, focused on improvement in the powerlifts. The objective of this four-week program (see Table 1) was to bring all participants to a similar level of training and fatigue prior to the taper week. This four week program has been used previously to successfully enhance 1RM performance of the powerlifts²⁰. Participants were instructed to separate each of the three training days with at least one full rest day between, i.e. if a strength training day occurred on Tuesday, the next strength training day would be Thursday. Participants could perform their habitual aerobic or conditioning focused training, but no further resistance training was allowed – although a limitation, this permitted greater recruitment from a limited local participant pool.




Figure 3. Taper B - Jump height individual responses

Table 1: Training program.

Day	Exercise	Week One			Week Two			Week Three			Week Four		
		Reps	Sets	Intensity	Reps	Sets	Intensity	Reps	Sets	Intensity	Reps	Sets	Intensity
1	Bench Press	4	3	80%	4	3	82.5%	3	4	85%	3	4	87.5%
1	Back Squat	6	4	75%	6	4	77.5%	4	4	80%	4	4	82.5%
1	Military Press	6	4	75%	6	4	77.5%	4	4	80%	4	4	82.5%
1	Barbell Row	10	3	70%	10	3	72.5%	8	4	75%	8	4	77.5%
2	Deadlift	4	3	80%	4	3	82.5%	3	4	85%	3	4	87.5%
2	Close Grip Bench Press	6	4	75%	6	4	77.5%	4	4	80%	4	4	82.5%
2	Deficit Deadlift	6	4	75%	6	4	77.5%	4	4	80%	4	4	82.5%
2	Good Morning	10	3	70%	10	3	72.5%	8	4	75%	8	4	77.5%
3	Back Squat	4	3	80%	4	3	82.5%	3	4	85%	3	4	87.5%
3	Paused Bench Press	6	4	75%	6	4	77.5%	4	4	80%	4	4	82.5%
3	Front Squat	6	4	75%	6	4	77.5%	4	4	80%	4	4	82.5%
3	Barbell Row	10	3	70%	10	3	72.5%	8	4	75%	8	4	77.5%

N.B. Intensity is percentage of 1RM; Deficit Deadlift was with feet raised on a 2” plate; Paused Bench Press had a two second pause on the chest.

Table 2: Taper programs.

Day	Exercise	Taper Week A			Taper Week B		
		Reps	Sets	Intensity	Reps	Sets	Intensity
1	Bench Press	4	2	90%	3	3	80%
1	Back Squat	3	3	82.5%	3	4	70%
2	Deadlift	4	2	90%	3	3	80%
2	Close Grip Bench Press	3	3	82.5%	3	4	70%
3	Back Squat	4	2	90%	3	3	80%
3	Paused Bench Press	3	3	82.5%	3	4	70%

N.B. Intensity is percentage of 1RM; Paused Bench Press had a two second pause on the chest.

Table 3: Performance results.

Performance Measures				
	CMJ Height (cm)	CMJ FT: CT	IMTP Relative Peak Force (N/kg)	IBP Relative Peak Force (N/kg)
A1	37.8 ± 5.3	0.738 ± 0.171	34.0 ± 4.8	18.6 ± 2.2
A2	39.4 ± 5.7*	0.788 ± 0.163*	35.9 ± 5.0*	18.4 ± 3.0
A3	40.2 ± 5.7*	0.803 ± 0.138*	35.9 ± 5.5*	18.4 ± 2.6
B1	38.2 ± 5.9	0.755 ± 0.160	35.4 ± 5.4	18.8 ± 2.9
B2	39.2 ± 5.2	0.793 ± 0.172*	35.9 ± 4.8	18.6 ± 2.8
B3	39.9 ± 5.3*	0.771 ± 0.169	35.2 ± 5.0	19.1 ± 2.6
P1	38.0 ± 5.5	0.747 ± 0.162	34.7 ± 5.0	18.7 ± 2.5
P2	39.3 ± 5.3*	0.791 ± 0.163*	35.9 ± 4.8*	18.5 ± 2.8
P3	40.0 ± 5.3*	0.787 ± 0.151*	35.5 ± 5.1	18.7 ± 2.5

(A = +5% Intensity Taper; B = -10% Intensity Taper; P = Pooled Data; Numbers indicate testing time point, 1 = Pre-training; 2 = Post-training, 3 = Post-taper).

* represents a small ES improvement compared to that conditions baseline. CMJ = countermovement; FT: CT = flight-time: contraction-time; IMTP = isometric mid-thigh pull; IBP = isometric bench press.

Table 4: Non-Performance Test Results.

Non-Performance Test Results					
	Cortisol (ng/ml)	Testosterone (pg/ml)	T/C Ratio (x 1,000)	Creatine Kinase (I/U)	DALDA “worse than’s”
A1	7.65 ± 2.90	150.77 ± 46.43	25.10 ± 14.59	296.4 ± 216.6	4.2 ± 2.6
A2	8.89 ± 4.74	155.70 ± 45.92	24.25 ± 15.41	220.8 ± 101.6	4.6 ± 2.9
A3	8.62 ± 3.79	151.47 ± 41.31	20.42 ± 7.13	246.7 ± 136.6	3.5 ± 3.2
B1	9.20 ± 6.72	156.72 ± 55.00	24.00 ± 15.60	282.5 ± 155.6	3.4 ± 2.2
B2	8.94 ± 4.29	146.87 ± 35.57	20.53 ± 10.62	319.5 ± 204.9	2.6 ± 2.0
B3	8.04 ± 5.19	138.16 ± 33.27	21.02 ± 9.15	223.4 ± 162.6	2.1 ± 1.8

(A = +5% Intensity Taper; B = -10% Intensity Taper; Numbers indicate testing time point, 1 = Pre-training; 2 = Post-training, 3 = Post-taper)