

Bond University  
Research Repository



**Parent-child interaction therapy**  
**An evidence-based treatment for child maltreatment**

Thomas, Rae; Zimmer-Gembeck, Melanie J.

*Published in:*  
Child Maltreatment

*DOI:*  
[10.1177/1077559512459555](https://doi.org/10.1177/1077559512459555)

*Licence:*  
Other

[Link to output in Bond University research repository.](#)

*Recommended citation(APA):*  
Thomas, R., & Zimmer-Gembeck, M. J. (2012). Parent-child interaction therapy: An evidence-based treatment for child maltreatment. *Child Maltreatment*, 17(3), 253-266. <https://doi.org/10.1177/1077559512459555>

**General rights**

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

For more information, or if you believe that this document breaches copyright, please contact the Bond University research repository coordinator.

Running Head: **PCIT AND CHILD MALTREATMENT**

Parent-Child Interaction Therapy: An Evidence-Based Treatment for Child Maltreatment

Rae Thomas<sup>1,2</sup> and Melanie J. Zimmer-Gembeck<sup>1</sup>

<sup>1</sup> Behavioural Basis of Health, School of Applied Psychology, Griffith University

<sup>2</sup> Centre for Research in Evidence-Based Practice, Faculty of Health Sciences and Medicine,  
Bond University

Author Notes:

Some of these data were presented at the annual meeting of the Society for Prevention Research in June 2011.

This research was partially supported by funding from Future Directions Prevention and Early Intervention Trials, Queensland Department of Child Safety, Australia.

We greatly appreciate the assistance of families who participated in this research. In addition, we wish to thank members of the Family Interaction Research Program at Griffith University for their willingness to share ideas and for continuing to work with and support children and their parents.

Correspondence concerning this article should be addressed to Rae Thomas, Centre for Evidence-Based Practice, Faculty of Health Sciences and Medicine, Bond University, Gold Coast, Queensland, Australia 4229. Tel: + 61 7 5595 4482; Fax: +61 7 5595 1271  
Electronic mail may be sent to [rthomas@bond.edu.au](mailto:rthomas@bond.edu.au).

### **Abstract**

It is common practice to augment efficacious treatment protocols for special populations (Durlak & DuPre, 2008), but this is often done before establishing that standard services are not appropriate. In this randomized controlled trial with families at risk or with a history of maltreatment ( $N = 151$ ), we investigated the effectiveness of *standard* 12-session Parent-Child Interaction Therapy (PCIT). This is in contrast to other PCIT studies with similar parents, which have allowed for longer and sometimes variable treatment length and with modifications to PCIT protocol. After treatment and compared to Waitlist, mothers reported fewer child externalizing and internalizing behaviors, decreased stress and were observed to have more positive verbalizations and maternal sensitivity. These outcomes were equivalent or better than outcomes of our previous PCIT trial with high-risk families (Thomas & Zimmer-Gembeck, 2011) when treatment length was variable and often longer. These findings support standard protocol PCIT as an efficacious intervention for families in the child welfare system.

## **Parent-Child Interaction Therapy: An Evidence-Based Treatment for Child Maltreatment**

Despite best efforts of child protection systems and increased government expenditure, child maltreatment rates continue to remain at unacceptably high levels (Australian Institute of Health and Welfare 2012; Hussey, Chang, & Kotch, 2006) and child protection systems struggle to provide effective interventions and treatments to high-risk families. Budgetary increases of over 13% per annum (Bromfield, Holzer, & Lamont, 2011) resulting in billions of dollars in child protection, prevention and intervention seem to do little to stem the tide. Generalist parenting programs known to be effective in increasing parenting skills and decreasing child behavior problems are often utilised in mainstream contexts, however, few of these have met the criteria for evidence-based treatment (EBT; Chambless & Ollendick, 2001), and there have been few interventions founded in theory about the causes of child maltreatment (Allen, Gharagozloo & Johnson, 2012). In contrast, Parent-Child Interaction Therapy (PCIT), a well-known parenting program for child externalizing behavior problems, has theoretical foundations that support its utility as a treatment approach for this population and has accumulated data to support its contention as an EBT for families who have experienced child maltreatment (Chaffin, Funderburk, Bard, Valee, & Gurwich, 2011; Thomas & Zimmer-Gembeck, 2011; Timmer, Ware, Urquiza, & Zebell, 2010).

However, efficacious interventions are often considered inappropriate for certain populations and may be adapted with the aim of better meeting the needs of those accessing the services (Axford & Little, 2009). Several behavior management and parenting EBTs have adapted the original design to target the needs of specialized populations. For example the Triple P Parenting Program has been adapted for obese and gifted children (Morawska & Sanders, 2009; West, Sanders, Cleghorn & Savies, 2010) and The Incredible Years program has been adapted for bereaved and substance abusing parents (Braiden, McDaniel, Duffy &

McCann, 2011; Stangera, Ryan, Hongyun & Budney 2011). This trend is also seen in many of the PCIT research studies working with subpopulations (Berkovits, O'Brien, Carter, & Eyberg, 2010; Pincus, Eyberg, & Choate, 2005). Therefore, it is not surprising that PCIT researchers providing services to parents who have maltreated their children have altered the standard PCIT design for this population (Chaffin et al., 2004; Chaffin et al., 2011; Timmer, Zebell, Culver & Urquiza, 2010). However, the assumption that particular subgroups of parents require different parenting interventions has not been tested. The purpose of the current study was to investigate the effectiveness of a *Standard* PCIT treatment protocol with mothers who were at high-risk or who had a history of maltreating their children.

### **Parent-Child Interaction Therapy for High-Risk Families**

PCIT was developed for children aged between 3 and 7 years diagnosed with an externalizing behavior problem and their parents (Hembree-Kigin & McNeil, 1995) and is founded in social learning theory and attachment theory. Parent-child dyads are observed through a one-way mirror, and by using a bug-in-the-ear device, parents are coached to attend to the child's behaviors consistently and predicably. Parents are taught behavior management strategies that focus on positive reinforcement rather than power assertion to reduce child oppositional and disruptive behaviors. The behavior management techniques in PCIT are designed to aid children's emotion regulation by providing parents with developmentally appropriate language and skills (Hembree-Kigin & McNeil, 1995).

Evidence suggests that the most proximal risks of child maltreatment are negative and coercive patterns of parent-child interactions and parents' lack of knowledge or inappropriate use of discipline (Cicchetti & Valentino, 2006). This is because they often result in escalating coercive exchanges, harsh discipline strategies, and aggressive communication techniques (Patterson, 1982). Further, parents with a history of maltreating their children are likely to be less sensitive in interactions and more likely to have an insecure or disorganized

caregiver-child attachment compared to other mothers (Baer & Martinez, 2006). Both community and lab-based PCIT treatment outcome studies have generally produced statistically significant and medium to large effect sizes in reductions in child externalizing behavior and parental stress, increases in positive observations of parent behavior (see Thomas & Zimmer-Gembeck, 2007 for a review), and more recently, increases in maternal sensitivity (Thomas & Zimmer-Gembeck, 2011). Because each of these therapeutic outcomes has also been identified as a risk factor associated with child maltreatment, PCIT has been identified as an EBT for families at child maltreatment risk (Kauffman Best Practices Project, 2004). However, in investigations of PCIT and child maltreatment, the standard PCIT protocol has been adapted in many different ways making it difficult for the community practitioner to clearly identify which PCIT protocol to implement with their high-risk and abusive parents.

For example, Chaffin and colleagues (2004) randomly assigned parent-child dyads with a history of child maltreatment to one of three treatment conditions; two PCIT conditions and a treatment as usual condition. Both PCIT groups were augmented with a six-session motivation component prior to commencing PCIT, however in addition to the motivation component, the second PCIT group participants were also offered individual counselling in addition to PCIT. Results indicated reductions in rates of future referrals to child welfare in the PCIT + motivation group compared to the treatment as usual group. However, contrary to expectations, the PCIT group participants who also received individual therapy were not significantly different than the PCIT+motivation group in terms of renotification rates. There was a trend suggesting that increasing services (individual counselling) to PCIT participants attenuated parent and child outcomes. More recently and with similar families, Chaffin et al. (2011) compared PCIT and services as usual with two orientation conditions (motivation and psychoeducation). Participants with a motivation orientation combined with PCIT had fewer

notifications to child welfare after treatment than either the PCIT group combined with psychoeducation or either of the service as usual groups. To evaluate the orientation addition to PCIT, the authors recommended future research compare non-orientated PCIT and PCIT with a motivation orientation.

PCIT with high-risk families has also been augmented with either an in-home coaching component or an in-home social support component (Timmer, Zebell et al., 2010). It was expected that augmenting PCIT with in-home coaching would enable greater remediation of inappropriate parent-child interaction, produce greater reductions in child externalizing behavior and parent stress, and greater increases in positive parent-child communications. However, at mid-treatment assessment there were no differences between groups on parental report of child behavior change, but the in-home coaching group parents did report reduced stress compared to the social support group participants. No differences between groups were found also for the rate of skill acquisition or quality of parent-child interactions at treatment completion. Therefore, augmenting PCIT with either in-home coaching or social support did not alter treatment outcomes at completion.

Finally, in a third randomized controlled trial (RCT) of PCIT, treatment was allowed to vary in length (Thomas & Zimmer-Gembeck, 2011). Participants concluded PCIT when they had attained skills to a specified level (Mastery Criteria) and could demonstrate changes in behavior management strategies. This approach differed from the other PCIT trials due to the open-ended treatment time-frame. Compared to a waitlist comparison group at 12-weeks of treatment, PCIT participants had significantly less externalizing child behavior problems and parental stress, and significantly greater observed positive communication. These positive outcomes were strengthened for PCIT participants after completion of the treatment protocol and similar to Chaffin et al. (2004), participants who completed PCIT were significantly less likely to be notified to child welfare for any maltreatment complaint than

those who did not complete PCIT. Despite overall positive results for participants in PCIT groups, no study with a high-risk cohort has investigated the effectiveness of Standard PCIT. Community practitioners need better guidance regarding which PCIT adaptation to use or whether the standard protocol might be similarly effective.

### **New Directions in Services to Improve Parenting and Decrease Child Abuse**

Although Governments continue to increase funding for child maltreatment (Bromfield et al., 2011), many of these funds go to much needed front-line child welfare services and therapeutic agencies are required to do more with fewer resources. Lengthy treatments are often cost-prohibitive and may not produce more benefit than treatments of shorter duration, potentially reducing the cost-effectiveness of more intensive treatments. In a meta-analysis of sensitivity and attachment interventions in early childhood (Bakermans-Kranenburg, van IJzendoorn, & Juffer, 2003), results showed that some of the most effective interventions were not lengthy and interventions with fewer contacts appeared somewhat more effective than interventions with many sessions. Second, it was concluded that some interventions were effective regardless of whether families experienced multiple problems. The present study was designed to address the effectiveness of an EBT (i.e., PCIT) with high-risk families using the standard and original protocol without adaptation.

Hence, the purpose of the current study was to assess the effectiveness of Standard PCIT with families engaged in or at risk of child maltreatment. Instead of providing additional components, we minimised the length of the intervention protocol to determine if outcomes were as positive as adapted PCIT interventions. PCIT has two phases. In this study progression from the first to the second occurred after 6 to 8 coaching sessions regardless of Mastery Criteria and treatment concluded when a maximum of 12 coaching sessions had been conducted. Participants were referred because they had a history or were assessed to be at high risk of child abuse.



Previous research has shown that PCIT is effective in reducing child externalizing behavior problems, parent stress, and child abuse potential, and increasing positive parent-child interactions and maternal sensitivity; these are all correlates of child maltreatment. In the current study, outcomes of the standard PCIT protocol were compared to those of a supported waitlist control group. As PCIT is a dyadic therapy, both parent and child outcomes are important. Therefore, the primary outcomes for the study were an expected decrease in child externalizing behavior and increase in observed parent sensitivity. Overall, we expected that using the standard PCIT protocol for families engaged in or at risk of child maltreatment would be as effective as adapted PCIT for the same population. We expected decreases in externalizing and internalizing child behavior, parent stress, parent depression, abuse potential and observed decreases in negative parent communication, and increases in positive communication and parent sensitivity. Due to consistency in measures and sample population, outcomes of the current trial were compared to our previous PCIT effectiveness trial (Thomas & Zimmer-Gembeck, 2011) to determine whether the standard PCIT treatment protocol was as effective as the lengthier version for high-risk parents.

## **Method**

### **Participants**

Participants were 151 female caregivers ( $M_{\text{age}} = 33.9$  years,  $SD = 7.31$ ) and their children ( $M_{\text{age}} = 4.57$  years,  $SD = 1.3$ ; 70.4% boys). For brevity and to simplify language, the term parent will be used to identify the group of female caregivers in this study. The majority of parents were born in Australia (74%) with 1.4% being of Aboriginal or Torres Strait Islander descent. Most mothers had completed some high school (81%) and 16.5% had some tertiary education.

The intervention was delivered by a well-known tertiary referral service and research program for families at high-risk of, or engaged in, child maltreatment. Families were referred after having completed alternative parenting courses and assessed by referrers as requiring

further intervention. A semi-structured interview was developed and administered by researchers at pre-assessment to assess child maltreatment risk using proximal risk factors including high levels of parent distress, inappropriate discipline strategies, and aggressive communication. All families indicated use of corporal punishment in their discipline strategies, expressed high levels of frustration and intolerance with child behavior and high levels of parental distress.

Participants were referred from child protection authorities (34.2%), government health services (19.7%), and education and non-government social service organizations (18.4%). Parent self-referrals also were accepted (27.6%), but the pre-assessment semi-structured interview had to reveal prior parenting interventions, high risk for child maltreatment and significant levels of child behavioral problems. Families referred from child protection were classified as having engaged in child maltreatment. We were not able to reliably ascertain child protection status for those families referred from sources other than the child protection authority. Therefore, all families were assessed at pre-assessment using a semi-structured interview and families referred from sources other than child protection were classified as high-risk.

Although, all participants in the current study referred from child protection authorities were considered as having engaged in child maltreatment, it has been argued that families who come to the attention of the child welfare authorities represent only the most severe end of the continuum of child maltreatment (Cicchetti & Toth, 2005; Manly, 2005) and it is often concluded that the actual child maltreatment rate exceeds official estimates. A reliance on substantiated or reported cases of child maltreatment for inclusion criteria, therefore, not only may produce research findings that are not fully generalizable to the broader population of parents who have maltreated their children, but also precludes some parents from interventions because the report of maltreatment was not substantiated or reported. Psychosocial maladaptation occurs in both substantiated and unsubstantiated cases (Drake, 1996). Hussey and colleagues (2005) found the behavioral and developmental

outcomes of a high-risk sample of 8-year old children with both substantiated and unsubstantiated reports of child maltreatment were indistinguishable. Therefore, the recruitment strategy of the current study may be more representative of the general population of maltreating parents than would be found when relying on referrals from child protection sources only (Hussey et al., 2005).

To determine whether the current participants differed at pre-assessment when compared to participants in previous published studies who were all referred from child welfare agencies, we compared published levels of pre-assessment measures to our findings. We found no differences in child abuse potential scores ( $t = 1.47, p = .14$ ; Chaffin et al., 2004), or internalizing behaviors ( $t = -1.04, p = .30$ ; Timmer, Ware et al., 2010). However, compared to Timmer, Ware et al. (2010) participants in the current study reported *greater* child externalizing behavior ( $t = -2.43, p = .01$ ), *greater* behavior intensity ( $t = -3.37, p = .0003$ ) and *greater* problematic behavior ( $t = -2.41, p = .02$ ).

In the current study, other than information regarding sexual abuse, access to specific maltreatment history for families referred from child protection authorities was not available. However, different subtypes of maltreatment have been reported to coexist in a majority of families (Cicchetti & Valentino, 2006) and it is likely that this was the case for almost all families in the current study. However, PCIT has been contraindicated for children with a history of sexual abuse. Therefore, children were excluded if there was any suspected sexual abuse history based on information revealed during the initial interview with parents or from child protection authorities.

The present study was part of a larger RCT of PCIT where participants were allocated to Time-Variable PCIT (TV/PCIT), Standard PCIT (S/PCIT) or Waitlist. The RCT comparing PCIT and a Waitlist group has been conducted since 2002. Between 2002 and 2006 participants were randomly assigned to TV/PCIT group or the Waitlist group. Outcomes of these data were

reported in (Thomas & Zimmer-Gembeck, 2011). Between 2007 and 2009, participants were randomly assigned to S/PCIT or Waitlist. Therefore, the Waitlist group was continuous throughout the RCT and randomization to this group occurred between 2002 and 2009. Consequently due to the length of time for allocation, more families were allocated to the Waitlist group compared to the S/PCIT group. Of the 91 families allocated to the Waitlist group, 64 (70%) completed 12-week assessments. Of those families who commenced S/PCIT, 41 (68.3%) completed 12-week S/PCIT assessment. Families randomly assigned to Waitlist were informed they could begin PCIT treatment at the conclusion of a 12-week wait. Figure 1 shows the flow of S/PCIT and Waitlist participants through the study between 2007 and 2009.

### **Procedures**

**Parent-Child Interaction Therapy.** PCIT was developed to improve parenting skills and parent-child interactions among families struggling with their children's (aged 3 to 7) behavior problems (e.g., Oppositional Defiant Disorder; Hembree-Kigin & McNeil, 1995). PCIT has two sequential phases known as Child Directed Interaction (CDI) and Parent Directed Interaction (PDI). Each phase teaches parents communication skills that foster positive parent-child relationships and strategies of differential reinforcement. PCIT skills are taught via didactic presentations to parents and direct coaching of parents while they are interacting with their children. The commencement of each phase includes a didactic session designed to teach the parent specific skills related to each phase of the therapy. The remainder of PCIT involves direct coaching sessions that provide the parent with immediate praise for appropriate responses to their child's behavior and remediation of inappropriate responses. Although the length of treatment time in PCIT is cited as variable, as transition from CDI to PDI occurs when Mastery Criteria for the first phase has been achieved, previous PCIT research reported the average length of treatment time as 13 sessions (Hembree-Kigin & McNeil, 1995; Schuhmann, Foote, Eyberg, Boggs & Algina, 1998).

Treatment is completed when the parent successfully and consistently meets Mastery Criteria by demonstrating strategies learned in PDI and expresses a clear understanding of their own change and role in the family system (Hembree-Kigin & McNeil, 1995). For information on training in PCIT please refer to the PCIT training websites: [www.pcittraining.tv/](http://www.pcittraining.tv/) or <http://www.pcit.org/>.

Participants completed pre-assessment questionnaires (at home between assessment sessions) and the observation task (in the clinic at the second assessment session) prior to random allocation to S/PCIT or Waitlist.

**Standard PCIT.** In previous trials of PCIT (Schuhmann et al., 1998; Thomas & Zimmer-Gembeck, 2011), parents were coached during CDI until Mastery Criteria were achieved for two consecutive sessions before commencing PDI. Treatment concluded when parents met Mastery Criteria for CDI and were observed to effectively implement behavior management strategies taught in PDI (Hembree-Kigin & McNeil, 1995; McNeil, Capage, Bahl, & Blanc, 1999). However in S/PCIT, participants were allocated 12 coaching sessions only regardless of proficiency. On average, S/PCIT participants completed a total of 6.5 CDI sessions ( $SD = 0.78$ , range 5 to 8) and 5.6 PDI sessions ( $SD = 0.97$ , range 4 to 7). In addition to the coaching sessions, all participants received two assessment sessions (pre-assessment and 12-week assessment) and two didactic teaching sessions preceding both CDI and PDI. The average number of PCIT session in total for S/PCIT was 14 ( $SD = 0.84$ , range 12-16). On average, S/PCIT participants who did not complete the intervention had a total of 6.6 sessions ( $SD = 4.7$ ).

**Waitlist.** Participants allocated to the Waitlist were contacted weekly by phone by an allocated PCIT psychologist for brief conversations regarding family and other concerns. Parents in the Waitlist group were asked to refrain from family therapy and therapeutic assistance with child behavior management for the duration of 12 weeks. At the end of 12 weeks, families were offered S/PCIT. Families who commenced S/PCIT after the Waitlist were

not included in the S/PCIT treatment group data of the current study.

**Training and treatment integrity.** Master and Doctoral level psychologists trained in PCIT implemented the intervention. In total, six psychologists (including the first author) provided the intervention. Prior to PCIT all psychologists had experience in providing psychological interventions to adults and children. Supervision was provided by the first author who was trained in PCIT by the PCIT CAARE team in Sacramento USA. A minimum of weekly supervision sessions were provided with additional consults available as necessary. Supervision included observations of PCIT sessions both when requested and at random to assist in PCIT implementation and integrity.

**Data collection.** At the first assessment session, parents were given the self-report and parent-report measures to complete at home. Parents returned forms the following week when they were scheduled to complete the second assessment session, the videotaped pre-assessment. Random allocation to S/PCIT or Waitlist occurred after the videotaped pre-assessment. Post-assessment data were collected after 12 weeks for all participants.

## Measures

**Child externalizing and internalizing symptoms.** Two measures were used to assess children's symptoms; the Eyberg Child Behavior Inventory (ECBI, parent-report) to measure child behavior problems, and the Child Behavior Checklist/4-18 (CBCL; parent-report) to measure externalizing and internalizing symptoms. The ECBI measures the intensity of behavior problems (ECBI Intensity) and the extent parents found the behaviors problematic in children (ECBI Problem). Response options for ECBI Intensity range from 1 (*never*) to 7 (*always*). The sum of these scores indicates the parent perception of the intensity of the child's behaviors. Second, on a dichotomous yes/no scale, the parents endorse whether each behavior is problematic. The sum of the number of endorsed items constitutes a problem scale. Norms for children 2 to 12 years have been established, with the threshold score for clinical problems being

132 for the intensity scale and 15 for the problem scale (Eyberg & Pincus, 1999). The CBCL is a widely used behavioral rating scale for children aged 4 to 18 years and contains 118 items describing a wide range of behavioral and emotional problems (Achenbach, 1991). For children younger than 4 years the Child Behavior Checklist/2-3 was administered. Responders are required to circle the numbers 0 (*not true*), 1 (*somewhat true*), or 2 (*very true*) for each item. Items are categorised into subscales and scores of these items are summed to determine subscale raw scores. Raw scores are converted to *T*-scores, which have a mean of 50 and a *SD* of 10. Hence, the *T*-score was utilised in the current study to enable comparisons between different versions of the CBCL. A *T*-score of 60 represents the lower band of the borderline clinical range (1 *SD* above the mean) with the upper band 63. Scales surpassing a *T*-score of 64 are considered to be in the clinical range. In the current study internal consistency for ECBI Intensity was Cronbach's  $\alpha = .94$ , ECBI Problem .90, CBCL Externalizing .88, and .83 for CBCL Internalizing.

**Parent stress.** The Parenting Stress Inventory (PSI; Abidin, 1990) consists of 101 items that form composite scores for the child and parent stress domains with 90 response options ranging from 0 (*strongly disagree*) to 5 (*strongly agree*) and a further 11 multiple choice. There are a further 19 items that assess specific life stressors. Scores for the PSI are summed for each scale. High scores on the child stress domain indicate that parents believe that they have more difficulty fulfilling their parental role as a result of qualities of their child. High scores on the parent stress domain indicate that the parent's functioning is a significant stressor in the parent-child relationship. Percentiles scores for the child domain scale range from 121 for the 85<sup>th</sup> percentile to 147 for the 99<sup>th</sup> percentile. The parent domain scores range from 160 for the 85<sup>th</sup> percentile to 190 for the 99<sup>th</sup> percentile. Scores for the parent and child stress domains were used in the current study and had good internal consistency ( $\alpha = .93$ , parent domain and .91, child domain).

**Parents' depression.** Maternal depression was measured using the Beck Depression Inventory II (BDI; Beck, Steer & Brown, 1996). The BDI-II is a 21-item self-report measure of depressive symptoms. Scores for multiple choice responses are summed to provide an overall score for depression (0-13 minimal depression, 14-19 mild, 20-28 moderate and 29-63 severe). The internal consistency for the BDI in the current study was  $\alpha = .93$ .

**Parents' child abuse potential.** The Child Abuse Potential Inventory (CAPI; Milner, 1986) was used to measure mothers' level of child abuse potential. The CAPI contains 160 items designed to differentiate maltreating from non-maltreating individuals. Parents are required to circle a dichotomous agree/disagree response option for each item. Items are summed for each subscale, and the abuse scale represents a composite of items from the other scales. The current study used the total Child Abuse score. Using survival analyses comparing CAPI abuse scores and child protection notifications, Chaffin and Valle (2003) reported pre-assessment CAPI abuse scores predicted future child protection notifications for maltreatment with a sample of parents with a history of child maltreatment and those at high-risk. The normative mean for the CAPI abuse scale is 91 and signal detection cut-off is 166. The Cronbach's alpha for the current study was .90.

**Parent observed behaviors.** The Dyadic Parent-Child Interaction Coding System III (DPICS; Eyberg et al., 2004) was used to assess the quality of parents' verbalizations when interacting with their children. The first 5 minutes of a 10 minute child-led videotaped play interaction between parents and children was coded by independent observers. The first 5 minute segment was chosen to be comparable to other published PCIT trials, which have relied on the first 5 minutes of the free-play situation to code DPICS scores. We also considered the first 5 minutes would simulate realistic communication skills rather than skills that increased over time due to parent habituation to the environment. Frequencies of parent verbalizations were tallied and included praises (labeled and unlabeled), descriptions/reflections (combination



of reflections and behavioral and information descriptions), questions (combination of descriptive/reflective questions and information questions) and commands (combination of indirect and direct commands). To adjust for variability in the total number of verbalizations across participants, the percentage of each DPICS category to total verbalizations was calculated for use in analyses.

In addition, the full 10-minute videotaped interactions were coded for sensitivity. The measure of parent sensitivity was developed by modifying one subscale of the Emotional Availability Scales (Biringen, Robinson, & Emde, 2000). Parents were rated from 1 (*highly insensitive*) to 9 (*highly sensitive*). Coding included consideration of the parent's affect, ability to respond to the child's signals, flexibility, and accessibility to the child.

Coders were third or fourth year psychology undergraduate students with no knowledge of PCIT and who were blind to treatment condition. The training regime for each of the observational coding systems included 2-hour blocks of time for 6 weeks. A minimum of 18 hours of training and practice occurred before coding began. Video segments of either DPICS or parent sensitivity were coded by two independent coders. Interrater reliability was assessed using 10 randomly selected video segments. High intraclass correlations were established for the 24 coders for the DPICS categories: praise (.98), reflections/descriptions (.97), questions (.99), commands (.97) and negative talk (.92). In addition, the intraclass correlation for the modified sensitivity scale was .96.

## Results

### Overview of Analyses

For the primary analysis of treatment outcomes, the child and parenting outcomes of families in S/PCIT were compared to Waitlist using 2 (Group: S/PCIT vs. Waitlist) x 2 (time: Pre-assessment vs. 12-week assessment) mixed factorial analysis of variance (ANOVA). Group intervention effect sizes (Cohen's *d*) were also calculated for the Group x Time interactions

(Deville, 2005). Missing data were managed in two ways. First, because a small number of participants (5 or fewer) were missing data for some items on some scales, missing values were imputed using multiple imputation (SPSS version 18) and all analyses were repeated. If all items on a scale were missing, these items were not imputed resulting in a smaller sample size for some analyses. Second, to account for attrition we also conducted Intent-to Treat (ITT) analyses using an imputed data set. ITT was conducted using the Last Observation Carried Forward (LOCF) method of replacing missing data. LOCF was conducted for participants who completed pre-assessment data and were randomized but failed to complete the 12-week assessment. Few differences in data analyses were found when comparing the original data to the imputed data and so all analyses reported are with the imputed data set.

Direct comparisons of S/PCIT and TV/PCIT data were conducted using 2 (group S/PCIT vs. TV/PCIT)  $\times$  2 (time: Pre-assessment vs. follow-up assessment) mixed factorial ANOVA with the imputed ITT data sets for both pre-assessment to 12-week assessment and pre-assessment to treatment completion.

Prior to primary analyses, the adequacy of the randomization was determined by comparing S/PCIT and Waitlist Groups at pre-assessment. Independent *t*-tests (reported in Table 1) revealed no differences between S/PCIT and Waitlist participants on any dependent measure at pre-assessment. Also, no significant differences were found between S/PCIT and Waitlist groups on referral source (child protection, health, education and non-government organization and self referral),  $\chi^2(3, 152) = 0.21, p = .976$ , or when participants were categorized as high (above 166) or low (below 166) abuse potential according to CAPI pre-assessment scores  $\chi^2(1, 150) = 0.04, p = .848$ . In addition, no differences were found between S/PCIT and Waitlist participants who completed the RCT and those who did not. Specifically, there were no differences between participants who completed or chose not to complete in pre-assessment measures for child behavior, parent stress, depression, child abuse potential, and observational

assessment scores. Further, no significant group differences between treatment completers and dropouts were found for child age, parent age, marital status, education level, employment status, or referral source.

On average, parents reported their child's behavior to be within the clinical range for behavior problems: CBCL externalizing, ECBI intensity scale, and ECBI problem scale. Parents reported children to be in the borderline range for internalizing symptoms. In addition, parents also reported child abuse potential well above the normative mean and close to signal detection, high levels of stress pertaining to themselves, their child, and mild depression.

### **S/PCIT Compared to Waitlist**

Overall, from pre- to 12-week assessment, S/PCIT participants showed more improvements than Waitlist participants. Results of the Group  $\times$  Time interactions effects and associated effect sizes are reported in Table 2. Specifically, S/PCIT participants reported greater reductions in their child's externalizing behaviors and internalizing symptoms compared to Waitlist participants with small to medium effects. A small but significant effect for reductions in parent stress attributed to the child was also reported by S/PCIT participants compared to Waitlist participants. However, larger effects were observed for S/PCIT participants compared to Waitlist for praise and descriptions and reflections, and medium to large effects in decreasing questions, commands and negative talk. Also, compared to Waitlist, a significant medium effect was observed for sensitivity, with greater improvement among S/PCIT participants compared to Waitlist. However, there were no significant differences between S/PCIT and Waitlist participants when changes in maternal depression, stress due to parent concerns, or total child abuse potential scores were compared.

### **S/PCIT Compared to TV/PCIT**

**S/PCIT compared to TV/PCIT at pre-assessment.** As done previously, we first determined whether participants differed at pre-assessment prior to conducting the primary

analyses. Independent *t*-tests of the pre-assessment scores of S/PCIT and TV/PCIT participants showed that TV/PCIT participants reported significantly greater internalizing symptoms ( $n = 108$ ,  $M = 60.06$ ,  $SD = 12.32$ ) than S/PCIT participants ( $n = 58$ ,  $M = 54.47$ ,  $SD = 10.1$ ),  $t(164) = 2.96$ ,  $p = .004$ . No differences were found for any other dependent variable (data are not shown in a Table).

**S/PCIT compared to TV/PCIT at pre-assessment to 12-weeks.** Table 3 shows the Group  $\times$  Time interaction effects from mixed factorial ANOVA used to compare differences between S/PCIT and TV/PCIT participants from pre-assessment to the 12-week assessment. Effect sizes are also shown in Table 3. Overall, S/PCIT participants had greater improvements in child externalizing behavior (CBCL) and ECBI problem scale compared to TV/PCIT participants (see Table 3). In addition, although marginally significant and a smaller effect, S/PCIT participants reported fewer child internalizing symptoms after treatment than TV/PCIT participants. With the exception of negative talk, S/PCIT participants outperformed TV/PCIT participants with medium to large effects for improvements in praise and description and reflections, greater reductions in questions and commands, and a significant medium effect for improvements in observed sensitivity.

**S/PCIT compared to TV/PCIT pre-assessment to treatment completion.** Because our previous trial of TV/PCIT used a time-variable format, completion of treatment did not occur within a specified time period. In contrast, all participants in S/PCIT completed treatment after 12 weeks. Therefore we used LOCF to compare S/PCIT participants' completion scores, which were all assessed at 12-weeks ( $n = 59$ ) to TV/PCIT participants' treatment completion scores, which were assessed as early as 10 weeks and as late as 53 weeks after starting treatment ( $n = 99$ , data not shown). There were improvements in all measures between the pre- to the completion assessment, and there were no differences in the improvements over time between S/PCIT and TV/PCIT participants for parent depression,

stress, child abuse potential, child externalizing behaviors, and internalizing symptoms. However, S/PCIT participants showed greater improvements on observational measures, including greater improvements in observed praise,  $F(1, 156) = 4.06, p = .046, d = 0.43$ , and descriptions and reflections,  $F(1, 156) = 8.53, p = .004, d = -0.13$ , compared to TV/PCIT participants. S/PCIT, compared to TV/PCIT participants, also had greater declines in observed questions,  $F(1, 156) = 8.93, p = .003, d = -0.62$ . Further, S/PCIT participants had marginally larger reductions in observed commands than TV/PCIT participants,  $F(1, 156) = 3.83, p = .052, d = -0.31$ . No significant difference between S/PCIT and TV/PCIT participants was found for observed sensitivity.

#### **Attrition Rates of PCIT Participants: Pre- to 12-week Assessment**

A comparison of attrition rates between PCIT groups and the Waitlist group was conducted using  $\chi^2$  analysis (data not shown in a table). Significant differences were found between S/PCIT and TV/PCIT. Of the families who commenced S/PCIT ( $n = 61$ ), 31.6% did not complete treatment compared to 51.4% ( $n = 109$ ) of TV/PCIT participants  $\chi^2(1, 170) = 4.95, p = .036$ . There was no significant difference in attrition rate between S/PCIT and Waitlist participants  $\chi^2(1, 150) = 0.17, p = .722$ .

### **Discussion**

PCIT has a robust evidence-base as an effective intervention for child externalizing behavior, parent stress, and non-optimal parent-child interactions (Eyberg et al., 2001; McNeil et al., 1999). In addition to providing further data for a well-established EBT for child externalizing behavior, this study provides further evidence that PCIT is an EBT for parents who have or are at high risk of maltreating their children. Three independent research cohorts have published data on the effectiveness of PCIT with maltreating parents and their children (Chaffin et al., 2011; Chaffin et al., 2004; Thomas & Zimmer-Gembeck, 2011;

Timmer, Ware et al., 2010). However, the current study is the first to demonstrate effectiveness of Standard PCIT with families at high risk or engaged in maltreatment.

Determining the effectiveness of EBTs with high or special needs subpopulations is an integral part of establishing intervention effectiveness. Evident in the array of modified interventions for specialized populations, it is commonly expected that original interventions require adjustments before implementation with selected populations (Chaffin et al., 2011; McNeil, Herschell, Gurwitsch, & Clemens-Mowrer, 2005; Pincus et al., 2005; Roberts, Mazzucchelli, Studman, & Sanders, 2006; Webster-Stratton & Reid, 2012). In our original RCT of PCIT (Thomas & Zimmer-Gembeck, 2011), we too expected that parents and families with complex problems, such as parents engaged in child maltreatment, would require lengthier treatment options than other families. This was expected because of the evidence that parents who maltreat their children have entrenched maladaptive behaviors, have children who are likely to have very elevated externalizing and internalizing symptoms (Dodge, Bates, & Pettit, 1990), and have experienced transgenerational negative parenting behaviors (Neppl, Conger, Scaramella & Ontai, 2009). Because of this, our original PCIT trial with maltreating parents and their children used a time-variable approach to treatment, whereby therapy concluded when participants had met Mastery Criteria and demonstrated consistent application of behavior management skills (Thomas & Zimmer-Gembeck, 2011). However, a meta-analysis conducted after initiating this trial (Bakersman-Kranenburg et al., 2003) suggested differently. Hence, we were in a unique position to modify PCIT to 12-weeks of coaching only, similar to the original PCIT design (S/PCIT; Hembree-Kigin & McNeil, 1995).

### **Improvements for Families in S/PCIT Compared to Waitlist**

Overall, the findings showed that S/PCIT had positive outcomes for high risk parents and their children when compared to a supported waitlist control. First, S/PCIT participants

reported greater reductions in symptoms and problems from pre- to post-intervention. Reductions were found for child externalizing and internalizing behaviors and stress attributed to the child. S/PCIT participants were also observed to decline in questions, commands and negative talk more than Waitlist participants. Similarly S/PCIT participants were also observed to increase significantly more in positive communication skills and maternal sensitivity than Waitlist.

All parents, whether in S/PCIT or on the Waitlist, reported declines in depressive symptoms between pre-assessment and 12-week assessment. A plausible explanation for improvements to participants' depression scores may be because all participants had weekly contact with a psychologist. Despite non-active intervention by the psychologist during the weekly telephone calls to Waitlist participants, any contact with a professional on a regular basis may be associated with the alleviation of some depressive symptoms.

Although parent stress attributed to the child (PSI Child Domain) decreased over time among intervention families relative to waitlist control, there was no significant decrease in parents' stress due to concerns other than those related to the child. This was measured with the Parent Domain of the PSI, which contains subscales such as isolation and health that are not addressed specifically in S/PCIT. Similarly to parent depression, child abuse potential significantly decreased from pre-assessment to 12-week assessment, however unexpectedly, improvement in these scores did not differ between groups. This contrasts our previous PCIT trial (Thomas & Zimmer-Gembeck, 2011) but is consistent with Chaffin et al., (2004) where decreases in CAPI distress, loneliness and rigidity scale were reported in both groups but no significant treatment group effect was found.

### **S/PCIT Compared to Time Variable PCIT (TV/PCIT)**

To investigate the effectiveness of limiting the PCIT coaching sessions to 12 only with parents who are at high risk or have maltreated their children, we compared the outcomes of

our S/PCIT group to the outcomes for families who participated in TV/PCIT group. Comparisons were possible because of the use of the same measures and consistencies in referral sources and intervention protocols. Compared to TV/PCIT outcomes, S/PCIT treatment outcomes at 12-weeks were either as positive as TV/PCIT outcomes or significantly better. Compared to TV/PCIT, S/PCIT participants had greater reductions in child externalizing behaviors and nominated problem behaviors, and marginally greater reductions in internalizing symptoms and child abuse potential. Similarly, S/PCIT participants outperformed TV/PCIT participants in observed increases in positive communication and reductions in negative communication. These are important differences, suggesting that S/PCIT is at least as beneficial and may be more beneficial than a longer form of PCIT, therefore reducing costs for both families and intervention providers.

### **Potential Practice and Policy Implications**

The identification of effective interventions, which are also time-limited and/or brief, has implications for policy-makers and community practitioners. First, given the high costs of intervention services for children and their families, interventions that demonstrate effectiveness with shorter treatment durations are an incentive for implementation in busy, underfunded community welfare organizations. Practitioners trained in standard protocols of efficacious interventions may not need to modify the intervention for their complex families. This is important for cost-effectiveness of training and implementing evidence-based treatments (EBT). Rather than offering multiple training options for an EBT based on subpopulations (i.e., subpopulation training supplements), EBT training organizations could succinctly deliver their standard protocol training therefore reducing training costs for organizations and simultaneously increasing the likelihood the organization would adopt the EBT. Further, rather than developing subpopulation modifications of EBTs, perhaps policy-



makers should fund large scale trials of efficacious standard protocols for subpopulations before decisions are made regarding whether intervention modifications are required.

Other important factors when providing therapy to high-risk families are attrition and therapy fatigue. Studies have reported that maintaining high-risk families in interventions is difficult (Friars & Mellor, 2007; Timmer, Urquiza, Zebell, & McGrath, 2005). Previous PCIT research in child maltreatment has reported high attrition rates. Within their sample of mothers with a history of maltreating their children, Timmer et al. (2005) reported an attrition rate of 53%. In our TV/PCIT study we had an attrition rate of 57% (Thomas & Zimmer-Gembeck, 2011). In the current study of S/PCIT however, attrition was reduced to 32%. This is similar to a PCIT study comparing two abbreviated versions of PCIT in a subclinical paediatric population (30%; Berkovits et al., 2010).

To handle missing data and attrition we managed the data in two ways. First, missing data were imputed using multiple imputation when analyses included all participants who completed treatment or waitlist follow-up assessments. Second, to maintain participants in analyses when they had not completed treatment, we used the method of LOCF. Because there were two repeated assessments only, this method assumes an individual's score on a measure at follow-up was the same at pre-test. LOCF is a transparent method for managing attrition and missing data, which is more likely to produce a conservative estimate of intervention effect sizes in studies where improvement, rather than deterioration, is expected (Prakash, Risser, & Mallinckrodt, 2008). Hence, we used a combination of methods to manage missing data in order to present good estimates of effects and to be conservative in our estimates when maintaining participants who did not complete treatment in the analyses.

Therapy fatigue (accessing services from too many interventions simultaneously) has been described as contributing to attrition (Kaminski, Valle, Filene, & Boyle, 2008). Reducing the number of treatment sessions and providing a defined intervention end point,

may encourage families to remain in the intervention until completion. When intervention length is open-ended there is potential for both practitioners and families to lose focus of treatment goals. It is possible that participants may not be able to maintain commitment to a variety of intervention approaches simultaneously, and organizations may not have the resources to develop and maintain equally high standards across all intervention components. Open-ended interventions (particularly those with ancillary services) may inadvertently convey messages of complexity of family problems and divert attention from initial presenting problems and goals of treatment (Kaminski et al., 2008). Reviews and meta-analyses have described interventions with fewer treatment sessions as more efficacious than lengthier interventions (Bakersman-Kranenburg et al., 2003; Kaminski et al., 2008). An organized case management approach at initial consultation, with clear goals and foci of treatment, based on family need rather than service centre provision, may assist families to resolve one goal at a time, thereby increasing a sense of empowerment and ability to change. Shortened treatment duration may create a sense of purpose and shared goals between practitioners and their clients resulting in commitment to treatment and focussed requirements of change.

### **Future Research**

A strength of the current study was its diverse sample resulting from the use of inclusive recruitment methods and minimal exclusionary criteria. In addition, few differences were found when pre-assessment levels of dependent variables were compared to the levels reported in previous PCIT studies with families referred solely from child protection authorities. For example, Chaffin et al. (2004, 2011) reported pre-assessment child abuse potential scores and these did not differ when compared to those of the current study participants. Also, Timmer, Ware et al. (2010) reported pre-assessment scores for ECBI intensity and problem scales and CBCL internalizing and externalizing scales. Participants in

the current study had pre-assessment scores higher than Timmer, Ware et al. (2010) for child externalizing behavior (ECBI intensity and problem scales and CBCL externalizing scale) and there were no differences between study participants in internalizing symptoms. However, Standard PCIT was compared to a Waitlist control group in the current study rather than services as usual. Hence, trials are warranted which compare Standard PCIT to treatment as usual or to PCIT augmented with motivational enhancement sessions.

Attrition in our study was approximately 30%, and although this is a significantly smaller proportion than has been reported in other published PCIT trials, it is still higher than ideal. Future research is needed to identify why families do not complete treatment. Also, the timeframe for recruitment of Waitlist participants was longer than for S/PCIT. This may have affected the study findings if there were historical changes in participants and did mean that sample sizes differed between groups. We were not able to compare treatment effects of S/PCIT to a Waitlist at 1-month follow-up, because Waitlist participants were offered treatment after a 12-week wait. However, this issue is not unique to the current study. Waitlist comparison groups are difficult to achieve in research with parents who have maltreated their children due to ethical concerns for a waitlist and sufficient control in an alternate treatment option. However, a supported waitlist comparison group of 12 weeks was included and the current study is the only study that has compared the effects of S/PCIT with families at risk of maltreatment to those on a waitlist. Further research is needed comparing S/PCIT to an active treatment group.

## **Conclusion**

PCIT is an established EBT for child externalizing behavior. In addition, the current study provides data to support Standard 12-week PCIT as an EBT for parents who are at high risk or have maltreated their children. The present findings also provide important information regarding treatment length. Data suggest that adding additional sessions and

ancillary services to extend the time in PCIT may not be necessary for families to experience and report similar or even greater improvements in child behaviors and parent stress attributed to the child, and to be observed to show improvements in parent-child interactions. Although PCIT is identified as an EBT for child maltreatment (Kauffman Best Practices Project, 2004), essential implementation issues such as dosage need further redress to support transportability of PCIT into child welfare organizations before the full benefits of PCIT for families and children can be realized.

### References

- Abidin, R.R. (1990). *Parenting Stress Index* (3<sup>rd</sup> Ed.). Virginia: Pediatric Psychology Press.
- Achenbach, T.M. (1991). *Manual for the Child Behavior Checklist/4-18 and 1991 Profile*. Burlington, VT: University of Vermont.
- Allen, B., Gharagozloo, L., & Johnson, J. C. (2012). Clinician utilization of empirically-supported treatments for maltreated children. *Child Maltreatment, 17*, 10-21. doi: 10.1177/1077559511426333
- Australian Institute of Health and Welfare (2012). Child protection Australia 2010–11. Child Welfare series no. 53. Cat. no. CWS 41. Canberra: AIHW.
- Axford, N., & Little, M. (2009). You can customise it – but just make sure the wheels stay on. *Journal of Children's Services, 4*, 2-3. doi:10.1108/17466660200900001
- Baer, J. C., & Martinez, C. D. (2006). Child maltreatment and insecure attachment: A meta-analysis. *Journal of Reproductive and Infant Psychology, 24*, 187-197. doi: 10.1080/02646830600821231
- Bakermans-Kranenburg, M. J., & van IJzendoorn, M. H., & Juffer, F. (2003). Less is more: Meta-analyses of sensitivity and attachment interventions in early childhood. *Psychological Bulletin, 129*, 195-215. doi:10.1037/0033-2909.129.2.195
- Beck, A. T., Steer, R. A., & Brown, G. K. (1996). *Beck Depression Inventory Manual* (2nd Edition). San Antonio, TX: The Psychological Corporation.
- Berkovits, M. D., O'Brien, K. A., Carter, C. G., & Eyberg, S. M. (2010). Early intervention for behavior problems in primary care: A comparison of two abbreviated versions of Parent-Child Interaction Therapy. *Behavior Therapy, 41*, 375-387. doi:10.1016/j.beth.2009.11.002
- Biringen, Z., Robinson, J. L., & Emde, R. N. (2000). The Emotional Availability Scales, 3<sup>rd</sup> edition. *Attachment and Human Development, 2*, 257-270.

- Braiden, H. J., McDaniel, B., Duffy, J., & McCann, M. (2011). A pilot study of the Incredible Years BASIC Parenting Programme with bereaved families. *Journal of Children's Services, 6.3*, 141-145. doi:10.1108/17466661111176015.
- Bromfield, L., Holzer, P., & Lamont, A. (April 2011). The economic costs of child abuse and neglect. Resource Newsletter. Australian Institute of Families Studies, retrieved from <http://www.aifs.gov.au/nch/pubs/sheets/rs2/rs2.pdf>
- Chaffin, M., Funderburk, B., Bard, D., Valle, L. A., & Gurwich, R. (2011). A combined motivation and Parent-Child Interaction Therapy package reduces child welfare recidivism in a randomized dismantling field trial. *Journal of Consulting and Clinical Psychology, 79*, 84-95. doi:10.1037/a0021227
- Chaffin, M., Silovsky, J. F., Funderburk, B., Valle, L. A., Brestan, E. V., Balachova, T., Jackson, S., Lensgraf, J., & Bonner, B. L. (2004). Parent-Child Interaction Therapy with physically abusive parents: Efficacy for reducing future abuse reports. *Journal of Consulting and Clinical Psychology, 72*, 500-510. doi:10.1037/0022-006X.72.3.500
- Chaffin, M., & Valle, L. A. (2003). Dynamic prediction characteristics of the Child Abuse Potential Inventory. *Child Abuse & Neglect, 27*, 463-481. doi: 10.1016/S0145-2134(03)00036-X
- Chambless, D. L., & Ollendick, T. H. (2001). Empirically supported psychological interventions: Controversies and evidence. *Annual Review of Psychology, 52*, 685-716. doi:0066-4308/01/0201-0685
- Cicchetti, D., & Toth, S. L. (2005). Child maltreatment. *Annual Review of Clinical Psychology, 1*, 409-438. doi:10.1146/annurev.clinpsy.1.102803.144029
- Cicchetti, D., & Valentino, K. (2006). An ecological-transactional perspective on child maltreatment: Failure of the average expectable environment and its influence on child development. In D. Cicchetti & D. Cohen (Eds.), *Developmental psychopathology 2<sup>nd</sup>*

- Edition, Volume three. Risk, disorder, and adaptation* (pp. 129-201). New Jersey: John Wiley & Sons. Inc.
- Devilley, G.J. (2005). ClinTools Software for Windows: Version 4 (computer programme). [www.clintools.com](http://www.clintools.com), Psytek Ltd.
- Dodge, K. A., Bates, J. E., & Pettit, G. S. (1990). Mechanisms in the cycle of violence. *Science*, 250, 1678-1683. doi:10.1126/science.2270481
- Drake, B. (1996). Unraveling “unsubstantiated”. *Child Maltreatment*, 1, 262-271.
- Durlak, J. A., & DuPre, E. P. (2008). Implementation matters: A review of research on the influence of implementation on program outcomes and the factors affecting implementation. *American Journal of Community Psychology*, 41, 327-350  
doi:10.1007/s10464-008-9165-0
- Eyberg, S.M., Duke, M., McDiarmid, M., Boggs, S., Robinson, E., & Washington, E. (2004). *Dyadic Parent-Child Interaction Coding System Third Edition*. Unpublished manuscript. University of Florida.
- Eyberg, S. M., Funderburk, B. W., Hembree-Kigin, T. L., McNeil, C. B., Querido, J. G., & Hood, K. K. (2001). Parent-Child Interaction Therapy with behavior problem children: One and two year maintenance of treatment effects in the family. *Child and Family Behavior Therapy*, 23, 1-19.
- Eyberg, S & Pincus, D. (1999). *Eyberg Child Behavior Inventory and Sutter-Eyberg Student Behavior Inventory-Revised. Professional manual*. Florida: Psychological Assessment Resources.
- Friars, P. M. & Mellor, D. J. (2007). Drop out from behavioral management training programs for ADHD: A prospective study. *Journal of Child and Family Study*, 16, 427-414.  
doi:10.1007/s10826-006-9096-z

- Hembree-Kigin, T. L., & McNeil, C. B. (1995). *Parent-child interaction therapy*. New York: Plenum Press.
- Hussey, J. M., Chang, J. J., & Kotch, J. B., (2006). Child maltreatment in the United States: Prevalence, risk factors, and adolescent health consequences. *Pediatrics*, *118*, 933-942, doi:10.1542/peds.2005-2452
- Hussey, J. M., Marshall, J. M., English, D. J., Knight, E. D., Lau, A. S., Dubowitz, H., Kotch, J. B. (2005). Defining maltreatment according to substantiation: Distinction without a difference? *Child Abuse and Neglect*, *29*, 479-492. doi:10.1016/j.chiabu.2003.12.005
- Kaminski, J. W., Valle, L. A., Filene, J. H., & Boyle, C. L. (2008). A meta-analytic review of components associated with parent training program effectiveness. *Journal of Abnormal Child Psychology*, *36*, 567-589. doi:10.1007/s10802-007-9201-9
- Kauffman Best Practices Project (2004, March). *Closing the Quality Chasm in Child Abuse Treatment: Identifying and Disseminating Best Practices: Findings of the Kauffman Best Practices Project to Help Children Heal From Child Abuse*. Charleston, SC: National Crime Victims Research and Treatment Center.
- Manly, J. T. (2005). Advances in research definitions of child maltreatment. *Child Abuse & Neglect*, *29*, 425-439. doi: 10.1016/j.chiabu.2005.04.001
- McNeil, C. B., Capage, L. C., Bahl, A., & Blanc, H. (1999). Importance of early intervention for disruptive behavior problems: Comparison of treatment and waitlist-control groups. *Early Education and Development*, *10*, 445-454. doi:10.1207/s15566935eed1004\_2
- McNeil, C. B., Herschell, A. D., Gurwitch, R. H., & Clemens-Mowrer, L. (2005). Training foster parents in Parent-Child Interaction Therapy. *Education & Treatment of Children*, *28*, 182-196.
- Milner, J. S. (1986). *The child abuse potential inventory: Manual (2<sup>nd</sup> edition)*. Webster, NC: Psytec.



- Morawska, A., & Sanders, M. R. (2009). An evaluation of a behavioural parenting intervention for parents of gifted children. *Behavior Research and Therapy*, *47*, 463–470. doi: 10.1016/j.brat.2009.02.008
- Neppl, T. K., Conger, R. D., Scaramella, C. V., & Ontai, L. L. (2009). Intergenerational continuity in parenting behavior: Mediating pathways and child effects. *Developmental Psychology*, *45*, 1241-1256. doi:10.1037/a0014850
- Patterson, G. R. (1982). *Coercive family process*. Eugene, OR: Castalia.
- Pincus, D. B., Eyberg, S. M., & Choate, M. L. (2005). Adapting Parent-Child Interaction Therapy for young children with separation anxiety disorder. *Education & Treatment of Children*, *28*, 163-181.
- Prakash, A., Risser, R. C., & Mallinckrodt, C. H. (2008). The impact of analytic method on interpretation of outcomes in longitudinal clinical trials. *International Journal of Clinical Practice*, *62*, 1147-1158.
- Roberts, C., Mazzucchelli, T., Studman, L., & Sanders, M. (2006). Behavioral family intervention for children with developmental disabilities and behavioral problems. *Journal of Clinical Child and Adolescent Psychology*, *35*, 180-193. doi:10.1207/s15374424jccp3502\_2
- Schuhmann, E., Foote, R., Eyberg, S. M., Boggs, S., & Algina, J. (1998). Parent-Child Interaction Therapy: Interim report of a randomized trial with short-term maintenance. *Journal of Clinical Child Psychology*, *27*, 34-35.
- Stangera, C., Ryan, S. R., Hongyun F., & Budney, A. J. (2011). Parent training plus contingency management for substance abusing families: A complier average causal effects (CACE) analysis. *Drug and Alcohol Dependence*, *118*, 119-126. doi: 10.1016/j.drugalcdep.2011.03.007

- Thomas, R., & Zimmer-Gembeck, M. J. (2007). Behavioral outcomes of Parent-Child Interaction Therapy and Triple P – Positive Parenting Program: A review and meta-analysis. *Journal of Abnormal Child Psychology*, *35*, 475-495. doi:10.1007/s10802-007-9104-9
- Thomas, R., & Zimmer-Gembeck, M. J. (2011). Accumulating evidence for Parent-Child Interaction Therapy in the prevention of child maltreatment. *Child Development*, *82*, 177-192. doi: 10.1111/j.1467-8624.2010.01548.x
- Timmer, S. G., Urquiza, A. J., Zebell, N. M., & McGrath, J. M. (2005). Parent-Child Interaction Therapy: Application to maltreating parent-child dyads. *Child Abuse and Neglect*, *29*, 825-842. doi:10.1016/j.chiabu.2005.01.003
- Timmer, S. G., Ware, L. M., Urquiza, A. J., & Zebell, N. M. (2010). The effectiveness of Parent-Child Interaction Therapy for victims of interpersonal violence. *Violence and Victims*, *25*, 486-503. doi:10.1891/0886-6708.25.4.486
- Timmer, S. G., Zebell, N. M., Culver, M. A., & Urquiza, A. J. (2010) Efficacy of adjunct in-home coaching to improve outcomes in Parent–Child Interaction Therapy. *Research on Social Work Practice*, *20*, 36-45. doi:10.1177/1049731509332842
- Webster-Stratton, C. H., & Reid, M. J. (2012). The Incredible Years evidence-based parenting and child programs for families involved in the child welfare system. In A. Rubin (Ed.), *Programs and interventions for maltreated children and families at risk*. (pp. 11-30). New Jersey: John Wiley & Sons. Inc.
- West, F., Sanders, M. R., Cleghorn, G. J., & Savies, P. S. W. (2010). Randomised clinical trial of a family-based lifestyle intervention for childhood obesity. *Behavior Research and Therapy*, *48*, 1170-1179. doi: 10.1016/j.brat.2010.08.008




Figure 1. Flow of S/PCIT and Waitlist participants through study.

Table 1  
*Comparison of Dependent Variables at Pre-assessment Between Standard PCIT and Waitlist*

Measures	Group	N	Pre-assessment		p	95% Confidence Interval of Difference	
			M	SD		Lower	Upper
Child Behavior Problems							
Parent report							
Externalizing behaviors	Standard	58	64.5	9.9	.93	-3.50	3.19
	Waitlist	90	64.3	10.1			
ECBI Intensity	Standard	60	149.8	37.8	.84	-13.40	10.84
	Waitlist	91	148.5	35.2			
ECBI Problem	Standard	60	19.1	8.0	.39	-3.75	1.50
	Waitlist	91	18.0	7.8			
Internalizing symptoms	Standard	58	54.5	10.1	.27	-1.50	5.42
	Waitlist	90	56.4	10.9			
Parent Characteristics							
Parent stress							
Stress due to the child	Standard	60	134.4	25.5	.65	-10.20	6.34
	Waitlist	91	132.5	24.5			
Stress due to the parent	Standard	60	147.7	30.1	.62	-11.87	7.13
	Waitlist	91	145.4	26.8			
Parent child abuse potential	Standard	59	153.9	100.5	.94	-32.39	34.78
	Waitlist	91	155.1	103.2			
Parent depression	Standard	59	14.0	10.5	.57	-2.53	4.62
	Waitlist	91	15.1	11.2			
Parent verbalizations							
Praise, %	Standard	59	3.6	3.3	.83	-1.08	1.35
	Waitlist	81	3.7	3.9			
Descriptions/reflections, %	Standard	59	43.8	11.3	.51	-2.71	5.40
	Waitlist	81	45.1	12.9			
Questions, %	Standard	59	37.3	12.5	.84	-4.89	3.95
	Waitlist	81	36.9	13.9			
Commands, %	Standard	59	13.4	9.2	.74	-3.84	2.72
	Waitlist	81	12.8	10.3			
Negative talk %	Standard	59	1.7	3.3	.42	-1.36	0.57
	Waitlist	81	1.3	2.1			
Parental sensitivity	Standard	59	5.6	1.3	.17	-0.79	0.14
	Waitlist	81	5.2	1.5			

*Note.* ECBI = Eyberg Child Behavior Inventory; Numbers for S/PCIT and Waitlist observations vary due to equipment failure (W = 4 tapeover, 2 no audio) or missing data (W = 4, S/PCIT = 2).

Table 2

*Comparison of Change Between Standard PCIT and Waitlist Pre-assessment to 12-week Assessment*

Measures	Group	N	Pre-assessment		12-week assessment		Group x time F	p	Effect Size d
			M	SD	M	SD			
Child Behavior Problems									
Parent report									
Externalizing behaviors	Standard	57	64.8	9.8	59.0	12.6	13.00	.000	-0.38
	Waitlist	89	64.5	10.1	62.9	11.1			
ECBI Intensity	Standard	60	149.8	37.9	133.7	38.1	5.61	.019	-0.27
	Waitlist	90	149.1	34.9	143.1	36.7			
ECBI Problem	Standard	60	19.1	8.0	13.5	8.6	21.31	.000	-0.61
	Waitlist	90	18.0	7.9	17.5	9.2			
Internalizing symptoms	Standard	57	54.6	10.1	49.8	11.5	6.25	.014	-0.30
	Waitlist	89	56.5	10.9	55.1	12.2			
Parent Characteristics									
Parent stress									
Stress due to the child	Standard	60	134.4	25.5	125.5	36.4	4.27	.041	-0.24
	Waitlist	91	132.5	24.5	130.5	25.8			
Stress due to the parent	Standard	60	147.7	30.1	144.7	37.2	0.29	.591	-0.07
	Waitlist	91	145.4	26.8	144.4	25.9			
Parent child abuse potential	Standard	59	153.9	100.5	137.1	110.7	1.02	.315	-0.01
	Waitlist	91	155.1	103.2	149.1	103.4			
Parent depression	Standard	59	14.0	10.6	12.0	11.26	2.06	.153	0.19
	Waitlist	90	15.1	11.3	11.0	9.88			
Parent verbalizations									
Praise, %	Standard	59	3.6	3.3	12.4	9.3	41.69	.000	1.40
	Waitlist	81	3.7	3.9	4.3	5.1			

Descriptions/reflections, %	Standard	59	43.8	11.3	61.5	12.8	39.22	.000	1.28
	Waitlist	81	45.1	12.9	46.8	12.9			
Questions, %	Standard	59	37.3	12.5	16.7	12.4	59.51	.000	-1.50
	Waitlist	81	36.9	13.9	35.7	13.2			
Commands, %	Standard	59	13.4	9.2	7.9	8.3	5.75	.018	-0.39
	Waitlist	81	12.8	10.3	10.8	8.6			
Negative talk %	Standard	59	1.7	3.3	0.8	1.7	10.19	.002	-0.61
	Waitlist	81	1.3	2.1	1.9	2.9			
Parental sensitivity	Standard	59	5.6	1.3	6.3	1.2	7.16	.008	-0.47
	Waitlist	81	5.3	1.5	5.4	1.4			

*Note.* ECBI = Eyberg Child Behavior Inventory; Numbers for S/PCIT and Waitlist observations vary due to equipment failure (W = 4 tapeover, 2 no audio) or missing data (W = 4, S/PCIT = 2).

Table 3

*Comparisons of Change between Standard and Time Variable PCIT Pre assessment to 12-week Assessment*

Measures	Group	N	Pre-assessment		12-week assessment		Group x time		Effect Size
			M	SD	M	SD	F	p	d
Child Behavior Problems									
Parent report									
Externalizing behaviors	TV/PCIT	107	68.5	10.7	66.1	11.8	10.34	.002	0.31
	S/PCIT	57	64.8	9.8	58.9	12.6			
ECBI Intensity	TV/PCIT	107	162.3	35.6	151.8	37.5	2.80	.096	0.15
	S/PCIT	60	149.8	37.9	133.7	38.1			
ECBI Problem	TV/PCIT	107	20.1	8.3	17.9	9.2	12.42	.001	0.41
	S/PCIT	60	19.1	8.0	13.5	8.6			
Internalizing symptoms	TV/PCIT	107	59.3	11.2	57.2	11.4	3.79	.053	0.24
	S/PCIT	57	54.6	10.1	49.8	11.5			
Parent Characteristics									
Parent stress									
Stress due to the child	TV/PCIT	107	139.5	26.1	138.4	36.5	2.59	.109	0.25
	S/PCIT	60	134.4	25.5	125.5	36.4			
Stress due to the parent	TV/PCIT	107	148.5	35.8	143.7	33.8	0.26	.613	-0.05
	S/PCIT	60	147.7	30.1	144.7	37.2			
Parent child abuse potential	TV/PCIT	107	184.4	102.5	181.8	108.2	3.69	.056	0.13
	S/PCIT	59	153.9	100.5	137.1	110.7			
Parent depression	TV/PCIT	107	15.7	11.2	14.0	11.0	0.08	.775	0.03
	S/PCIT	59	14.0	10.6	12.0	11.3			
Parent verbalizations									
Praise, %	TV/PCIT	99	3.6	4.2	8.5	9.2	7.92	.006	0.55
	S/PCIT	59	3.6	3.3	12.4	9.3			

Descriptions/reflections, %	TV/PCIT	99	48.9	11.9	57.1	14.5	15.82	.000	0.91
	S/PCIT	59	43.8	11.2	63.5	12.8			
Questions, %	TV/PCIT	99	34.6	11.7	24.2	15.2	15.22	.000	0.79
	S/PCIT	59	37.3	12.5	16.7	12.4			
Commands, %	TV/PCIT	99	11.3	8.9	8.9	7.3	6.59	.011	0.37
	S/PCIT	59	13.4	9.2	7.9	8.3			
Negative talk %	TV/PCIT	99	1.6	2.8	1.2	2.6	2.11	.149	0.22
	S/PCIT	59	1.7	3.3	0.8	1.7			
Parental sensitivity	TV/PCIT	99	5.1	1.6	5.4	1.6	4.03	.046	0.34
	S/PCIT	59	5.6	1.3	6.3	1.2			

*Note.* ECBI = Eyberg Child Behavior Inventory; TV/PCIT = Time-variable PCIT; S/PCIT = Standard PCIT; Numbers for TV/PCIT and S/PCIT observations vary due to equipment failure (TV = 2 tapeover, 1 no audio) or missing data (TV = 5, S/PCIT = 2).