Tracking Training Load and Its Implementation in Tactical Populations: A Narrative Review

Maupin, Daniel; Schram, Ben; Orr, Rob Marc

Published in:
Strength and Conditioning Journal

DOI:
[10.1519/SSC.0000000000000492](https://doi.org/10.1519/SSC.0000000000000492)

Licence:
Other

[Link to output in Bond University research repository.](#)

Recommended citation(APA):
Maupin, D., Schram, B., & Orr, R. M. (2019). Tracking Training Load and Its Implementation in Tactical Populations: A Narrative Review. *Strength and Conditioning Journal*, 41(6), 1-11.
<https://doi.org/10.1519/SSC.0000000000000492>

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

For more information, or if you believe that this document breaches copyright, please contact the Bond University research repository coordinator.

Tracking Training Load and Its Implementation in Tactical Populations: A Narrative Review

Danny Maupin D. Pty., ATC

PhD Student, Faculty of Health Sciences and Medicine Bond University

Bond Institute of Health and Sport, Bond University, Gold Coast, QLD, Australia 4216

daniel.maupin@student.bond.edu.au

2 Promethean Way Robina, QLD 4226

Phone: (07) 5595 1111

Fax: (07) 5595 3524

Dr. Ben Schram PhD., D. Pty.

Assistant Professor, Faculty of Health Sciences and Medicine Bond University

Tactical Research Unit, Bond University, Gold Coast, QLD, Australia, 4216

bschram@bond.edu.au

2 Promethean Way Robina, QLD 4226

Phone: (07) 5595 1111

Fax: (07) 5595 3524

Dr. Robin Orr PhD.

Associate Professor, Faculty of Health Sciences and Medicine Bond University

Tactical Research Unit, Bond University, Gold Coast, QLD, Australia, 4216

rorr@bond.edu.au

2 Promethean Way Robina, QLD 4226

Phone: (07) 5595 1111

Fax: (07) 5595 3524

Danny Maupin is a PhD Student in the Faculty of Health Sciences and Medicine at Bond University.

Ben Schram is an Assistant Professor in the Faculty of Health Sciences and Medicine at Bond University.

Robin Orr is an Associate Professor in the Faculty of Health Sciences and Medicine at Bond University.

Key Words: Police, Military, Firefighters, Injury Prevention, Overload

Abstract: Tactical populations often participate in demanding physical training and perform strenuous workplace tasks, increasing injury risk. Mitigating injury risk is vital for maintaining trained personnel and should be a focus for tactical populations. One such method, tracking training load, has not been studied in-depth in tactical populations, despite documented effectiveness in elite sport. Most injuries to tactical personnel are overuse in nature and therefore may be prevented by optimizing training load.

While the methods used in elite sport may not be directly transferrable to tactical environments, they may be used to inform injury mitigation strategies in tactical populations.

Accepted

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Tracking Training Load and Its Implementation in Tactical Populations: A Narrative Review

Key Words: Police, Military, Firefighters, Injury Prevention, Overload

Abstract: Tactical populations often participate in demanding physical training and perform strenuous workplace tasks, increasing injury risk. Mitigating injury risk is vital for maintaining trained personnel and should be a focus for tactical populations. One such method, tracking training load, has not been studied in-depth in tactical populations, despite documented effectiveness in elite sport. Most injuries to tactical personnel are overuse in nature and therefore may be prevented by optimizing training load. While the methods used in elite sport may not be directly transferrable to tactical environments, they may be used to inform injury mitigation strategies in tactical populations.

Introduction

Tactical populations are inclusive of, but not limited to, firefighters, police officers, and military personnel. Despite intermittent sedentary occupational tasks, especially police, (e.g. sitting in a patrol car ¹), tactical populations are required to perform tasks in stressful, and physically demanding situations ¹. This requirement often occurs while these personnel are carrying external loads, such as body armor and other personnel protection equipment ². External loads often vary and can range from 10kg in general police units ², to 22kg in firefighters ³, while military personnel can carry average loads of 45 kg ⁴. The addition of these loads not only causes an increase in metabolic demand ⁵, but can also lead to an increased risk of injury ⁶. In police in particular, these loads further increase upon admission to specialist units ^{7,8}. Specialist police units, due to the requirement to carry additional equipment, such as riot shields and more substantial body armor, can be required to carry loads ranging from around 20kg ⁹ to 40 kg ¹⁰, which can cause even greater metabolic demands and higher risk of injury ^{6,11}.

While each tactical population undergoes high physical stress that increases the risk of injury, the stresses placed on each of these tactical populations vary. Firefighters are typically faced with significant environmental hazards ¹², military personnel are more likely to carry considerably heavier loads ¹³, and police officers are more likely to encounter resistant and uncooperative suspects ¹⁴. Furthermore, and apart from the nature of the occupation itself, sport and physical training can be a leading source of injury in tactical populations ¹⁵. However, sport and physical conditioning are integral to many tactical populations, with appropriate physical training required to increase occupational capability either directly (increased task performance) or indirectly (improved general health) ^{2,16}.

Given the high injury risk of tactical populations ⁶, it is vital that injury mitigation strategies are researched. One such strategy that has not been explored in depth is the monitoring and optimization of training load (TL), despite documented effectiveness in elite sport ¹⁷. The purpose of this narrative review is to explore the potential benefits and limitations of tracking TL in tactical populations with the intent of reducing injury risk while maintaining or increasing fitness. This review will summarize information regarding injuries in tactical populations, and the potential usefulness of optimizing training loads. Information regarding training load, how it can be measured, as well as previous research examining the relationship between training load and injuries, fitness, and performance will also be presented.

Injuries in Tactical Populations

With differing occupational and physical training demands, the location and incidence of injuries experienced can vary; however, injury types are known to be similar across these populations ¹⁸. In a study by Taylor et al. ¹⁹ it was found that firefighters suffered injuries at a rate of 177 per 1000 full-time employees per annum. Firefighters were most likely to suffer joint sprains and muscle strains, accounting for 66.5% of all injuries, followed by contusion/crush injuries, mental disorders, and open wounds. The most common sites of injury being the knee, lower back, shoulder and ankle ¹⁹.

In law enforcement, a critical review ¹⁸ found injury rates ranging from 240 to 2500 per 1000 personnel per annum, with multiple articles reviewed reporting the most common region of injury being the upper extremity (32.95% to 43.42% of injuries), while another stated the most common site was Other Unspecified sites (63.41%), followed by torso and back (20.49%). This review found the most common injury type to be sprains and strains (42.36% to 94.59%), followed by Other Muscle pains and Other Natures of injury ¹⁸.

With regards to military, incidence rates have been reported to range from 23 incidents, for active duty personnel, to 34 incidents, for reserve personnel, per 100 person-years of active service, with “minor personal injury” being the most common for both groups²⁰. Their injuries were found to most commonly be overuse or stress syndrome, muscle strains, ankle sprains, and stress fractures which typically occurred at or below the knee²¹.

Injuries in these populations can have serious downstream effects, such as a temporary or permanent loss of experience and skills, or even worse, higher workforce strain due to the requirement of covering injured personnel^{22,23}. Injuries to tactical populations can also result in high monetary costs to treat and, if needed, replace (e.g. temporarily or permanently) injured personnel, placing an increased financial burden on the organization^{22,23}. For example, the financial cost for medical care and salaries of soldiers in the United States (U.S) Army is estimated to be about \$1.5 billion (USD) per year²³. By exploring causative factors for these injuries, and implementing programs to reduce these injuries, some of these negative effects may be mitigated.

Numerous causative factors have been studied in tactical populations, such as poor metabolic fitness levels²⁴, high body mass index (BMI)²⁵, history of previous injury²⁶, and smoking²⁷. In 2000, Kaufman et al.²¹ discussed how training load (TL) could potentially be another causative factor, and how optimizing TL may decrease injury risk while promoting fitness in a military population. Previous research has shown that an excessive increase in physical activity is linked with an increased injury risk^{26,28}. In a study by Trank et al.²⁸, it was found that U.S. Navy Recruits who completed the highest running mileage (>25 miles (40.23 km)) also had a significantly higher risk of injury, with no significant change in 1.5 mile (2.41 km) run performance. While the need to track TL is evident, a cursory search of the literature shows that little research has been performed to date with regards to quantifying TL in tactical populations. In fact, a systematic review found only two articles regarding TL tactical populations, but these articles focused mainly on increases in physical activity²⁹. Despite minimal information pertaining to the tactical

environment, it has become a popular tool in elite sporting teams. Similar to the study by Trank et al.²⁸, a medium effect size ($r = 0.59$; CI and p not reported) was found between higher running distances during a given month in endurance runners and number of injury days the following month³⁰. By implementing methods that track TL, sporting teams are able to adjust training stimuli at an individual level, and on a weekly basis to reduce injury risk and also increase performance¹⁷. Given the significant costs associated with injuries, and the increased risk of injury in tactical populations caused by higher physical loads, a means of monitoring TL and their associated effects on injury risk could prove beneficial for tactical populations^{26,28}.

Training Load

The International Olympic Committee consensus statement defines TL as a cumulative amount of stress on an individual from single, or multiple, training sessions over time³¹. TL is typically measured using either external methods (e.g. total distance run, or speed) or internal methods (e.g. heart rate or rating of perceived exertion)¹⁷. External load (EL) has previously been defined as “any external stimulus applied to the athlete that is measured independently of their internal characteristics”, while internal load (IL) is “load measurable by assessing internal response factors within the biological system, which may be physiological, psychological, or other”³¹. Tracking EL can communicate the total amount of work completed, while information yielded from IL typically conveys how the individual is adapting physiologically³¹. Emphasis is placed upon the importance of utilizing both measures, as together they provide a more comprehensive view of the physical stress an individual is experiencing¹⁷. For example, an individual may perform the same overall output (as measured by EL, such as distance) on two different days, however their ability to respond to this output (as measured by IL, such as heart rate) may have changed. Potential differences between responses to training may provide insights as to

whether the individual is in a state of 'readiness' and able to tolerate high TL or 'fatigue' and potentially at an increased risk of both injury and decreased performance ³².

Rapid changes in both EL and IL have previously been linked to higher risk of injury in the literature ¹⁷.

One example shows 40% of injuries were the result of a rapid change in TL in the preceding week in Australian football ³³. Factors further compounding the issue were proposed by Gabbett that athletes who over-train or under-train are at higher injury risks ¹⁷. These issues have led to the development of monitoring both IL and EL over the short-term (acute, typically 1 week) and long-term (chronic, 3-6 weeks) periods, and weekly changes to track overall and rapid changes in workload ¹⁷. There are various methods of measuring TL reported in the literature, each of which have benefits and limitations; the following sections will explore these various methods.

External Measurements of Training Load

EL is measured in an attempt to define the total work completed ³¹. EL measurements can include distance run, tracking volume-load during strength training sessions, and number of sprints completed, among others ¹⁷. Measuring EL has been aided with the advent of Global Positioning System (GPS) devices and the subsequent accessibility of these relatively low cost devices, which allow for accurate measurement of the above variables ³⁴. The use of GPS to measure EL is common practice in many professional sports including Australian Football League (AFL), rugby league, and soccer ³⁴. Research has shown the effectiveness and validity of GPS to measure EL in these sports ^{35,36}. However, there are concerns of the precision of GPS with regards to high rates of change in velocity and while GPS technology has seen an increase in data accuracy, there are still some limitations in regards to very high speed running (>20 km/hr) ³⁷. While there may be potential validity issues at high speed, has been estimated that the speed of military marches while on level ground is 5.3 km/h ³⁸, which may allow GPS

1
2
3
4 138 devices to accurately record data. Practitioners and clinicians may be able to utilize these devices
5
6 139 effectively during training stations and while personnel are on duty and be able to record distance
7
8
9 140 covered.

10
11
12 141 Another concern with GPS units is the high cost associated with them ³⁹. Research has shown that
13
14 142 utilizing GPS on a cellular phone may provide an alternate way to measure distance, as the accuracy
15
16
17 143 between the two was comparable with an approximately 3% variation identified when measuring
18
19 144 distance and average speed ⁴⁰. Of importance, this study compared the cellular device to GPS devices
20
21 145 with lower sampling rates and more limited accuracy, therefore more variation may exist when
22
23 146 comparing cellular devices to GPS devices with higher sampling rates ⁴⁰. Utilizing cellular phones or
24
25
26 147 fitness watches may be an alternative tool for tactical practitioners to pursue if the acquisition of GPS
27
28 148 units proves too costly. One key area of difference between tactical populations and athletics may be
29
30 149 potential security concerns and GPS units. Recently data collected by a wearable GPS unit showed the
31
32 150 location of a secure U.S. military facility as discussed in an article published by Liz Sly in the Washington
33
34
35 151 Post on January 29th, 2018. Tactical personnel will need to ensure that the devices they are using are
36
37
38 152 secure and will not lead to similar issues.

39
40
41 153 Apart from GPS devices to track EL, there are other, more cost-effective methods. One such method is
42
43 154 utilizing volume-load per the NSCA's definition (number of repetitions x external load [kg]), a convenient
44
45 155 method as it does not require any additional equipment ⁴¹. Whereas repetition-volume is the total
46
47 156 number of repetitions performed during a workout, volume-load also incorporates the weight lifted and
48
49
50 157 is likely better at quantifying the total amount of work completed ⁴¹. For example, if a police officer
51
52 158 completes 4 sets of 12 repetitions of bench press at 100 kg, their volume load will be 4800 "work units",
53
54
55 159 while their repetition-volume will be 48. Volume-load can be used during physical training sessions, when
56
57
58 160 tactical personnel are participating in strength training, to track and adjust based on previous and future
59
60 161 loads. More complex methods have that determine mechanical work during resistance exercises may be
61
62
63
64
65

utilized during strength sessions; however these require additional devices, such as force plates, accelerometers, and inertial sensors³⁹. While these devices have been shown to have good reliability and validity, they are expensive methods to implement and may not be accessible in all populations³⁹. Another method track TL that may be of benefit when training large groups include measuring total running distance, or weight lifted utilizing subjective measures³⁹. This method may be particularly useful in military populations upon return from patrol, due to their ability and operational requirement in tracking distances covered. In support of subjective measures, a systematic review on endurance athletes³⁰ on endurance runners found a medium effect size ($r=0.59$; CI and p not reported)⁴² ($r^2=0.36$, CI and $p=0.001$)⁴³ between subjective reports of high total training distance and rates of injury or pain. Lastly, there are other methods which have been used to track TL, though these are unlikely to be used in tactical populations due to logistical concerns and time constraints³⁹. For example, despite time motion analysis being found to be effective in tracking fatigue in professional soccer players⁴⁴, it is unlikely that tactical populations would have the time or resources to record and perform film review of all training and occupational settings. Similarly, despite training diaries being used to track load, concern exists with regards to the accuracy of self-reporting⁴⁵; especially if trying to recall events that occurred at the beginning of long hours of training⁴⁶.

Internal Measurements of Training Load

While EL measurements attempt to measure the amount of work done, IL measurements are more focused on the individual's response to the TL¹⁷. Numerous methods to track IL including heart rate (HR), session Rating of Perceived Exertion (sRPE), and questionnaires. Historically heart rate (HR) has been used to track IL due to an almost linear relationship with VO_{2max} ⁴⁷. This allows practitioners to quantify how each individual is responding to the imposed work load⁴⁸. Likewise, blood lactate is

related to VO_{2Max} and may be used to track training load and intensity⁴⁹. As this measure has also been linked to load carriage performance in soldiers⁵⁰, it may be a useful tool to track intensity and performance. While these methods are accurate, they can be invasive and may require a high level of technical skill or the necessary technology to be performed, and therefore may not be feasible in a tactical population. As an example, while measuring HR at one set point is a simple task, requiring only a stopwatch, measuring and recording HR consistently over a period time may require more sophisticated technology, such as HR chest straps³⁹. Utilizing fitness watches to measure heart rate may be an alternative; however, research has shown that these devices are less accurate at higher heart rates compared to chest straps⁴⁷. Incorporating the Rating of Perceived Exertion (RPE) scale may be useful, as it is both low cost and easy to administer, especially to larger groups⁴⁸.

The RPE scale (Figure 1) was designed by Borg⁵¹, as an attempt to measure the level of physical strain during activity. It was proposed by Foster et al.⁵², that the this scale can be modified to a 0-10 scale (Figure 2), and a score given at the end of activity. This score can then be multiplied by the duration of the session for a measure of TL, called session RPE or sRPE⁵². For example, if an athlete completed a 60-minute weight lifting session, and reported the session was a 6 on the modified RPE, the athlete would have a sRPE of 360 (60 x 6) arbitrary units (AU). Previous research has shown that this method correlates well with blood lactate concentration and heart rate after exercise^{53,54}. In a study by Gabbet et al.⁵³ the correlations between sRPE and both heart rate and blood lactate concentration were found to be $r=0.89$ and $r=0.86$ respectively, showing a high correlation between measures. These results suggest sRPE, using the modified scale, can be used to accurately measure IL by tactical practitioners after a variety of activities, such as physical training sessions. While tactical personnel do tend to perform physical activity throughout the day, previous research has confined the “session duration” to fixed training hours experienced each day (0600 – 2200), allowing comparison of RPE scores across days⁵⁵. There was good agreement between RPE and recorded training impulse (TRIMP) scores ($R^2= 0.57-0.77$), showing that this

may be a valid method to compare RPE scores in tactical populations though further research will be necessary⁵⁵.

Training impulse (TRIMP) was developed by Bannister et al.⁵⁶ as a method of quantifying a training session. This method is calculated by using training duration, maximal heart rate, resting heart rate, average heart rate, and a sex-dependent exponential coefficient⁵⁶. This equation was further modified by Manzi et al.⁵⁷, by introducing an individual based exponential factor. The modification was done in attempts to better determine individual physiological response to exercise⁵⁷. The use of this equation, TRIMP_i, has been shown to be a valid method to track fitness and performance changes in long distance runners⁵⁷ as well as aerobic fitness in team sport⁵⁸. Another method of calculating TRIMP, Edward's TRIMP, multiplies the duration accumulated in five HR zones (zone 1 = 50-60% HR_{Max}, zone 2 = 60-70%, zone 3 = 70-80%, zone 4 = 80-90%, and zone 5 = 90-100%) by a coefficient for each zone and then sums the result⁵⁹. This equation was utilized in military training and found agreement ($R^2 = 0.34-0.49$) between TRIMP and distance, suggesting this may be a valid method to calculate internal load, though it may be limited due to the variable training activities and prolonged time frames found in military populations⁵⁵. Utilizing TRIMP may be an effective means track IL over time in tactical populations, particularly Edward's TRIMP as an individual factor will not need to be calculated and it accounts for activity completed in various HR zones.

Heart rate variability (HRV), the beat-to-beat fluctuation of resting HR, is another measure of IL⁶⁰. The autonomic nervous system (ANS) plays a crucial and complex role in maintaining the body's homeostasis⁶⁰. While the ANS is involved in many different mechanisms, its ability to control HR and role in overuse injuries is a key factor when it comes to monitoring TL⁶⁰. Examining the physiology of this relationship is beyond the scope of this article, but to paraphrase, using the parasympathetic and sympathetic nervous systems, the ANS is able to increase and decrease HR, and release chemical mediators in response to injuries, including overuse injuries⁶¹. HRV aims to indirectly measure the ANS by analyzing the change in

time between heart beats during resting HR ⁶⁰. Significant changes from an individual's baseline may show that the ANS is under stress, potentially due to overtraining or the onset of an overuse injury ⁶⁰. Given the many roles of ANS, changes in HRV may be indicative of other types of stress than overuse injuries. However, this may be beneficial to those working with tactical populations given the amount of non-physical stress, such as psychological, that are experienced ^{62,63}. Currently, research is inconclusive about the relationship between HRV and overtraining, with evidence both for and against ⁶³⁻⁶⁵, but this may be a valuable tool to measure the amount of stress, both physical and other, that are being experienced by tactical personnel.

Various wellness questionnaires, which examine factors such as sleep quantity and sleep quality, feelings of fatigue and soreness, have also been used in an attempt to measure IL ³⁹. One such survey, the Recovery-Stress Questionnaire for Athletes (RSQ), was created to identify the extent of an athlete's physical and mental exhaustion and recoverability ⁶⁶. This questionnaire consists of 19 subscales that discuss various topics such as general and emotional stress, burnout, physical complaints and physical recovery among others ⁶⁶. The survey is periodically given out, with one study administering the survey four times during a three week period ⁶⁷. It has previously been shown to be able to track the general parameters of training stress of athletes across multiple sports ⁶⁶, though it is unknown how this scale will apply to those in tactical populations or if they will require a more specific questionnaire for their demands. However, given the high amounts of psychological stress experienced by tactical populations ⁶², the RSQ may be provide useful information regarding not only the physical, but emotional state of personnel.

Another questionnaire that attempts to quantify recovery is the total quality recovery process (TQR) ⁶⁸. This process is divided into two parts, the first being a perceived recovery scale (TQR perceived) (Figure

1). Structured like the Borg RPE Scale, this 6-20 scale attempts to measure an individual's perceived recovery at the end of the day⁶⁸. The second aspect of TQR (TQR action) is a more objective measure of recovery where individuals earn recovery points (20 points maximum) based on nutrition and hydration (10 points maximum), sleep and rest (4 points), and relaxation and emotional support (3 points) (Figure 3)^{68,69}. As this method combines both perceived and an objective score for recovery, it may be useful in tactical populations, though future research will be needed to validate this process.

Insert Figure 1 Here

Insert Figure 2 Here

Insert Figure 3 Here

Lastly, while the above methods may be more complex, previous research in military populations has examined muscle soreness and physical fatigue using 0-10 scales⁵⁵. While this study, did not examine these measures in relation to injury, these simple questions may be able track how tactical personnel are managing their current TL⁵⁵. Various questionnaires exist that track both the physical and psychological stress and recovery experienced by individuals^{66,68}. Tactical practitioners can utilize these measures periodically to track their personnel. If measures such as these prove to be too time-consuming implementing muscle soreness and physical fatigue scales may be an alternative method to track IL⁵⁵.

Insert Table 1 Here

Acute:Chronic Workload Ratio

Acute:Chronic Workload Ratio (ACWR) is an attempt to provide a more complete picture of an individual's cumulative load by comparing an acute workload to a chronic workload and quantifying the

changes over time¹⁷. As mentioned above, it may be the rate of change in load, more than the absolute load, that relates to injury risk³¹. The importance of rate of change in load can be historically seen through the 10% rule which states that increases in exercise frequency, duration, or intensity should be limited to 10% of the previous week⁷⁰. While the timeframes may vary, in sports the acute workload is typically represented by one week, while the chronic load is a summation of the previous 3-6 weeks¹⁷. If the acute workload is greater than the chronic workload (signifying increases in training load), the ratio will be greater than one, and the individual will be in a fatigued state¹⁷. Conversely, if the acute workload is less than chronic workload, the ratio will be less than one, and injury may be less likely however performance may plateau or decline with inadequate stimulus¹⁷. The chronic workload has also been called 'fitness' in the literature, while the acute workload has been called 'fatigue'¹⁷. Evidence shows that high chronic workload or 'fitness', may protect against spikes in acute workload or 'fatigue'^{17,71,72}. In a study by Malone et al., it was found that a high chronic TL (4750 AU) measured by sRPE protected against injury from maximal velocity running⁷¹. In another study by Hulin et al.⁷², it was found that a high chronic workload (>16095 meters covered) combined with a moderate ACWR (1.03-1.38; Relative Risk (RR)=6.2±2.2) resulted in a lower injury risk than low chronic workload (<16095 meters covered) combined with a moderate ACWR (1.03-1.38; RR=9.3±2.2). RR states how many times more likely an outcome will occur in an exposed group compared to a non-exposure group⁷³. In the case of the Hulin et al. article⁷², those who experienced a high chronic workload were 6.2 times more likely to experience an injury, while those in the low chronic workload were 9.3 times more likely to experience an injury. ACWR may be more appropriate to utilize than the 10% rule, as it is not a rigid law, but can be adjusted based on past training.

One limitation of the ACWR model is that it does not account for the decaying effects of fatigue and fitness over time, and therefore may be able to be improved^{39,74}. A new method utilizing exponential weighted moving averages (EWMA), instead of rolling averages, was recently put forward by Williams et

al⁷⁵. This method attempts to account for the decaying effects by assigning a decreasing weighting for older values⁷⁵. A study by Murray et al.⁷⁴ compared EWMA ACWR to rolling average ACWR and found that, while both models showed a significant increase in injury risk with high ACWR, EWMA was more sensitive to detect increases in injury likelihood at higher ACWRs. When compared with an ACWR of 1.0-1.49, a rolling average ACWR of 2.0 was shown to have a relative injury risk of 6.52 [4.83-8.80], a statistically significant finding. Utilizing the same comparison, but with EWMA, the relative risk increased to 21.28 [20.02-22.62] - demonstrating a much higher relative injury risk. Incorporating an EWMA ACWR over a rolling average ACWR may be a more sensitive and advantageous method⁷⁵.

Clinicians and practitioners can utilize ACWR with any of the above-mentioned variables. By incorporating ACWR with measures such as a total distance, SRPE, or volume load, practitioners may be better able to predict how well their personnel will respond to future programs. If knowing that over the last 3 weeks, personnel have only covered an average of 2000m per week, practitioners can calculate that covering 4000m will result in an ACWR of 2.00 and therefore increase injury risk (acute load divided by chronic load, or 4000m divided by 2000m). Practitioners can adjust their scheduled training sessions either based off previous work load or off current occupational demands (e.g. dismounted patrols) to keep personnel within a safe range.

While ACWR may be a useful tool for predicting injury risk, there are potential limitations affecting its validity. One such issue is the potential spurious correlations that may exist between ACWR and injury risk⁷⁶. This is likely due to the fact that the acute load is used in the calculation of the chronic load, as the acute load is on both sides of the equation⁷⁶. A second impacting factor is the confounding effect training schedule has on the ability of ACWR to predict injury risk⁷⁷. This was calculated with regards to sporting teams, so it remains to be seen if this relationship exists in tactical population. However, though ACWR may be an effective method to calculate injury risk, it should not be solely relied upon by clinicians and practitioners.

The Association of Training Load and Injuries

Research has hypothesized that musculoskeletal injuries may be linked to adverse responses in TL⁶⁰, while specifically researched targets regarding TL and injury in the athletic domain have been studied. Research has shown that two week cumulative sRPE loads >3700 AU⁷⁸, and high three week running distances (73721 – 86662 meters)³⁶ are likely to increase injury risk in Australian Rules football players while studies across multiple sports have shown that ACWR ratios ranging from 0.8 – 1.3 may minimize injury risk while ensuring increases in fitness^{35,72}. Utilizing either cumulative loads or ACWR provides information to medical staff and strength and conditioning (S&C) coaches which may assist with reducing injury risk and maximizing performance in tactical environments. It is unlikely that these numbers will be directly transferable given the differences that exist between the tactical and athletic domain, though it should be noted that members of tactical population regularly play sport^{79,80}.

The Associations of Training Load and Fitness and Performance

While TL may assist with mitigating injury risk, it also is related to fitness and performance. Fitness is a key requirement for tactical personnel and is associated with successful occupational task performance², reduced injury risk,⁸¹ and mental health benefits¹⁶. However, while training to increase fitness is needed, it may pose a risk if not monitored correctly, as supported by findings that in military personnel, physical training and sports participation are leading causes of injuries¹⁵. Tracking TL may be able to ensure that tactical populations are training at a high enough intensity to increase fitness while not unnecessarily increasing risk of injury¹⁷. In AFL players, it was found that those who had completed high TL (1600-2000 AU per week) increased their 2km time trial compared to those who completed very high (>2000 AU per week), moderate (1250-1599 AU per week), and low (<1250 AU per week) TL as measured by sRPE⁷⁸. In a study by Gabbett et al.⁵³, it was found that while increases in TL of 175 to 620

1
2
3
4 350 AU, as measured by sRPE, did not result in any increase risk of injury, it did result in decreased
5
6 351 performances in agility. These results provide evidence that tracking TL cannot only influence injury risk,
7
8 352 but fitness and performance as well. Given the inherent dangers tactical populations can encounter ^{7,82},
9
10 353 decreases in performance or losses of fitness do not simply result in a lost game, but can have life and
11
12 354 death consequences. By implementing a method of monitoring TL tactical populations can aim for, and
13
14 355 train at a level that maximizes performance gains while minimizing injury risk ^{39,83}.
15
16
17
18
19 356 While there are notable differences between tactical populations and elite athletes, the TL methodology
20
21 357 used in elite sport may provide a template for tactical personnel. Preliminary studies in military
22
23 358 populations show the associated increase in injury risk with increase in physical activity ^{26,28}. While these
24
25 359 studies have shown the potential harmful impacts of excessive physical activity, modern methods such
26
27 360 as sRPE, ACWR or any of the other various methods to track EL and IL used in elite sports ³⁹, appear to be
28
29 361 suitable to implement to track and adjust TL ¹⁷.
30
31
32
33
34 362 Practitioners must recognize that excessive TL is just one potential causative factor of injury in tactical
35
36 363 populations. While there are various other causative factors, i.e. smoking ^{27,81}, previous injury ^{26,27}, and
37
38 364 poor fitness ⁸¹, little research has been performed regarding monitoring TL, especially using the methods
39
40 365 seen in elite sport ^{17,39}. The current depth of research is lacking given the impact overloading or
41
42 366 underloading has on injuries in tactical populations. Multiple studies have shown the impact of high
43
44 367 physical activity and repetitive microtrauma has on musculoskeletal injuries in tactical populations ^{28,84}.
45
46
47
48 368 A large amount of these injuries are due to repetitive microtrauma, or overuse, accounting for almost 4
49
50 369 times as many injuries as acute trauma ⁸⁴. Military recruits especially are at risk for injury due to sudden
51
52
53 370 and large increases in TL due to high amounts of running and other physical activity with possibly little to
54
55 371 no prior training ^{85,86}. Given the impact injuries have on both institutional costs, approximately \$1.5
56
57 372 billion (USD) per year in United States Army ^{22,23}, and organizational workload, it is vital that tactical
58
59 373 populations begin to research and implement these practices.
60
61
62
63
64
65

Conclusion

Given both the high physical workload and injury rate within tactical populations, it is important that steps are put in to place to maximize performance in decrease injury risk. A wide range of variables including HR, sRPE, HRV, distance covered, volume-load among others may allow practitioners to monitor TL within their personnel. In addition, ACWRs could be devised to minimize any spikes in training load which may increase injury risk. A comprehensive understanding of TL may allow for optimizing the programming of activities such as pack marches and patrols, to allow for the addition of, or reduction of training where required. For example, if a military unit is planning to complete a 5 km dismounted patrol in 4 weeks, a S&C coach can analyze the distance covered previously during field and physical training sessions. Due to the minimal research performed within the tactical domain to date, specific ratios are not currently available, however evidence from elite sport highlights that ACWRs from 0.8-1.3 may be optimal for both performance gains and injury reduction.

Figure 1. Borg RPE Scale and Total Quality Recovery Scale. Reproduced from Kentta and Hassmen (48)

Figure 2. Modified Borg RPE Scale. Reproduced from Foster et al. (25)

Figure 3. TQR Action Scoring Guide. Reproduced from Kentta and Hassmen (49)

References

1. Dawes JJ, Orr RM, Flores RR, Lockie RG, Kornhauser C, Holmes R. A physical fitness profile of state highway patrol officers by gender and age. *Ann Occup Environ Med.* 2017;29.
2. Robinson J, Roberts A, Irving S, Orr R. Aerobic fitness is of greater importance than strength and power in the load carriage performance of specialist police. *Int J Exerc Sci.* 2018;11(4):987-998.
3. Carlton A, Gorey R, Orr R. The Impact of Suppressing a Structural Fire on Firefighter Hydration. *J Aust Strength Cond.* 2016;24(5):27-33.

4. Orr R, Pope R, Coyle J, Johnston V. Occupational loads carried by Australian soldiers on military operations. *J Health Saf Environ*. 2015;31(1):451-467.
5. Divencenzo HR, Morgan AL, Laurent CM, Keylock KT. Metabolic demands of law enforcement personal protective equipment during exercise tasks. *Ergonomics*. 2014;57(11):1-6.
6. Knapik JJ, Reynolds KL, Harman E. Soldier load carriage: historical, physiological, biomechanical, and medical aspects. *Mil Med*. 2004;169(1):45.
7. Pryor RR, Colburn D, Crill MT, Hostler DP, Suyama J. Fitness characteristics of a suburban special weapons and tactics team. *J Strength Cond Res*. 2012;26(3):752.
8. Hunt AP, Orr RM, Billing DC. Developing physical capability standards that are predictive of success on Special Forces selection courses. *Mil Med*. 2013;178(6):619.
9. Carbone PD, Carlton SD, Stierli M, Orr R. The impact of load carriage on the marksmanship of the tactical police officer: a pilot study. *J Aust Strength Cond*. 2014;22(2):50-57.
10. Keeler JM. *The Effect of Tactical Tasks and Gear on Muscle Activation of SWAT Officers* [Master of Science]: Theses and Dissertations--Kinesiology and Health Promotion, University of Kentucky; 2014.
11. Drain J, Orr R, Attwells R, Billing D. *Load Carriage Capacity of the Dismounted Combatant-A Commanders' Guide*. Defence Science and Technology Organisation Victoria (Australia) Human Protection and Performance Div;2012.
12. Rossi R. Fire fighting and its influence on the body. *Ergonomics*. 2003;46(10):1017-1033.
13. Orr R, Pope R, Johnston V, Coyle J. Soldier load carriage: An investigation into the load carriage conditioning practices of the Australian Regular Army. Paper presented at: Australian Military Medicine Association 2012.
14. Brandl SG, Stroshine MS. The Physical Hazards of Police Work Revisited. *Police Q*. 2012;15(3):262-282.
15. Jones B, Knapik J. Physical Training and Exercise-Related Injuries. *Sports Med*. 1999;27(2):111-125.
16. Crowley SK, Wilkinson LL, Wigfall LT, et al. Physical fitness and depressive symptoms during army basic combat training. *Med Sci Sports Exerc*. 2015;47(1):151-158.
17. Gabbett TJ. The training-injury prevention paradox: should athletes be training smarter and harder? *Br J Sports Med*. 2016;50(5).
18. Lyons K, Radburn C, Orr R, Pope R. A profile of injuries sustained by Law Enforcement Officers: A Critical Review. *Int J Environ Res Public Health*. 2017.
19. Taylor NA, Dodd MJ, Taylor EA, Donohoe AM. A retrospective evaluation of injuries to Australian urban firefighters (2003 to 2012): injury types, locations, and causal mechanisms. *J Occup Environ Med*. 2015;57(7):757-764.
20. Pope R, Orr R, MacDonald D. Incidence rates of reported work health and safety incidents and injuries in part-time and full-time Australian Army personnel. *J Mil Veterans Health*. 2015;23(4):45-46.
21. Kaufman KR, Brodine S, Shaffer R. Military training- related injuries: Surveillance, research, and prevention. *Am J Prev Med*. 2000;18(3):54-63.
22. Hilyer JC, Brown KC, Sirles AT, Peoples L. A flexibility intervention to reduce the incidence and severity of joint injuries among municipal firefighters. *J Occup Med*. 1990;32(7):631-637.
23. Nindl BC, Williams TJ, Deuster PA, Butler NL, Jones BH. Strategies for optimizing military physical readiness and preventing musculoskeletal injuries in the 21st century. *US Army Med Dep J*. 2013:5-23.
24. Lisan P, Connor FG, Deuster PA, Knapik JJ. Functional movement screen and aerobic fitness predict injuries in military training. *Med Sci Sports Exerc*. 2013;45(4):636.

- 1
2
3
4 445 25. Robinson M, Siddall A, Bilzon J, et al. Low fitness, low body mass and prior injury predict injury
5 446 risk during military recruit training: a prospective cohort study in the British Army. *BMJ Open*
6 447 *Sport Exerc Med.* 2016;2(1).
7
8 448 26. Almeida SA, Williams KM, Shaffer RA, Brodine SK. Epidemiological patterns of musculoskeletal
9 449 injuries and physical training. *Med Sci Sports Exerc.* 1999;31(8):1176.
10 450 27. Bulzacchelli MT, Sulsky SI, Rodriguez-Monguio R, Karlsson LH, Hill MOT. Injury During U.S. Army
11 451 Basic Combat Training: A Systematic Review of Risk Factor Studies. *Am J Prev Med.*
12 452 2014;47(6):813-822.
13 453 28. Trank TV, Ryman DH, Minagawa RY, Trone DW, Shaffer RA. Running mileage, movement
14 454 mileage, and fitness in male US Navy recruits. *Med Sci Sports Exerc.* 2001;33(6):1033-1038.
15 455 29. Eckard TG, Padua DA, Hearn DW, Pexa BS, Frank BS. The relationship between training load and
16 456 injury in athletes: a systematic review. *Sports Med.* 2018;48(8):1-33.
17 457 30. Johnston R, Cahalan R, O'keeffe M, O'sullivan K, Comyns T. The associations between training
18 458 load and baseline characteristics on musculoskeletal injury and pain in endurance sport
19 459 populations: A systematic review. *J Sci Med Sport.* 2018(9):910-918.
20 460 31. Soligard T, Schwellnus M, Alonso J-M, et al. How much is too much? (Part 1) International
21 461 Olympic Committee consensus statement on load in sport and risk of injury. *Br J Sports Med.*
22 462 2016;50(17):1030-1041.
23 463 32. Halson S. Monitoring Training Load to Understand Fatigue in Athletes. *Sports Med.*
24 464 2014;44(Supplement 2):139-147.
25 465 33. Piggott B. *The relationship between training load and incidence of injury and illness over a pre-*
26 466 *season at an Australian Football League Club:* Faculty of Computing, Health and Science, Edith
27 467 Cowan University; 2008.
28 468 34. Cummins C, Orr R, O'Connor H, West C. Global positioning systems (GPS) and microtechnology
29 469 sensors in team sports: a systematic review. *Sports Med.* 2013;43(10):1025-1042.
30 470 35. Ehrmann FE, Duncan CS, Sindhusake D, Franzsen WN, Greene DA. GPS and Injury Prevention in
31 471 Professional Soccer. *J Strength Cond Res.* 2016;30(2):360-367.
32 472 36. Colby MJ, Dawson B, Heasman J, Rogalski B, Gabbett TJ. Accelerometer and GPS-derived running
33 473 loads and injury risk in elite Australian footballers. *J Strength Cond Res.* 2014;28(8):2244-2252.
34 474 37. Johnston RJ, Watsford ML, Kelly SJ, Pine MJ, Spurrs RW. Validity and interunit reliability of 10 Hz
35 475 and 15 Hz GPS units for assessing athlete movement demands. *J Strength Cond Res.*
36 476 2014;28(6):1649-1655.
37 477 38. Veenstra BJ, Wyss T, Roos L, Delves SK, Buller M, Beeler N. An evaluation of measurement
38 478 systems estimating gait speed during a loaded military march over graded terrain. *Gait Posture.*
39 479 2018;61:204-209.
40 480 39. Bourdon PC, Cardinale M, Murray A, et al. Monitoring athlete training loads: consensus
41 481 statement. *Int J Sports Physiol Perform.* 2017;12(Suppl 2):S2-161-S162-170.
42 482 40. Benson AC, Bruce L, Gordon BA. Reliability and validity of a GPS-enabled iPhone TM "app" to
43 483 measure physical activity. *J Sports Sci.* 2014;33(14):1-8.
44 484 41. Haff GG, Triplett NT. *Essentials of Strength Training and Conditioning.* 4th ed: Human Kinetics;
45 485 2015.
46 486 42. Lysholm J, Wiklander J. Injuries in runners. *Am J Sports Med.* 1987;15(2):168-171.
47 487 43. Bovens A, Janssen G, Vermeer H, Hoeberigs J, Janssen M, Verstappen F. Occurrence of Running
48 488 Injuries in Adults Following a Supervised-Training Program. *Int J Sports Med.* 1989;10:S186-S190.
49 489 44. Mohr M, Krstrup P, Bangsbo J. Match performance of high-standard soccer players with special
50 490 reference to development of fatigue. *J Sports Sci.* 2003;21(7):519-528.
51 491 45. Sallis JF, Saelens BE. Assessment of Physical Activity by Self-Report: Status, Limitations, and
52 492 Future Directions. *Res Q Exerc Sport.* 2000;71(2):1-14.
53
54
55
56
57
58
59
60
61
62
63
64
65

46. Hoyt RW, Opstad PK, Haugen A-H, DeLany JP, Cymerman A, Friedl KE. Negative energy balance in male and female rangers: effects of 7 d of sustained exercise and food deprivation—. *Am J Clin Nutr*. 2006;83(5):1068-1075.
47. Strath SJ, Swartz AM, Bassett DR, O'Brien WL, King GA, Ainsworth BE. Evaluation of heart rate as a method for assessing moderate intensity physical activity. *Med Sci Sports Exerc*. 2000;32(9 Suppl):S465.
48. Impellizzeri FM, Rampinini E, Coutts AJ, Sassi A, Marcora SM. Use of RPE-based training load in soccer. *Med Sci Sports Exerc*. 2004;36(6):1042-1047.
49. Farrell PA, Wilmore J, Coyle E, Billing J, Costill DL. Plasma Lactate Accumulation and Distance Running Performance. *Med Sci Sports Exerc*. 1993;25(10):1091-1097.
50. Simpson RJ, Graham SM, Connaboy C, Clement R, Pollonini L, Florida-James GD. Blood lactate thresholds and walking/running economy are determinants of backpack-running performance in trained soldiers. *Appl Ergon*. 2017;58:566-572.
51. Borg G. Perceived exertion as an indicator of somatic stress. *Scand J Rehabil Med*. 1970;2(2):92.
52. Foster C, Florhaug JA, Franklin J, et al. A new approach to monitoring exercise training. *J Strength Cond Res*. 2001;15(1):109-115.
53. Gabbett TJ, Domrow N. Relationships between training load, injury, and fitness in sub-elite collision sport athletes. *J Sports Sci*. 2007;25(13):1507-1519.
54. Scott TJ, Black CR, Quinn J, Coutts AJ. Validity and reliability of the session-RPE method for quantifying training in Australian football: a comparison of the CR10 and CR100 scales. *J Strength Cond Res*. 2013;27(1):270-276.
55. O'Leary TJ, Saunders SC, McGuire SJ, Venables MC, Izard RM. Sex Differences in Training Loads during British Army Basic Training. *Med Sci Sports Exerc*. 2018.
56. Bannister E. *Modeling elite athletic performance*. Champaign, IL: Human Kinetics.
57. Manzi V, Iellamo F, Impellizzeri F, D'Ottavio S, Castagna C. Relation between individualized training impulses and performance in distance runners. *Med Sci Sports Exerc*. 2009;41(11):2090.
58. Malone S, Collins K. Relationship Between Individualized Training Impulse and Aerobic Fitness Measures in Hurling Players Across a Training Period. *J Strength Cond Res*. 2016;30(11):3140.
59. Edwards S. High performance training and racing. In: *The Heart Rate Monitor Book*. Sacramento: Fleet Feet Press; 1994:113-123.
60. Gisselman AS, Baxter GD, Wright A, Hegedus E, Tumilty S. Musculoskeletal overuse injuries and heart rate variability: Is there a link? *Med Hypotheses*. 2016;87:1-7.
61. Hamill RW, Shapiro RE, Vizzard MA. Chapter 4 - Peripheral Autonomic Nervous System. In: Robertson D, Biaggioni I, Burnstock G, Low PA, Paton JFR, eds. *Primer on the Autonomic Nervous System (Third Edition)*. San Diego: Academic Press; 2012:17-26.
62. Ogle AD, Young JA. USAF Special Tactics Operator Combat Exposure and Psychological Stress. *Mil Psychol*. 2016;28(3):123-133.
63. Mourot L, Bouhaddi M, Perrey S, et al. Decrease in heart rate variability with overtraining: assessment by the Poincaré plot analysis. *Clin Physiol Funct Imaging*. 2004;24(1):10-18.
64. Le Meur Y, Pichon A, Schaal K, et al. Evidence of parasympathetic hyperactivity in functionally overreached athletes. *Med Sci Sports Exerc*. 2013;45(11):2061.
65. Hedelin R, Wiklund U, Bjerle P, Henriksson-Larsén K. Cardiac autonomic imbalance in an overtrained athlete. *Med Sci Sports Exerc*. 2000;32(9):1531-1533.
66. Davis H, Orzeck T, Keelan P. Psychometric item evaluations of the Recovery-Stress Questionnaire for athletes. *Psychol Sport Exerc*. 2007;8(6):917-938.
67. Kellmann M, Günther K-d. Changes in stress and recovery in elite rowers during preparation for the Olympic Games. *Med Sci Sports Exerc*. 2000;32(3):676-683.

- 1
2
3
4 540 68. Kenttä G, Hassmén P. Overtraining and recovery. A conceptual model. *Sports Med.* 1998;26(1):1-
5 541 16.
6 542 69. Kenttä G, Hassmén P. Train smart: Avoid Overtraining Syndrome. SISU Idrottsbocker; 1999;
7 543 Stockholm.
8 544 70. Sheppard MJ, Triplett NT. *Essentials of Strength Training and Conditioning*. Fourth edition ed.
9 545 Champaign, IL: Human Kinetics; 2016.
10 546 71. Malone S, Roe M, Doran DA, Gabbett TJ, Collins K. High chronic training loads and exposure to
11 547 bouts of maximal velocity running reduce injury risk in elite Gaelic football. *J Sci Med Sport.*
12 548 2017;20(3):250-254.
13 549 72. Hulin BT, Gabbett TJ, Lawson DW, Caputi P, Sampson JA. The acute: chronic workload ratio
14 550 predicts injury: high chronic workload may decrease injury risk in elite rugby league players. *Br J*
15 551 *Sports Med.* 2015(50):231-236.
16 552 73. Viera AJ. Odds ratios and risk ratios: what's the difference and why does it matter? *South Med J.*
17 553 2008;101(7):730.
18 554 74. Murray NB, Gabbett TJ, Townshend AD, Blanch P. Calculating acute: chronic workload ratios
19 555 using exponentially weighted moving averages provides a more sensitive indicator of injury
20 556 likelihood than rolling averages. *Br J Sports Med.* 2016.
21 557 75. Williams S, West S, Cross MJ, Stokes KA. Better way to determine the acute:chronic workload
22 558 ratio? *Br J Sports Med.* 2017;51(3):209-210.
23 559 76. Lolli L, Batterham AM, Hawkins R, et al. Mathematical coupling causes spurious correlation
24 560 within the conventional acute-to-chronic workload ratio calculations. *Br J Sports Med.* 2017.
25 561 77. Bornn L, Ward P, Norman D. Training Schedule Confounds the Relationship between
26 562 Acute:Chronic Workload Ratio and Injury. MIT Sloan Sports Analytics Conference; 2019; Boston,
27 563 MA.
28 564 78. Harrison PW, Johnston RD. Relationship Between Training Load, Fitness, and Injury Over an
29 565 Australian Rules Football Preseason. *J Strength Cond Res.* 2017;31(10):2686-2693.
30 566 79. Strowbridge N, Burgess K. Sports And Training Injuries In British Soldiers: The Colchester
31 567 Garrison Sports Injury And Rehabilitation Centre. *J R Army Med Corps.* 2002;148(3):236.
32 568 80. Adams MS, Croft A. Sports Injuries in British Troops deployed on Operation Resolute (Bosnia). *J*
33 569 *R Army Med Corps.* 1997;143(1):35.
34 570 81. Knapik JJ, Sharp MA, Canham-Chervak M, Hauret K, Patton JF, Jones BH. Risk factors for training-
35 571 related injuries among men and women in basic combat training. *Med Sci Sports Exerc.*
36 572 2001;33(6):946.
37 573 82. Galdorisi G. The 2016 Australian Defence white paper: A view from across the pacific. *Asia-*
38 574 *Pacific Defence Reporter (2002).* 2016;42(5):34-38.
39 575 83. Morton RH. Modeling training and overtraining. *J Sports Sci.* 1997;15(3):335.
40 576 84. Hauschild V, Hauret K, Richardson M, Jones BH, Lee T. *A Taxonomy of Injuries for Public Health*
41 577 *Monitoring and Reporting.* Army Public Health Center Aberdeen Proving Ground United
42 578 States;2017.
43 579 85. Orr R, Ford K, Stierli M. Implementaion of an Ability-Based Training Program in Police Force
44 580 Recruits. *J Strength Cond Res.* 2016;30(10):2781-2787.
45 581 86. Sherrard J, Lenné M, Cassell E, Stokes M, Ozanne-Smith J. Injury prevention during physical
46 582 activity in the Australian Defence Force. *J Sci Med Sport.* 2004;7(1):106-117.
47 583
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Ratings of perceived exertion (RPE)	Total quality recovery (TQR)
6	6
7 Very, very light	7 Very, very poor recovery
8	8
9 Very light	9 Very poor recovery
10	10
11 Fairly light	11 Poor recovery
12	12
13 Somewhat hard	13 Reasonable recovery
14	14
15 Hard	15 Good recovery
16	16
17 Very hard	17 Very good recovery
18	18
19 Very, very hard	19 Very, very good recovery
20	20

Rating	Descriptor
0	Rest
1	Very, Very Easy
2	Easy
3	Moderate
4	Somewhat Hard
5	Hard
6	.
7	Very Hard
8	.
9	.
10	Maximal

Athletes can fill out this guide over the course of a week to assess their own recovery behaviors. Once they have scored a full day, the total reveals whether they are paying adequate attention to their physical and mental recovery needs. **17-20** daily points is optimal; **15-16** points is good but shows room for improvement; **14 or fewer** points means the athlete needs a serious individual evaluation of recovery behaviors.

[illegible]