Gait speed characteristics and Its spatiotemporal determinants in nursing home residents: A cross-sectional study

Fien, Samantha; Henwood, Timothy; Climstein, Mike; Rathbone, Evelyne; Keogh, Justin William Leslie

Published in:
Journal of Geriatric Physical Therapy

DOI:
[10.1519/JPT.0000000000000160](https://doi.org/10.1519/JPT.0000000000000160)

Licence:
Other

[Link to output in Bond University research repository.](#)

Recommended citation(APA):
Fien, S., Henwood, T., Climstein, M., Rathbone, E., & Keogh, J. W. L. (2019). Gait speed characteristics and Its spatiotemporal determinants in nursing home residents: A cross-sectional study. *Journal of Geriatric Physical Therapy*, 42(3), E148-E154. <https://doi.org/10.1519/JPT.0000000000000160>

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

For more information, or if you believe that this document breaches copyright, please contact the Bond University research repository coordinator.

Journal of Geriatric Physical Therapy

Gait Speed Characteristics and Its Spatio-Temporal Determinants in Nursing Home Residents: A Cross-Sectional Study --Manuscript Draft--

Manuscript Number:	JGPT-D-17-00054R1
Full Title:	Gait Speed Characteristics and Its Spatio-Temporal Determinants in Nursing Home Residents: A Cross-Sectional Study
Article Type:	Research Reports/Quantitative
Keywords:	allied health professionals; gait speed; nursing home; spatio-temporal determinants
Corresponding Author:	Samantha Fien, BExSc Bond University Robina, AUSTRALIA
Corresponding Author Secondary Information:	
Corresponding Author's Institution:	Bond University
Corresponding Author's Secondary Institution:	
First Author:	Samantha Fien, BExSc
First Author Secondary Information:	
Order of Authors:	Samantha Fien, BExSc
	Tim Henwood, PhD
	Mike Climstein, PhD FASMF, FACSM, FAAESS, AEP
	Evelyne Rathbone, MSc, GradStat
	Justin William Leslie Keogh, PhD
Order of Authors Secondary Information:	
Manuscript Region of Origin:	AUSTRALIA
Abstract:	<p>Background and purpose: Low and slowing gait speeds among nursing home residents are linked to a higher risk of disability, cognitive impairment, falls, and mortality. A better understanding of the spatio-temporal parameters of gait that influence declining mobility could lead to effective rehabilitation and preventative intervention. This study aims were to objectively quantify the spatio-temporal characteristics of gait in the nursing home setting and define the relationship between these parameters and gait speed.</p> <p>Methods: One hundred nursing home residents were enrolled into the study and completed three habitual gait speed trials over a distance of 3.66 m. Trials were performed using an instrumented gait analysis. The manner in which the spatiotemporal parameters predicted gait speed was examined by univariate and multivariable regression modelling.</p> <p>Results: The nursing home residents had a habitual gait speed of 0.63 ± 0.19 m/s, a stride length of 0.83 ± 0.15 m, a support base of 0.15 ± 0.06 m and step time of 0.66 ± 0.12 s. Multivariable linear regression revealed stride length, support base and step time predicted gait speed ($R^2 = 0.89$, $p < 0.05$). Step time had the greatest influence on gait speed with each 0.1 s decrease in step time resulting in a 0.09 m/s (95% CI 0.08 - 0.10) increase in habitual gait speed.</p> <p>Conclusion: This study revealed step time, stride length and support base are the strongest predictors of gait speed among nursing home residents. Given the impact of low and slowing gait speed in this population, future research should concentrate on developing and evaluating intervention programs that were specifically designed to focus on improving step time, stride length and support base in nursing home residents. As gait speed has been shown to be predictive of many adverse events in older adults, we would also suggest that routine assessments of gait speed, and if</p>

	possible their spatiotemporal characteristics be done on all nursing home residents in an attempt to identify residents with low or slowing gait speed.
Response to Reviewers:	Response to the editor and reviewer's comments can be found in the cover letter in a table that includes point-by-point response to each comment.

Gait Speed Characteristics and Its Spatio-Temporal Determinants in Nursing Home Residents: A Cross-Sectional Study

Samantha Fien (BExSc)¹, Timothy Henwood (PhD)^{2,1}, Mike Climstein (PhD
FASMF, FACSM, FAAESS, AEP)^{3,4}, Evelyn Rathbone (MSc, GradStat)¹,
Justin William Leslie Keogh (PhD)^{1,5,6}

1. Faculty of Health Science and Medicine, Bond University, Robina, QLD,
Australia;
2. Southern Cross Care, North Plympton, SA, Australia;
3. Exercise, Health and Performance Faculty Research Group, The University
of Sydney, Sydney, NSW, Australia;
4. Water Based Research Unit, Faculty of Health Sciences, Bond University,
Gold Coast, QLD, Australia;
5. Human Potential Centre, AUT University, Auckland, New Zealand;
6. Cluster for Health Improvement, Faculty of Science, Health, Education and
Engineering, University of the Sunshine Coast, Sippy Downs, QLD, Australia.

Corresponding Author

Samantha Fien
Faculty of Health Science and Medicine,
Bond University,
Robina, Australia
4229
Ph. +617 5595 4487
Fax: +617 5595 3524

E. samantha.fien@student.bond.edu.au

Acknowledgements

We would like to thank the management, staff and residents of the nursing home facilities for their assistance and participation in this project.

Disclosure statement

The Authors declare that there is no conflict of interest. This research did not receive any specific grant from funding agencies in the public, commercial, or not-for-profit sectors. The authors certify that they comply with the ethical guidelines for authorship and publishing of the Journal of Geriatric Physical Therapy.

Gait Speed Characteristics and Its Spatio-Temporal Determinants in Nursing Home Residents: A Cross-Sectional Study

ABSTRACT

Background and purpose: Low and slowing gait speeds among nursing home residents are linked to a higher risk of disability, cognitive impairment, falls, and mortality. A better understanding of the spatio-temporal parameters of gait that influence declining mobility could lead to effective rehabilitation and preventative intervention. This study aims were to objectively quantify the spatio-temporal characteristics of gait in the nursing home setting and define the relationship between these parameters and gait speed.

Methods: One hundred nursing home residents were enrolled into the study and completed three habitual gait speed trials over a distance of 3.66 m. Trials were performed using an instrumented gait analysis. The manner in which the spatiotemporal parameters predicted gait speed was examined by univariate and multivariable regression modelling.

Results: The nursing home residents had a habitual gait speed of 0.63 ± 0.19 m/s, a stride length of 0.83 ± 0.15 m, a support base of 0.15 ± 0.06 m and step time of 0.66 ± 0.12 s. Multivariable linear regression revealed stride length, support base and step time predicted gait speed ($R^2 = 0.89$, $p < 0.05$). Step time had the greatest influence on gait speed with each 0.1 s decrease in step time

1 22 resulting in a 0.09 m/s (95% CI 0.08 - 0.10) increase in habitual gait speed.
2
3

4 23 **Conclusion:** This study revealed step time, stride length and support base are
5
6 24 the strongest predictors of gait speed among nursing home residents. Given the
7
8 25 impact of low and slowing gait speed in this population, future research should
9
10 26 concentrate on developing and evaluating intervention programs that were
11
12 27 specifically designed to focus on improving step time, stride length and support
13
14 28 base in nursing home residents. As gait speed has been shown to be predictive
15
16 29 of many adverse events in older adults, we would also suggest that routine
17
18 30 assessments of gait speed, and if possible their spatiotemporal characteristics
19
20 31 be done on all nursing home residents in an attempt to identify residents with
21
22 32 low or slowing gait speed.
23
24
25
26
27
28

29 33 Keywords: health professionals; gait speed; nursing home; spatio-temporal
30
31 34 determinants
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

INTRODUCTION

Walking is a key physical performance task for people of all ages, including older adults. The majority of older adults, especially those living in nursing home (residential aged care) settings have decreased physical activity¹ and poor physical function as indicated by their reduced gait speed, muscle strength and balance.² Older adults with slower gait speeds are at higher risk of disability, cognitive impairment, institutionalization, falls, and mortality.³ While a variety of gait speed thresholds exist, healthy community dwelling older adults tend to experience poorer health when their habitual gait speed is < 0.8 m/s,^{3,4} whereas for nursing home residents > 80 years of age, a threshold of < 0.5 m/s has been proposed.^{4,5} A recent systematic review which included 34 studies quantifying the gait speed of residents living in nursing homes reported a mean habitual pace gait speed of 0.48 m/s (95% confidence interval (CI) 0.40-0.55).⁶ Gait speeds this low suggest that most nursing home residents are limited in mobility and independence, have decreased stability and are at increased risk for many other age-related conditions.⁶

Currently, little is known in relation to the physical determinants or risk factors for low gait speed in low-functioning older adults and those living in nursing home facilities. While McGough et al.⁷ and Keogh et al.⁸ have reported that measures of physical function, balance, lifetime physical activity levels and sitting time correlate with gait speed in these less functioning older cohorts, no studies have quantified the spatio-temporal determinants (e.g. step length, step rate) that determine gait speed in a nursing home population.⁹ Figure 1 presents a pictorial representation of the relationship between selected spatio-

temporal parameters and gait speed. When compared to younger adults, older adults walk more slowly; have a shorter step length and a broader support base than their younger counterparts, with these differences more pronounced in older adults living in nursing home and/or with a high risk of falls.¹⁰⁻¹³

Insert Figure 1 here.

A greater understanding of nursing home residents' gait speed and spatio-temporal determinants may assist health professionals to identify nursing home residents at high risk of adverse events and allow for a more specific-tailored physical therapy and rehabilitation program for each individual.¹¹ We would argue this is very important as indicated by a systematic review that found exercise-related improvements in older adults' gait speed are typically smaller in magnitude and more variable than the improvements in muscular strength.¹⁴ Specifically, semi-regular monitoring of gait speed and spatiotemporal parameters could assist appropriate health professionals to prescribe resistance, balance and gait exercises that target their clients' major spatiotemporal limitations. The collection of this gait spatiotemporal data may also allow the exercise therapist to provide their client task-relevant augmented feedback (e.g. visual cues, instant or delayed feedback) during these exercises to improve the transfer of training to their habitual walking performance^{15,16}.

This study aims were to objectively quantify the spatio-temporal determinants and gait speed of nursing home residents and to gain some insight into whether these spatiotemporal parameters may predict their gait speed.

METHODS

Recruitment and Study Design

The study employed a cross-sectional design, with data collected over an eight-month period across three nursing home facilities in South East Queensland, Australia. The facilities were from different providers that were either part of a small chain of nursing homes or a not for profit organisation. The flow of recruitment to assessment is represented in Figure 2. Facilities were approached via email and telephone follow-up seeking an expression of interest for participation. Following an expression of interest, nursing homes were visited and the study explained to the Service Manager. Once the service manager approved the participation of their nursing home in the project, eligible participants were identified at a meeting between the project lead researcher and the Service Manager, head Registered Nurse and head Diversional Therapist. Ethical approval for this study was attained from the University Human Ethics Research Committee (RO 1823) and gatekeeper's approval obtained through the nursing homes.

Insert Figure 2 here.

Based on study's eligibility, participants were eligible for inclusion if they were:

a) aged 65 years and over, b) residing in a nursing home facility, c) ambulate independently or without a walking aid and d) could provide informed consent.

The exclusion criteria included: a) end-stage terminal and/or life expectancy <6-months (ethical reasons), b) two person transfer or increased falls risk during ambulation (as assessed by the nursing home staff), c) unable to communicate or follow instructions (personal needs beyond the scope of this project) and d) behaviors that would endanger the participant or research staff.

All participants were approached personally about participation and given the opportunity to ask questions or raise concerns about the study. Following this discussion and reading of the participant information sheet, participants provided their informed consent if they wished to participate. A total of 100 participants took part in the study, with the primary investigator responsible for observing and administering all of the assessments.

Primary Outcome Measures: Gait Speed

Gait speed was assessed using a computer interfaced electronic system (model GaitMat II, EQInc, USA) which required participants to walk across a level pressure mat system that was 3.66 m (11.91 ft.) long.¹⁷ The concurrent validity of the spatio-temporal determinants of gait recorded with the GaitMat II is extremely high when compared to the criterion method of 3-D motion capture ($R = .99$).^{18,19} The Gait Mat II was chosen due to it being much more feasible to use in nursing homes and 3-D motion capture. The GaitMat II system automatically measured gait speed and spatio-temporal determinants, with this data automatically stored in a Microsoft Excel spreadsheet.

Participants completed the trials at their habitual gait speed in their regular footwear. The following instructions were provided, "Walk towards the end of the room in the centre of the mat at a pace that is comfortable for you". All measures were initiated from a standing start 2 m (6.56 ft.) from the GaitMat II platform in order to reduce the effect that acceleration or deceleration may have on the outcomes.^{20,21} The average gait speed (m/s) from three attempts was used for data analysis. Participants were allowed as much rest as required between attempts, with rest periods typically being up to one minute.

Secondary Outcome Measures

A full spectrum of spatio-temporal gait determinants outputted was recorded.

These spatio-temporal gait determinants included step length, stride length, support base, step time, swing time, stance time, single support time and double support time and are defined by the GaitMat II manual found in Table 1. Insert Table 1 here.

In addition handgrip strength, the Mini-Cog test²² and a simple five-item questionnaire (SARC-F) ^{23,24} were collected for the purpose of cohort characteristics description. Nursing home facility records provided other relevant descriptors including the number of medical conditions and medications.

Data Management and Statistical Analysis

All data were initially checked for normality prior to analysis. As data were normally distributed, descriptive statistics are presented as mean and standard deviations for continuous variables. A one-way ANOVA and post-hoc Tukey and Scheffe tests were performed to investigate between nursing home differences. Linear regression analyses were performed to gain insight into the potential determinants of gait speed (i.e. gait spatio-temporal determinants) in residents. Univariate analyses of all gait spatio-temporal determinants were employed to identify possible determinants of gait speed (two-tailed). Factors with a significance $p \leq 0.10$ determined from simple linear regression analyses were included in the multiple linear regression model. This multivariable model determined which combination of variables best-predicted gait speed in

residents. The 95% confidence interval (95% CI) was included for the coefficients in the multivariable model. All data were analysed using SPSS statistic software (version 22) with statistical significance set at $p < 0.05$ a priori.

RESULTS

Participants

One hundred of 166 (60.24%) invited, eligible residents were recruited to the study. There were no significant differences between all variables for nursing home cohorts in this study ($p > 0.05$), thus data combined into one group for analysis. Cohort data are present in Table 2. The average age of the 100 residents was 85.7 (7.1) years with a mean gait speed of 0.63 (0.19) m/s, an average of 11.0 (4.9) medical conditions and 14.0 (5.8) prescribed medications. There were no significant differences with gait speed and spatio-temporal parameters, handgrips strength, sarcopenia status and medications and chronic diseases across males and females. However, males were significantly younger ($p = 0.038$) and had a lower Mini-Cog assessment ($p = 0.002$) in comparison to females.

Insert Table 2 here.

The majority of participants (79%, $n = 79/100$) presented with below normal habitual gait speeds (< 0.80 m/s), whilst 26% ($n = 26/100$) ambulated at below the mean reported for nursing facilities residents (< 0.48 m/s, 95% CI 0.396-0.554).⁶

Results of the univariable linear regression analyses identified three spatio-temporal factors as being predictive of gait speed: stride length ($p < 0.001$), support base ($p < 0.001$) and step time ($p = 0.002$) (see Table 3). Of

these factors, stride length contributed to the largest change in gait speed, with each 0.1 m increase in stride length resulting in an average 0.09 m/s (95% CI 0.06 – 0.13) faster habitual gait speed.

Insert Table 3 here.

The multivariable linear regression model that included stride length, support base and step time predicted 89% ($R^2 = 0.89$) of the variation in gait speed (see Table 3). Specifically, step time contributed to the largest change in gait speed with every 0.1 s decrease in step time resulting in a mean increase in gait speed of 0.09 m/s (95% CI 0.08 – 0.10). A 0.1 m increase in stride length was also associated with a mean increase of 0.08 m/s (95% CI 0.07 – 0.09) in gait speed. The third determinant identified in the multivariable regression, support base appeared to have a smaller effect on gait speed, with a 0.1 m decrease in support base resulting in a mean gait speed increase of 0.04 m/s (95% CI 0.02 – 0.07).

DISCUSSION

This study demonstrated that nursing home residents who can self-ambulate with or without a walking aid still walk at a gait speed (0.63 ± 0.19 m/s) and possess spatiotemporal parameters that place them at high risk of falls and other adverse age-related events.^{3,11,12} A total of 79 participants presented with below normal habitual gait speeds (< 0.80 m/s), which is a threshold defined to screen for sarcopenia in older adults aged 80 years and older.²⁵ A total of 27 participants also walked at a threshold below 0.48 m/s which a meta-analysis of 48 studies found to be the mean gait speed for older adults in nursing homes.⁶

Results of the regression analyses also indicated that nursing home residents who ambulated at a slower habitual gait speed were more likely to have an increased step time, shorter stride length and a wider support base than their more ambulatory counterparts. While the finding that some spatio-temporal parameters do predict gait speed is not overly surprising, to our knowledge this is the first study to investigate the potential for spatio-temporal determinants to determine gait speed in the nursing home setting. The importance of the spatio-temporal parameters in determining gait speed also appear consistent with Sterke et al.¹¹ and Taylor et al.¹² who demonstrated that slower older walkers with increased falls risk had shorter stride lengths, longer double support times and a wider support base when compared to aged matched individuals with no falls history.

The significant ability of spatio-temporal determinants such as step time, stride length and support base to determine gait speed in the current study ($R^2 = 0.89$) and falls in previous studies^{11,12} would appear to reflect a variety of biomechanical concepts. For example, the ability of step time and stride length to determine gait speed may be explained by the impulse-momentum relationship and/or the nursing home residents' reduced ability to maintain balance during the gait cycle. It is fair to conclude that reduced lower-body muscle strength and power may mean that the nursing home residents require greater single/double support times to produce the necessary impulse (force multiplied by time) to propel their body forward during the gait cycle.²⁶ Their reduced force production ability and greater stance time would then contribute to a reduced stride length, increased step time and ultimately a reduced

habitual gait speed. Poor stride length may also be suggestive of shuffling gait and low plantar flexor and hip flexor strength.²⁶ Therefore, certain exercises such as calf raises and leg raises coupled with gait training may need to be incorporated in resistance training programs to improve gait speed in nursing home residents.²¹

The clinical significance of this study is that gait speed characteristics and spatio-temporal determinants are becoming more easily measured and analysed in nursing home settings. Health professionals can then use this individualized gait data to identify residents at risk of adverse events and intervene where appropriate by providing an individualized exercise intervention for each resident. In doing so, residents are likely to benefit more from these exercise programs as they are better tailored to the specific spatio-temporal parameters underlying the participants' poor gait speed and/or falls risk. Such an approach may also improve exercise adherence as these programs can better concentrate on improving gait performance in activities of daily living (ADL).²⁷

Collectively, the manner in which the nursing home residents walk (as described by their spatio-temporal determinants) have major implications to exercise therapy and rehabilitative approaches to improving gait speed in this cohort. With the clinical implications being that if health professionals can continually monitor and assess gait speed and spatio-temporal determinants, we may be able to decrease or prolong the amount of residents who are induced into the vicious cycle of reduced physical activity and decreased

mobility and physical performance that have a direct effect on their health and survival.

However, it is also possible that the tendency for the slower nursing home residents to have shorter stride lengths, increased step times and wider support bases may be indicative of a compensation for their reduced strength and dynamic stability. Longer strides and a narrower support base increase the distance that the centre of mass travels outside the anterior-posterior and medial-lateral bases of support, respectively.²⁶ As Sherrington et al.²⁸ has reported that exercise programs that do not sufficiently challenge balance may actually increase rather than decrease the risk of falling of nursing home residents, we would recommend that nursing home residents with short strides and wide support bases focus initially on improving their static and dynamic balance. Once balance has been improved in the anterior-posterior and medial lateral directions, these residents may further prioritize resistance and gait retraining to safely improve their gait speed and overall mobility.

Given the interplay between decreasing mobility and increasing disability, the monitoring of gait speed by health professionals (and if possible the primary spatio-temporal determinants) on at least an annual basis in the nursing home setting has been recommended.^{8,29} For those nursing home residents identified with poor and slowing mobility, systematic review evidence suggests that regular progressive resistance and balance training can improve their habitual gait speed by 0.07 m/s (95% CI 0.02-0.11) when compared to non-exercising controls.³⁰ While these reported improvements in gait speed are positive, there may be two potential criticisms of the studies reviewed in this meta-analysis

(and the wider literature). The first is that the studies have typically used quite generic exercise prescriptions that focus on improving muscular hypertrophy and strength in a variety of muscle groups. Based on emerging evidence that reduced gait speed in older adults is primarily a result of reduced ankle plantarflexor rather than hip or knee extensor moment and muscle power,²⁶ a greater focus on increasing the muscle strength and power of the plantarflexors compared to the traditional focus on the knee and hip extensors may be warranted. In addition, the majority of studies in this area that have included balance training primarily used static balance tasks that require the older adults to hold a position for a period of 10-20 s e.g. two feet stands on unstable surface or with eyes closed or semi-tandem/tandem stance. Based on our results, we suggest that dynamic balance ability, which would appear more closely related to the balance requirements of human gait, be taught by health professionals on a weekly basis. Therefore, nursing home residents may obtain greater gait speed benefit from performing dynamic balance tasks (e.g. stepping and perturbation response) than static balance tasks.³¹⁻³³

The Gait Mat II provided a feasible, reliable and valid tool to measure gait speed and spatio-temporal determinants in nursing home adults.^{2,18,19} New equipment has been developed since the data collection of this study. One suggestion for future studies would be to use inertial sensors³⁴ which may be a more portable and affordable gait assessment. Such advancements in inertial sensor technology would more easily allow health professionals to routinely monitor gait speed and spatiotemporal parameters in the nursing home setting,

which may further increase allow the development of targeted exercise programs for each nursing home resident.

Participation selection bias is a limitation that may have influenced our findings as the inclusion criteria deemed that a participant should have the ability to walk with or without an aid. A total of 55% of residents in the nursing home facilities were ineligible because of the inability to mobilize or because they were deemed too high a risk to participate. Because of this bias in selecting individuals who were ambulant (with or without assistive devices), the gait speed and spatiotemporal parameters obtained in this study may not be generalized to all nursing home residents. Nevertheless, the participants in the current study were still below the cut off for physical performance and at risk of further decreased disability, cognitive decline and mortality,³⁵ with every 0.1 m/s reduction in gait speed equating to a 10% decrease in older adult's ability to perform ADLs.³⁶ It must also be acknowledged that static or dynamic balance ability were not directly assessed in this study. Therefore, while our proposition of poor dynamic balance contributing to the reduced gait speed of nursing home participants has some experimental support,^{37,38} we cannot explicitly state that is the case with our participants.

CONCLUSIONS

This is the first study to investigate gait speed characteristics and spatiotemporal determinants in the nursing home setting. While our cross-sectional study suggests that step time, stride length and support base are highly predictive of gait speed in nursing home residents, longitudinal research is

1 322 required to determine if changes in these three spatio-temporal determinants
2
3 323 may be predictive of changes in gait speed. If these longitudinal relationships
4
5 324 between gait speed and spatio-temporal determinants can be found, health
6
7 325 professionals may be better able to alter aspects of their exercise prescription
8
9 326 and augmented feedback approach to improve outcomes for nursing home
10
11 327 residents.
12
13 328
14
15 329
16
17 330
18
19 331
20
21 332
22
23 333
24
25 334
26
27 335
28
29 336
30
31 337
32
33 338
34
35 339
36
37 340
38
39 341
40
41 342
42
43 343
44
45 344
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Acknowledgements

We would like to thank the management, staff and residents of the nursing home settings for their assistance and participation in this project.

Disclosure statement

The Authors declare that there is no conflict of interest. This research did not receive any specific grant from funding agencies in the public, commercial, or not-for-profit sectors. The authors certify that they comply with the ethical guidelines for authorship and publishing of The Journal of Geriatric Physical Therapy and that the rights of human subjects were protected.

References

1. Reid N, Eakin E, Henwood T, et al. Objectively measured activity patterns among adults in residential aged care. *Int J Environ Res Public Health*. 2013;10(12):6783-6798.
2. Kuys SS, Peel NM, Klein K, Slater A, Hubbard RE. Gait speed in ambulant older people in long term care: a systematic review and meta-analysis. *JAMDA*. 2014;15(3):194-200.
3. Abellan van Kan G, Rolland Y, Andrieu S, et al. Gait speed at usual pace as a predictor of adverse outcomes in community-dwelling older people an International Academy on Nutrition and Aging (IANA) Task Force. *J Nutr Health Aging*. 2009;13(10):881-889.
4. Cruz-Jentoft AJ, Baeyens JP, Bauer JM, et al. Sarcopenia: European consensus on definition and diagnosis: report of the European Working Group on Sarcopenia in Older People. *Age Ageing*. 2010;39(4):412-423.
5. Weidung B, Bostrom G, Toots A, et al. Blood pressure, gait speed, and mortality in very old individuals: a population-based cohort study. *JAMDA*. 2015;16(3):208-214.
6. Peel NM, Kuys SS, Klein K. Gait speed as a measure in geriatric assessment in clinical settings: a systematic review. *J Gerontol A Biol Sci Med Sci*. 2013;68(1):39-46.
7. McGough EL, Logsdon RG, Kelly VE, Teri L. Functional mobility limitations and falls in assisted living residents with dementia. *J Geriatr Phys Ther*. 2012;195(1):78-86.

- 1 404 8. Keogh JW, Senior H, Beller EM, Henwood T. Prevalence and risk factors
2
3 405 for low habitual walking speed in nursing home residents: an
4
5 406 observational study. *Arch Phys Med Rehabil.* 2015;96(11):1993-1999.
6
7
8 407 9. Espy DD, Yang F, Bhatt T, Pai YC. Independent influence of gait speed
9
10 408 and step length on stability and fall risk *Gait Posture.* 2010;32(3):378-
11
12 409 382.
13
14
15 410 10. Hausdorff JM, Edelberg HK, Mitchell SL, Goldberger AL, Wei JY.
16
17 Increased gait unsteadiness in community-dwelling elderly fallers. *Arch*
18 411
19 *Phys Med Rehabil.* 1997;78(3):278-283.
20 412
21
22 413 11. Sterke CS, van Beeck EF, Loonman CWN, Kressig RW, van der
23
24 414 Cammen TJM. An electronic walkway can predict short-term fall risk in
25
26 415 nursing home residents with dementia. *Gait Posture.* 2012;36(1):95-101.
27
28
29 416 12. Taylor ME, Delbaere K, Mikolaizak AS, Lord SR, Close JCT. Gait
30
31 417 parameter risk factors for falls under simple and dual task conditions in
32
33 418 cognitively impaired older people. *Gait Posture.* 2013;37(1):126-130.
34
35
36 419 13. Agner S, Bernet J, Brulhart Y, Radlinger L, Rogan S. Spatiotemporal gait
37
38 420 parameters during dual task walking in need of care elderly and young
39
40 421 adults. a cross-sectional study. *Z Gerontol Geriatric.* 2015;48(8):740-746.
41
42
43 422 14. Valenzuela T. Efficacy of progressive resistance training interventions in
44
45 423 older adults in nursing homes: a systematic review. *JAMDA.*
46
47 424 2012;13(5):418-428.
48
49
50 425 15. Begg RK, Tirosh O, Said CM, et al. Gait training with real-time
51
52 426 augmented toe-ground clearance information decreases tripping risk in
53
54
55
56
57
58
59
60
61
62
63
64
65

1 427 older adults and a person with chronic stroke. *Front Hum Neurosci.*
2
3 428 2014;8:243.
4
5
6 429 16. Shen X, Mak MKY. Balance and gait training with augmented feedback
7
8 430 improves balance confidence in people with parkinson's disease.
9
10 431 *Neurorehabilitation and Neural Repair.* 2014;28(6).
11
12
13 432 17. McDonough AL, Batavia M, Chen FC, Kwon S, Ziai J. The validity and
14
15 433 reliability of the GAITRite system's measurements: a preliminary
16
17 434 evaluation. *Arch Phys Med Rehabil.* 2001;82(3):419-425.
18
19
20 435 18. Huang W-N, W., , VanSwearingen JM, Brach JS. Gait variability in older
21
22 436 adults: observational rating validated by comparison with a computerized
23
24 437 walkway gold standard. *Phys Ther.* 2008;88(10):1146-1153.
25
26
27 438 19. Barker S, Craik R, Freedman W, et al. Accuracy, reliability, and validity of
28
29 439 a spatiotemporal gait analysis system. *Med Eng Phys.* 2006;28(5):460-
30
31 440 467.
32
33
34
35 441 20. Kressig RW, Gregor RJ, Oliver A, et al. Temporal and spatial features of
36
37 442 gait in older adults transitioning to frailty. *Gait Posture.* 2004;20(1):30-35.
38
39
40 443 21. Fien S, Henwood T, Climstein, M., & Keogh, J. W. L. Feasibility and
41
42 444 benefits of group based exercise in residential aged care adults: a pilot
43
44 445 study for the GrACE programme. *PeerJ.* 2016;4:e2018(eCollection
45
46 446 2016).
47
48
49 447 22. Borson S, Scanlan JM, Watanabe J, Tu SP, Lessig M. Improving
50
51 448 identification of cognitive impairment in primary care. *Int J Geriatr*
52
53 449 *Psychiatry.* 2006;21(4):349-355.
54
55
56
57
58
59
60
61
62
63
64
65

- 1 450 23. Malmstrom TK, & Morley JE. SARC-F: A simple questionnaire to rapidly
2
3 451 diagnose sarcopenia. *JAMDA*. 2013;14(8):1-2.
4
5 452 24. Cao L, Chen S, Zou C, et al. A pilot study of the SARC-F scale on
6
7 453 screening sarcopenia and physical disability in the Chinese older people.
8
9 454 *J Nutr Health Aging*. 2014;18(3):277-283.
10
11 455 25. Lourenco RA, Perez-Zepeda M, Gutierrez-Robeldo L, Garcia-Garcia FJ,
12
13 456 Manas LR. Performance of the european working group on sarcopenia in
14
15 457 older people algorithm in screening older adults for muscle mass
16
17 458 assessment. *Age and Ageing*. 2015;44(334-338).
18
19 459 26. Kulmala JP, Korhonen MT, Kuitunen S, et al. Which muscles
20
21 460 compromise human locomotor performance with age? *J R Soc Interface*.
22
23 461 2014;11(100):20140858.
24
25 462 27. Henwood T, Neville C, Baguley C, Clifton K, Beattie E. Physical and
26
27 463 functional implications of aquatic exercise for nursing home residents
28
29 464 with dementia *Geriatr Nurs*. 2015;36(1):35-39.
30
31 465 28. Sherrington C, Tiedemann A, Fairhall N, Close JCT, Lord SR. Exercise
32
33 466 to prevent falls in older adults: an updated meta-analysis and best
34
35 467 practice recommendations. *NSW Public Health Bull*. 2011;22(4):78-83.
36
37 468 29. Fien S, Henwood T, Climstein M, Keogh JWL. Clinical importance of
38
39 469 assessing walking speed in older adults in general practice. *Aust Fam*
40
41 470 *Physician*. 2016;45(4):250-251.
42
43 471 30. Chou CH, Hwang CL, Wu YT. Effect of exercise on physical function,
44
45 472 daily activities, and quality of life in the frail older adults: a meta analysis.
46
47 473 *Arch Phys Med Rehabil*. 2012;93(2):237-244.
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

- 1 474 31. Nnodim JO, Strasburg D, Nabozny M, et al. Dynamic balance and
2
3 475 stepping versus Tai Chi training to improve balance and stepping in at-
4
5 476 risk older adults. *J Am Geriatr Soc.* 2006;54(12):1825-1831.
6
7
8 477 32. Granacher U, Zahner L, Gollhofer A. Strength, power, and postural
9
10 478 control in seniors: Considerations for functional adaptations and for fall
11
12 479 prevention. *Eur J Sport Sci.* 2008;8(6):325-340.
13
14
15 480 33. Carty CP, Cronin NJ, Lichtwark GA, Mills PM, Barrett RS. Mechanisms of
16
17 481 adaptation from a multiple to a single step recovery strategy following
18
19 482 repeated exposure to forward loss of balance in older adults. *PLoS One.*
20
21 483 2012;7(3):e33591.
22
23
24
25 484 34. Greene BR, Redmond SJ, Caulfield B. Fall risk assessment through
26
27 485 automatic combination of clinical fall risk factors and body-worn sensor
28
29 486 data. *IEEE J Biomed Health Inform.* 2016;[Epub ahead of print].
30
31
32 487 35. Houles M, Abellan Van Kan G, Rolland Y, Andrieu S, Bauer J, et al. Gait
33
34 488 speed at usual pace as a predictor of adverse outcomes in community-
35
36 489 dwelling older people [French] La vitesse de marche comme critere de
37
38 490 fragilite chez la personne agee vivant au domicile. *Cah l'Annee Gerontol*
39
40 491 2010;2(1):13-23.
41
42
43
44 492 36. Hausdorff J. Gait dynamics, fractals and falls: finding meaning in the
45
46 493 stride-to-stride fluctuations of human walking. *Hum Mov Sci.*
47
48 494 2007;26(4):557-589.
49
50
51
52 495 37. Gusi N, Carmelo Adsuar J, Corzo H, del Pozo-Cruz B, Olivares PR,
53
54 496 Parraca JA. Balance training reduces fear of falling and improves
55
56
57
58
59
60
61
62
63
64
65

1	497	dynamic balance and isometric strength in institutionalised older people:
2		
3	498	a randomised trial <i>Journal of Physiotherapy</i> . 2012;58(2):97-104.
4		
5	499	38. Mohammadi M, Kaldirimci M, Ebrahim Kazemi S, Mizrak. O.,
6		
7	500	Tugrulhansam C. The effect of pilates exercise on gait speed and
8		
9	501	strength of lower limb in elderly male. <i>Advances in Applied Science</i>
10		
11	502	<i>Research</i> . 2015;6(7):1-6.
12		
13	503	39. Walsh J. GaitMat II User's Manual. <i>EQ, Inc Chalfont, PA</i> .Revision 3.2.1.
14		
15	504	
16		
17	505	
18		
19	506	
20		
21	507	
22		
23	508	
24		
25	509	
26		
27	510	
28		
29	511	
30		
31	512	
32		
33	513	
34		
35	514	
36		
37	515	
38		
39	516	
40		
41	517	
42		
43	518	
44		
45	519	
46		
47	520	
48		
49		
50		
51		
52		
53		
54		
55		
56		
57		
58		
59		
60		
61		
62		
63		
64		
65		

Figure Legends

Figure 1. Deterministic model of gait speed outlining the direct relationships between the spatio-temporal determinants and gait speed.

Figure 2. Consort flow diagram of the recruitment process within the nursing home facilities.

Figure 1

Figure 1. Deterministic Model of Gait Speed Outlining the Direct Relationships Between the Spatio-Temporal Determinants and Gait Speed.

Figure 2

Figure 2. Consort Flow Diagram of the Recruitment Process Within the Nursing Home Settings.

Table 1: Spatio-temporal gait determinants and definitions as defined by the GaitMat II manual.

Spatio-temporal gait determinants	Definitions
Step length	The distance from the first switch closure of one footprint to that of the footprint on the contralateral side. ³⁹
Stride length	The distance from the first switch closure of one footprint to the next footprint on the ipsilateral side. ³⁹
Support base	The medial lateral distance across the mat to the innermost switch closure for one footprint from the innermost switch closure of the previous footprint on the contralateral side. ³⁹
Step time	The time to the earliest switch closure of a footfall from the earliest switch closure of the previous footfall on the contralateral side. ³⁹
Swing time	The time to the earliest switch closure of a footfall from the latest switch opening of the previous footfall on the ipsilateral side. ³⁹
Stance time	The time to the latest switch opening of a footfall from the earliest switch closure of the same footfall. ³⁹
Single support time	The time to the earliest switch closure of the next footfall on the contralateral side from the latest switch opening of the previous footfall on the contralateral side. ³⁹
Double support time	The time to the latest switch opening of the previous footfall on the contralateral side from the earliest switch closure of a footfall. ³⁹

Table 2: Characteristics of the Cohort of 100 Nursing Home Residents.

Parameter	Group Mean (SD)	Females (n=67) Mean (SD)	Males (n=33) Mean (SD)
Age, y	85.7 (7.1)	86.1 (6.6)*	85.0 (8.1)
Handgrip Strength, kg	11.1 (4.9)	10.7 (4.2)	11.7 (6.2)
Mini COG, #	1.2 (0.4)	1.8 (0.4)*	1.3 (0.5)
SARC-F, #	5.5 (3.3)	4.9 (3.3)	6.6 (3.1)
Medical Conditions, #	11.0 (4.9)	11.4 (4.9)	10.2 (4.9)
Medications, #	14.0 (5.8)	13.8 (6.1)	14.4 (5.3)
Gait speed, m/s	0.63 (0.19)	0.65 (0.20)	0.58 (0.16)
Step length, m	0.41 (0.08)	0.42 (0.07)	0.41 (0.07)
Stride length, m	0.83 (0.15)	0.84 (0.16)	0.81 (0.14)
Support base, m	0.15 (0.06)	0.16 (0.06)	0.15 (0.07)
Step time, s	0.66 (0.12)	0.64 (0.12)	0.70 (0.12)
Swing time, s	0.42 (0.07)	0.41 (0.07)	0.44 (0.08)
Stance time, s	0.91 (0.20)	0.88 (0.19)	0.98 (0.20)
Single support Time, s	0.42 (0.07)	0.41 (0.06)	0.43 (0.08)
Double support Time, s	0.24 (0.07)	0.23 (0.06)	0.27 (0.08)

= number; Mini COG = Mini Cognitive test; SARC-F = Sarcopenia Five-Item

Questionnaire

* = Statistical significance $p < 0.05$.

Table 3: Univariable and Multivariable Linear Regression Model of the Spatio-Temporal Predictors for Habitual Gait Speed in 100 Residents Living in Nursing Homes.

Factor	Univariable		Multivariable	
	Coefficient (95% CI)	p-value	Coefficient (95% CI)	p-value
Stride length, m	0.93 (0.55 – 1.31)	< 0.001	0.83 (0.74 – 0.92)	< 0.001
Support base, m	-0.51 (-0.77 to -0.26)	< 0.001	-0.44 (-0.68 to -0.21)	< 0.001
Step time, s	-0.70 (-1.13 to -0.27)	0.002	-0.92 (-1.03 to -0.81)	< 0.001

CI = Confidence Interval.

R2 of multiple regression equals 0.892.

Note: All results significant $p<0.05$.

[Click here to access/download](#)

Supplemental Data File (.doc, .tif, pdf, etc.)
coi_disclosure - fien.pdf