Influence of Inpatient Dietary Restriction on Recovery from and Reoccurrence of Acute, Uncomplicated Diverticulitis

Crichton, Megan; Nucera, Romina; Jenkins, Julie; Canavan, Russell; Mahoney, Sophie; Eberhardt, Fiona; Dalwood, Phoebe; Dahl, Camilla ; Marshall, Skye

Published in:
Clinical Nutrition

DOI:
[10.1016/S0261-5614\(19\)32194-6](https://doi.org/10.1016/S0261-5614(19)32194-6)

Licence:
CC BY-NC-ND

[Link to output in Bond University research repository.](#)

Recommended citation(APA):
Crichton, M., Nucera, R., Jenkins, J., Canavan, R., Mahoney, S., Eberhardt, F., Dalwood, P., Dahl, C., & Marshall, S. (2019). Influence of Inpatient Dietary Restriction on Recovery from and Reoccurrence of Acute, Uncomplicated Diverticulitis. *Clinical Nutrition*, 38(S1), S191. [MON-PO360]. [https://doi.org/10.1016/S0261-5614\(19\)32194-6](https://doi.org/10.1016/S0261-5614(19)32194-6)

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

For more information, or if you believe that this document breaches copyright, please contact the Bond University research repository coordinator.

Influence of inpatient dietary prescription on recovery from and reoccurrence of acute, uncomplicated diverticulitis

A 3-year prospective cohort study

skye_marshall@bond.edu.au; @DrSkyeMarshall

The background

Internationally, physicians and surgeons continue to prescribe restricted diets for inpatient management of acute, uncomplicated diverticulitis.

Recent GRADE Clinical Recommendations^a concluded that a liberalized diet (i.e. no dietary restriction) appears safe for the inpatient management of acute, uncomplicated diverticulitis; but more research is needed.

Recent qualitative research^b suggests dietary restrictions can cause long-term psychological pain in diverticulitis patients.

“Clear fluids for 24hrs then upgrade to full fluids for 24hrs, followed by a low-fibre diet until discharge”

The purpose

This study aims to assess the impact of restricted versus liberalized inpatient dietary prescription on recovery from and reoccurrence of acute, uncomplicated diverticulitis.

The methods

Prospective observational study of adult patients admitted to two metropolitan hospitals in Queensland, Australia, from 2016-2019. Hierarchical multiple linear regression was used to determine the impact of restricted versus liberalized diets on hospital length of stay (a proxy for recovery). Binomial logistic regression was used to determine the impact of restricted versus liberalized diets on 30-day and 6-month reoccurrence of diverticulitis.

The participants

- 57% female
- Age: $\mu 59.0 \pm 13.7$ yrs
- BMI: $\mu 27.8 \pm 4.6$ kg/m²
- Alcohol: $\mu 1.4 \pm 2.7$ standard drinks/day
- Admission body temp: $\mu 36.9 (0.7)$ Celsius
- 34% admitted with diarrhoea
- 31% admitted with constipation
- 0% treatment failure
- 45% Restricted Diet
- 55% Liberalised diet
- 55% were prescribed ≥ 3 different diet codes during admission
- 100% were discharged on a “low dietary fibre” diet
- $\mu 36 \pm 19$ (scale 0-105, higher worse) Gastrointestinal symptoms

N=91 Participants

Dietary prescription didn't affect diverticulitis reoccurrence!

Neither liberalized nor restricted diets were predictors of risk of diverticulitis reoccurrence at 30-days or 6-months post-discharge ($p > 0.05$).

Prescribing a liberalized diet was associated with decreased hospital length of stay (indicator of recovery) by 28 hours! ($p < 0.001$)

In a model adjusted for patient characteristics, clinical presentation, and gastrointestinal symptoms, prescribing a liberalized diet decreased hospital length of stay (recovery) by 17 hours! ($p < 0.001$)

The key message

This study confirms and strengthens existing evidence^a: Adult patients admitted to hospital with acute, uncomplicated diverticulitis (i.e. no perforation, non-localized abscess, drains placed or surgery required) should be placed on a liberalized diet (i.e., allowing consumption of solid food) and not placed on a restricted diet (i.e., bowel rest/nil by mouth, clear or full fluids).