Consumers' questions about antipsychotic medication: revealing safety concerns and the silent voices of young men

Weersink, Rianne A.; Taxis, Katja; McGuire, Treasure M.; van Driel, Mieke L.

Published in:
Social Psychiatry and Psychiatric Epidemiology

DOI:
[10.1007/s00127-014-1005-y](https://doi.org/10.1007/s00127-014-1005-y)

Licence:
Other

[Link to output in Bond University research repository.](#)

Recommended citation(APA):
Weersink, R. A., Taxis, K., McGuire, T. M., & van Driel, M. L. (2015). Consumers' questions about antipsychotic medication: revealing safety concerns and the silent voices of young men. *Social Psychiatry and Psychiatric Epidemiology*, 50(5), 725-733. <https://doi.org/10.1007/s00127-014-1005-y>

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

For more information, or if you believe that this document breaches copyright, please contact the Bond University research repository coordinator.

Title Page

Title: Consumers' questions about antipsychotic medication: revealing safety concerns and the silent voices of young men

Authors:

- Rianne A. Weersink
 - o Department of Pharmacy, Unit of Pharmacotherapy and Pharmaceutical Care, University of Groningen, Antonius Deusinglaan 1, 9713 AV Groningen, The Netherlands
 - o Discipline of General Practice, School of Medicine, The University of Queensland, Level 8 Health Sciences Building, Herston QLD 4029, Brisbane, Australia
- Katja Taxis
 - o Department of Pharmacy, Unit of Pharmacotherapy and Pharmaceutical Care, University of Groningen, Antonius Deusinglaan 1, 9713 AV Groningen, The Netherlands
- Treasure M. McGuire
 - o Mater Pharmacy Services, Level 3, Mater Children's Hospital, Mater Health Services, South Brisbane, 4101, Australia
 - o School of Pharmacy, PACE Precinct, Building 870, The University of Queensland QLD 4072, Australia
 - o Faculty of Health & Medical Sciences, Bond University, Gold Coast, 4229, Australia
- Mieke L. van Driel
 - o Discipline of General Practice, School of Medicine, The University of Queensland, Level 8 Health Sciences Building, Herston QLD 4029, Brisbane, Australia, Phone: +61 7 336 55144, Fax: +61 7 3346 5178, m.vandriel@uq.edu.au

Word count

Abstract: 251 words

Text: 3,245 words

Abstract

Purpose: Little is known about consumer information needs regarding antipsychotic medicines. Medicines call centre (MCC) derived data are underutilised; and could provide insight into issues of importance to consumers. This study aimed to explore consumers' information needs about antipsychotic medication sought from a national MCC in Australia.

Methods: Questions received by the National Prescribing Service (NPS) *Medicines Line* relating to antipsychotic medication from September 2002 to June 2010 were examined by antipsychotic subclass and in relation to other medication queries.

Results: We identified 6,295 calls related to antipsychotic medication. While female callers predominated, the percentage of males with antipsychotic questions was statistically significantly higher than for other medication calls (33.9% vs 22.6%; $p < 0.001$). There were distinct gender differences in medicines information seeking across age ranges. Younger men asked about second-generation antipsychotics, shifting towards first-generation antipsychotics after 45 years of age. Female interest in both subclasses was comparable, irrespective of age. Most callers asking about antipsychotics sought information for themselves (69.4%). Callers were primarily concerned about safety (57.0%), especially adverse drug reactions (28.8%), and were more often prompted by a worrying symptom (23.8%) compared with the rest of calls (17.2%). Trends of antipsychotic questions received corresponded with antipsychotic prescription data.

Conclusions: The number of calls received by this MCC over time reveals an ongoing consumer need for additional, targeted information about antipsychotics. Noticeable was the relatively high frequency of young male callers asking about antipsychotics, indicating that call centres could be a way to reach these traditionally poor users of health services.

Keywords: antipsychotic agents, information seeking behaviour, consumer health information, medicines call centre, health information seeking, medicines information seeking

Introduction

Mental illness constitutes a large and increasing burden of disease in Australia and worldwide [1]. Psychotic disorders, in particular schizophrenia that requires lifelong treatment, contribute significantly to health burden and costs [2]. Prescribing of antipsychotics has increased markedly in Europe, North America and Australia in recent years [3-5]. It is therefore not surprising that consumers' need for information about antipsychotics is also growing.

When looking for information about medicines, consumers use a diverse range of information sources such as their doctor, pharmacist, the internet or consumer medication information (CMI) leaflets [6-8]. Another valuable resource is a medicines call centre (MCC) [9-11]. Information provided with prescription medicines is often perceived as inadequate by consumers [12]; and a MCC can be helpful to fill the information gap. Although there is extensive literature covering the area of *health* information seeking behaviour [13-15]; studies about *medicines* information seeking are limited. There is, specifically, a lack of studies about consumers' antipsychotic medication information needs. Data derived from a MCC is an underutilised resource and could provide insight into antipsychotic-related issues of concern to consumers. The aim of this study was to profile antipsychotic medication information seekers and explore their information needs, as presented to a national MCC in Australia.

Methods

Study site

Medicines Line is a national MCC, funded by the National Prescribing Service (NPS), to promote quality use of medicines by providing information that is independent, evidence-based and safe. Consumers could access the service for information about prescription, over-the-counter or complementary medicines for the cost of a local call, Monday to Friday 9am to 6pm, from anywhere in Australia, or leave a voicemail for a return call or to request consumer medicines information leaflets. Callers were also encouraged to discuss the information provided with their doctor or pharmacist. While marketing was primarily via the NPS website, the service was promoted in the media as part of National Medicines Week in 2007 and 2008. Between its launch in September 2002 and 30 June 2010, the service was operated by experienced clinical pharmacists at Mater Health Services, Brisbane.

Information collected during the call was recorded by the pharmacist on a standardised, scannable form, with the call narrative entered into a Microsoft Access™ database, linked by the unique scanform number. Information included demographic data of the caller, the patient and the relationship between caller and patient. Details of questions were collected, such as enquiry type and motivation for calling.

Medicines involved in the call were recorded and categorised into major groups, based on the Anatomical Therapeutic Chemical (ATC) classification of the World Health Organization [16]. Major groups were further divided into minor therapeutic classes, based on the Monthly Index of Medical Specialties (MIMS) Australia classification [17]. MIMS is a source for medicine's product information, available for Australian healthcare professionals.

Data extraction

1 Calls in the database were classified as 'antipsychotic-related calls', where the question
2 concerned a medication from the minor group 'antipsychotic agents'. The remainder of calls about
3 medication from all other minor and major groups were classified as 'rest of calls'.
4
5

6
7 As the focus of the MCC was service provision, only three demographic variables relating to the
8 caller and/or patient were requested: gender, age and postcode. Age was categorised into nine age
9 groups, based on the Australian Bureau of Statistics National Health Survey [18]. Caller postcodes
10 were grouped by state or territory, relative population frequency and by Accessibility/Remoteness
11 Index of Australia (ARIA), i.e. a measure of remoteness (from service centres) for all places in
12 Australia [19].
13
14
15
16
17
18
19
20

21 The following question characteristics were also extracted from the database:
22

- 23 • Date of call
- 24
25 • Enquiry type: Category of question asked by the caller. Only one of twenty options could be
26 chosen by the pharmacist handling the call (e.g. administration, adverse drug reaction (ADR),
27 interaction, dose, pregnancy). These items were collapsed into four main groups; safety,
28 efficacy, judicious use and other.
29
30
31
32
33
34
35 • Prompt: Callers were asked what motivated them to contact the service. This response was
36 categorised into one of nine drivers of medicines information seeking behaviour including
37 inadequate information, conflicting information, second opinion and worrying symptom.
38
39
40
41
42 • Medicine(s) involved in the query: Although the patient may have been taking multiple
43 medicines, a maximum of three were entered into the database in order of relevance to the
44 question. While in some cases, e.g. interaction questions, all three medicines (including one or
45 more antipsychotics) were equally relevant to the question, the first medicine entered was
46 assigned as the 'primary' medicine. Antipsychotic-related calls were grouped into the subclasses
47 of 'first-generation antipsychotics' and 'second-generation antipsychotics' based on this
48 primary medicine. Lithium, also categorised as an antipsychotic according to the ATC and MIMS-
49 classifications [16,17], was excluded from the group of antipsychotic-related calls as it has a
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

different and unique mechanism of action. Calls related to lithium have been included in the group 'rest of calls'.

- Prior information source: Information source(s) the caller had consulted prior to contacting *Medicines Line* (e.g. doctor, pharmacist, other health professional, support group, media, internet or consumer medicines information (CMI)).

Analyses

The MCC service Access™ database was exported to SPSS® version 21 for statistical analyses. Frequency tables were computed for caller demographics and question characteristics; these were compared by antipsychotic subclass. Data from antipsychotic-related calls were also compared with the rest of calls in the database. A chi-square test was used to compare dichotomous and categorical variables and an independent sample t-test was used to compare continuous variables. A p-value of ≤ 0.05 was applied to indicate statistical significance.

In addition, for the primary medicine of interest, the temporal relationship between antipsychotic classes of interest to consumers in this cohort relative to the dispensing of antipsychotic prescriptions (actual medicine use) was explored. Comparisons were made between annual calls about antipsychotics (first- and second-generation antipsychotics) and national prescription data from the Australian Statistics on Medicines (ASM) which provides annual dispensed prescription totals for medicines, based on a combination of Pharmaceutical Benefits Scheme (PBS) data and community pharmacy data [20]. Percentages per antipsychotic subclass were calculated using the total number of *Medicines Line* calls and the sum of all dispensed item types for that class in ASM, for 2002 to 2010 inclusive, as denominators.

Ethical Approval

This study was undertaken as part of a program of research that has received 'Low and Negligible Risk' ethical approval from the Human Research Ethics Committee, Mater Health Services Brisbane Limited (HREC Protocol Ref No: LNR submission 2012-68).

Results

A total of 125,919 calls were available for analysis concerning 195,855 medications. Antipsychotic-related calls accounted for 5.0% (n=6,295) of total calls. Of these calls, most (89.1%) involved a question about a single antipsychotic while the remaining 10.9% included two or more antipsychotics. This equates to an average of 1.1 (mode 1; range 1-3) antipsychotic medications per enquiry.

Among the 6,295 calls involving at least one antipsychotic, an antipsychotic was the primary medicine in 4,398 calls. This primary medicine of interest was a second-generation antipsychotic in 81.0% (n=3,564) of these calls, most commonly olanzapine, quetiapine and risperidone. Of the 834 calls (19.0%) where a first-generation antipsychotic was the primary medicine, the focus was on chlorpromazine, haloperidol and pericyazine.

In Figure 1, the calls received by *Medicines Line* for subgroups of antipsychotics (first- and second-generation) per year is compared to the sum of all prescriptions dispensed for these subgroups in Australia. This comparison shows a similar trend over time between the percentage of calls per antipsychotic subgroup (medicine of interest in this cohort) and the percentage of prescriptions dispensed per antipsychotic subgroup from 2002 to 2010 (national medication use) [20].

Besides antipsychotics, other medications frequently involved in antipsychotic-related questions were sodium valproate (5.6% of antipsychotic calls), diazepam (4.6%) and lithium (4.0%) – all these agents act on the nervous system.

Characteristics of callers

Table 1 lists characteristics of callers to the service. While females were the predominant caller, the percentage of male callers with questions about antipsychotics (33.9%) was significantly higher compared with the 22.6% of males calling for other medication ($p<0.001$). The mean age of callers for antipsychotic-related queries was 46.0 years (SD 14.6; range 16-98). They were significantly younger ($p<0.001$) than callers for the 'rest of calls' (mean age 50.8 years, SD 17.9; range 5-100).

1 Approximately 70% of callers for both antipsychotic and other medication information rang on
2 behalf of themselves. The remainder called for a significant 'other', mostly a family member. Calls
3 originated from all states and territories in Australia. Postcode classification by relative call
4 frequency and ARIA [21,22] revealed that service uptake was equally distributed across all areas of
5 accessibility and remoteness, with people calling from 'remote' areas approaching the relative
6 population frequency.
7

8 Age and gender of callers for antipsychotic-related questions (per subgroup) were compared with
9 those of callers for other medication (Figure 2). The mean caller age for first-generation
10 antipsychotics agents was 48.5 years (SD 14.8) compared to 45.4 years (SD 14.5) for second-
11 generation antipsychotics ($p<0.001$). However, even though the ratio of female to male callers
12 asking about second-generation or first-generation antipsychotics remained fairly consistent
13 (approximately 65%:35%; $p=0.934$), there were distinct gender differences in medicines information
14 seeking across age ranges. Younger men, 15 to 44 years, displayed a strong interest in second-
15 generation antipsychotics, shifting towards first-generation antipsychotics after 45 years of age.
16 Female interest in both subclasses was comparable, irrespective of age.
17

18 When comparing data from the subgroups of antipsychotics with the rest of calls (Figure 2),
19 several key issues emerged. For the rest of calls, the majority of callers were female for each age
20 group. This is in contrast with antipsychotic-related calls, in particular with calls about second-
21 generation antipsychotics. The main difference was in the 25-34 year age group, where the
22 proportion of males with second-generation antipsychotic queries was 10.9% compared with the
23 3.2% of males asking about other medication. Female callers requesting antipsychotic information
24 were generally older, most often 45-54 years old.
25

26 Reasons for calling

27 Table 2 lists the motivation for calling the MCC. Across all calls, the major reasons were
28 inadequate information, a worrying symptom or requiring a second opinion. Antipsychotic-related
29

calls were more often prompted by a worrying symptom (23.8%; $n=1,491$) than the rest of calls (17.2%; $n=20,442$) ($p<0.001$). There was no difference in the information needs of people calling for themselves or for another person.

Prior information source

Callers ringing about antipsychotic medication ($n=6,295$) cited 7,645 information sources used before contacting *Medicines Line* compared with the remaining 119,624 callers who nominated 143,271 sources. This suggests that consumers frequently use more than one source when seeking medicines information. In both cohorts, health professionals were important 'prior information sources', consulted in more than half of calls. The doctor was the main prior information source (40.6% for antipsychotic-related calls ($n=2,557$) and 33.2% for rest of calls ($n=39,681$)).

Narrative of enquiries

Medication safety was the main caller concern, accounting for almost 60% of calls ($n=3,578$) about antipsychotics (Table 2). Noticeable from individual enquiry types was the high number of calls received about adverse drug reactions (ADRs), with a frequency of 28.8% ($n=1,810$) for antipsychotic-related calls versus 18.7% ($n=22,280$) for rest of calls ($p<0.001$). The frequencies of other individual enquiry types were reasonably consistent compared to other medicines.

When call narrative was explored for key themes, safety-related questions focused on side-effects, risks in special populations and interactions. Some examples:

- 'Are any of my medicines (sertraline, haloperidol) causing my drowsiness and bleeding nose?'
- 'Will I get addicted to Zyprexa (olanzapine)?'
- 'What should I do to treat the constipation caused by my chlorpromazine and quetiapine?'
- 'How long will my side effects of pain in the legs and sedation due to paliperidone last?'

Questions about the risks in special populations included 'Would I be able to breastfeed my new born baby if I have been on olanzapine throughout my pregnancy and will most likely continue on it after

baby is born?'; while interaction questions often involved another psychiatric medication,
complementary or lifestyle drug e.g. 'Can my son take Aurorix (moclobemide) if he is on Zyprexa
(olanzapine)?' or 'Do my complementary medicines interact with olanzapine?' Safety concerns
about antipsychotics were followed by a desire for information on medication effectiveness. For
instance: 'How quickly will Neulactil (pericyazine) start to work?', or 'Would ziprasidone be a better
drug than olanzapine with regards to weight gain and diabetes?'

Discussion

Our study profiled antipsychotic medication information seekers and identified an apparent unmet need for information about antipsychotic medicines, particularly in young males.

Characteristics of (antipsychotic) medication information seekers

The profile of callers asking about medicines in general and antipsychotics in particular is predominantly female, seeking information for themselves or a significant other. This is consistent with the literature suggesting that women tend to seek health information more often than men [13,14,23], and for either themselves or a 'family' member [24]. It also aligns with data from other MCCs [25,26]. However, our study found a relatively high proportion of (young) males calling *Medicines Line* for information about antipsychotics. The literature is less clear in relation to caller age. Some studies about health seeking behaviour suggest that younger individuals are more likely to health seek [14,27,28] while others note a positive relationship with increasing age [13,29]. In our study, while callers seeking information on antipsychotics were spread across a wide age range supporting both paradigms, they were, on average, a younger cohort. This could be explained by the age of onset of psychosis, which is often in early adulthood [30]. This is also when affected individuals are having their first experiences of taking antipsychotic medications, potentially triggering information seeking behaviour.

Studies suggest that young males are less likely to help seek [13,29] or access health services [29], in particular for mental health disorders [29,31,32]. However, it is unlikely that they have fewer questions than, for instance, young women. Perhaps they are not getting the answers they need through conventional channels? It is possible that young men feel more comfortable contacting a call centre instead of directly facing a healthcare professional. This is in line with mental health patients' preference for anonymous sources for health and medicines information [33-35]. Consultation via a telephone service could therefore be a way for them to voice questions about

their health and for health services to reach this group of consumers. Further research is needed to confirm and explore this finding.

MCCs as a source for (antipsychotic) medication information

In a country as vast as Australia, there are challenges of providing equitable access to quality health services in rural and remote areas. A medicines information service with direct phone access to a pharmacist has the potential to overcome the tyranny of distance and meet the needs of rural as well as urban consumers, independent of location. However, this does not always equate to actual service utilisation. In this study, calls relating to antipsychotics not only originated from all Australian states and territories, but also importantly, people calling from 'remote' areas approached the relative population frequency. This supports that the MCC service model had utility for mental health consumers in rural and remote areas where there is likely to be less access to mental health services.

Although the two variables in Figure 1 use different denominators, over an eight year period, there was a close parallel between the percentage of calls about antipsychotic subgroups (medicine of interest in this cohort) and the percentage of antipsychotic prescriptions dispensed annually (national medicine use). This longitudinal perspective, together with the quantum of calls received over time, reveals the ongoing need consumers have for quality information about (all subgroups of) antipsychotic medication. This need was also recognized in other MCC studies [25,26].

Link between uncertainty and motivation to seek medicines information

Uncertainty, related to an information gap or a symptom of concern, has been shown to drive health seeking behaviours such as seeking medicines information [13,14,36,37]. Inadequate information was the main reason for contacting this MCC, in both antipsychotic-related and the remainder of calls. Consumers receiving inadequate information tend to be dissatisfied with the

information available to them and want to know more, supporting the existence of a knowledge gap for almost half of those seeking information on antipsychotics [38-40].

The second main motivation for calling the MCC with antipsychotic medication questions was a 'worrying symptom'. The higher number of questions prompted by a worrying symptom in relation to antipsychotics compared with 'rest of calls' is supported by the frequent occurrence of adverse effects with antipsychotics [41]. In almost half of total antipsychotic-related calls, the prior information source previously consulted was not a health care professional but, for instance, the internet. It is of concern that this cohort had not sought help from their health carers about this worrying symptom prior to contacting the service. The call, however, provided an opportunity for a MCC pharmacist to decrease caller uncertainty about their medication-related issue, and appropriately triage callers requiring review by their doctor. Another important prompt in both types of calls was 'second opinion'. Studies that assessed the motivation of people to seek health information on the internet also highlighted 'second opinion' as a prompt for health seeking behaviour - the hope to find something 'new' [42,43]. Since there is limited literature available about motivation for health seeking behaviour, these findings could provide a foundation for further research.

Narrative of medicine information seeking

Safety was the main concern of callers with those asking for information about antipsychotics more frequently wanting information about adverse drug reactions than callers for other medication. Safety as a goal of medicines information seeking is in agreement with literature about medicines information needs of consumers [8,44] and with findings from other MCCs [9,26,45]. Medicines information provided by healthcare professionals or accompanying medication, such as the Consumer Medicines Information (CMI) leaflet, should therefore focus more on safety issues, in particular strategies to minimise or monitor risk. Antipsychotics are medications with a high side-effect burden [41], which can be worrying and trigger health seeking behaviour. Qualitative studies

1 involving mental health patients revealed a knowledge gap among patients, especially about adverse
2 drug reactions from antipsychotics [38-40]. Patients experiencing these tend to be less compliant
3 with their therapy [46,47]. It is, therefore, important that such questions from consumers are
4 answered *before* their concerns escalate and they cease their therapy. Further research should be
5 aimed at assessing the specific information needs of consumers about adverse effects of
6 antipsychotics and their preferences for receiving this information.
7
8
9
10
11
12
13
14
15

16 Limitations

17 The *Medicines Line* service model was developed as a cost-effective strategy to meet the
18 medicines information needs of Australian consumers; it was not created as a data repository for
19 research. As a result, there are missing data, especially in relation to the voicemail requests for CMI
20 leaflets. The percentage of missing data is however not more than 5% for every characteristic and is
21 reasonably comparable for both antipsychotic-related calls and rest of calls.
22
23
24
25
26
27
28
29

30 An addition, callers to a MCC are only part of the total group of medicine users. We do not know
31 if the non-health information seeking consumer wants more or different information about their
32 medicines. However, our study includes a significant proportion of questions from young males
33 (often direct users of the antipsychotic medicines), who are known to be infrequent users of health
34 services.
35
36
37
38
39
40
41
42
43
44

45 Conclusion

46 The number of calls received by this MCC reveals an ongoing need for additional information on
47 antipsychotics from consumers. Noticeable was the relatively high frequency of young males
48 contacting the MCC for safety information about antipsychotic medication, a cohort known to be
49 poor users of health services. Acknowledging their preference for telephone consultation can help
50 reach this group. Overall, MCCs are a valuable and underutilised repository for research on
51 consumer medicines information seeking behaviour.
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Acknowledgements

We would like to acknowledge NPS MedicineWise (formerly National Prescribing Service, Australia), funder of NPS *Medicines Line* and service provider since July 2010. We would also like to thank Mater Health Services for providing the raw service data from September 2002 to 30 June 2010; and Gabrielle Hartley, Mater Pharmacy Services, for database assistance.

This study was supported by a travel scholarship for Rianne Weersink from the Chiel Hekster Fund and the Stipendium Fund of the Koninklijke Nederlandse Maatschappij ter bevordering der Pharmacie (KNMP), The Netherlands.

Conflict of Interest

The sponsors had no role in study design; in the collection, analysis and interpretation of data; in the writing of the report; and in the decision to submit the paper for publication. On behalf of all authors, the corresponding author states that there is no conflict of interest.

References

1. World Health Organisation (2001) The World Health Report 2001: Mental Health - New Understanding, New Hope. World Health Organisation, Geneva
2. Rössler W, Joachim Salize H, van Os J, Riecher-Rössler A (2005) Size of burden of schizophrenia and psychotic disorders. *Eur Neuropsychopharmacol* 15 (4):399-409
3. Mental health services in Australia (2012) Mental-health related prescriptions. Australian Institute of Health and Welfare. <http://mhsa.aihw.gov.au/resources/prescriptions/>. Accessed 8 December 2012
4. Data and Modelling Section Pharmaceutical Policy and Analysis Branch (2011) Expenditure and prescriptions twelve months to 30 June 2011. Pharmaceutical Benefit Scheme. <http://www.pbs.gov.au/info/statistics/expenditure-and-prescriptions-30-06-2011>. Accessed 21 January 2013
5. Verdoux H, Tournier M, Begaud B (2010) Antipsychotic prescribing trends: a review of pharmaco-epidemiological studies. *Acta Psychiatr Scand* 121 (1):4-10. doi: 10.1111/j.1600-0447.2009.01425.x
6. Tio J, LaCaze A, Cottrell WN (2007) Ascertaining consumer perspectives of medication information sources using a modified repertory grid technique. *Pharm World Sci* 29 (2):73-80
7. Närhi U (2007) Sources of medicine information and their reliability evaluated by medicine users. *Pharm World Sci* 29 (6):688-694
8. Ho CH, Ko Y, Tan ML (2009) Patient needs and sources of drug information in Singapore: is the Internet replacing former sources? *Ann Pharmacother* 43 (4):732-739. doi: 10.1345/aph.1L580
9. Melnyk P, Shevchuk Y, Remillard A (2000) Impact of the dial access drug information service on patient outcome. *Ann Pharmacother* 34 (5):585-592
10. Bertsche T, Hammerlein A, Schulz M (2007) German national drug information service: user satisfaction and potential positive patient outcomes. *Pharm World Sci* 29 (3):167-172

11. Marvin V, Park C, Vaughan L, Valentine J (2011) Phone calls to a hospital medicines information helpline: analysis of queries from members of the public and assessment of potential for harm from their medicines. *Int J Pharm Pract* 19 (2):115-122. doi: 10.1111/j.2042-7174.2010.00081.x
12. Svarstad BL, Mount JK, Tabak ER (2005) Expert and consumer evaluation of patient medication leaflets provided in US pharmacies. *J Am Pharm Assoc* 45:443-451
13. Dutta-Bergman MJ (2003) Developing a profile of consumer intention to seek our health information beyond a doctor. *Health Mark Q* 21 (1-2):91-112
14. Lambert SD, Loiselle CG (2007) Health information-seeking behavior. *Qual Health Res* 17(8):1006-1019
15. Anker AE, Reinhart AM, Feeley TH (2011) Health information seeking: A review of measures and methods. *Patient Educ Couns* 82 (3):346-354. doi: <http://dx.doi.org/10.1016/j.pec.2010.12.008>
16. World Health Organization Collaborating Centre for Drug Statistics Methodology (2010) Guidelines for ATC classification and DDD assignment, 2011. World Health Organisation, Oslo17.
- UBM Medica (2012) MIMS. MIMS Australia. <http://www.mims.com.au/index.php/about-mims/mims-australia>. Accessed 27 September 2012
18. ABS (2009) National Health Survey: Summary of Results, 2007-2008. Australian Bureau of Statistics. <http://www.abs.gov.au/AUSSTATS/abs@.nsf/mf/4364.0> Accessed 21 January 2013
19. Information and Research Branch Department of Health and Aged Care (2001) Measuring Remoteness: Accessibility/Remoteness Index of Australia (ARIA) New Series, vol Number 14. Department of Health and Ageing, Canberra
20. Australian Government Department of Health and Ageing (2004-2012) Australian statistics on medicines 2002-2010. Department of Health and Ageing. <http://www.pbs.gov.au/info/browse/statistics>. Accessed 4 February 2013
21. ABS (2012) Year Book Australia 2012; population and growth 2010. Australian Bureau of Statistics. <http://www.abs.gov.au/ausstats/abs@.nsf/mf/1301.0>. Accessed 11 February 2013

22. ABS (2007) Australian Standard Geographical Classification (ASGC) Remoteness Structure (RA) Digital Boundaries, Australia, 2006. Australian Bureau of Statistics, Canberra
23. Smith JA, Braunack-Mayer A, Wittert G (2006) What do we know about men's help-seeking and health service use? *Med J Aust* 184 (2):81-83
24. Warner D, Procaccino JD (2004) Toward wellness: Women seeking health information. *J Am Soc Inform Sci Technol* 55 (8):709-730
25. Pohjanoksa-Mantyla MK, Antila J, Eerikainen S, Enakoski M, Hannuksela O, Pietila K, Airaksinen M (2008) Utilization of a community pharmacy-operated national drug information call center in Finland. *Res Social Adm Pharm* 4 (2):144-152. doi: 10.1016/j.sapharm.2007.05.001
26. Huber M, Kullak-Ublick GA, Kirch W (2009) Drug information for patients—an update of long-term results: type of enquiries and patient characteristics. *Pharmacoepidemiol Drug Saf* 18 (2):111-119. doi: 10.1002/pds.1682
27. Gourash N (1978) Help-seeking: a review of the literature. *Am J Community Psychol* 6 (5):413-423
28. Ramanadhan S, Viswanath K (2006) Health and the information nonseeker: a profile. *Health Commun* 20 (2):131-139
29. Mackenzie CS, Gekoski WL, Knox VJ (2006) Age, gender, and the underutilization of mental health services: the influence of help-seeking attitudes. *Aging Ment Health* 10 (6):574-582
30. Kessler RC, Amminger GP, Aguilar-Gaxiola S, Alonso J, Lee S, Ustun TB (2007) Age of onset of mental disorders: a review of recent literature. *Curr Opin Psychiatry* 20 (4):359-64
31. Galdas PM, Cheater F, Marshall P (2005) Men and health help-seeking behaviour: literature review. *J Adv Nurs* 49 (6):616-623
32. Biddle L, Gunnell D, Sharp D, Donovan JL (2004) Factors influencing help seeking in mentally distressed young adults: a cross-sectional survey. *Br J Gen Pract* 54 (501):248-253
33. Pohjanoksa-Mäntylä M, Bell JS, Helakorpi S, Närhi U, Pelkonen A, Airaksinen M (2011) Is the Internet replacing health professionals? A population survey on sources of medicines information

among people with mental disorders. Soc Psychiatry Psychiatr Epidemiol 46 (5):373-379. doi:
10.1007/s00127-010-0201-7

34. Powell J, Clarke A (2006) Internet information-seeking in mental health. Br J Psychiatry 189
(3):273-277

35. Khazaal Y, Chatton A, Cochand S, Hoch A, Khankarli MB, Khan R, Zullino DF (2008) Internet use by
patients with psychiatric disorders in search for general and medical informations. Psychiatr Q 79
(4):301-309. doi: 10.1007/s11126-008-9083-1

36. McGuire TM (2005) Consumer medicines call centres: a medication liaison model of
pharmaceutical care. Dissertation, University of Queensland

37. Rosen NO, Knäuper B (2009) A little uncertainty goes a long way: State and trait differences in
uncertainty interact to increase information seeking but also increase worry. Health Commun 24
(3):228-238. doi: 10.1080/10410230902804125

38. Happell B, Manias E, Roper C (2004) Wanting to be heard: mental health consumers' experiences
of information about medication. Int J Ment Health Nurs 13 (4):242-248

39. Powell J, Clarke A (2006) Information in mental health: qualitative study of mental health service
users. Health Expect 9 (4):359-365

40. Gray R, Rofail D, Allen J, Newey T (2005) A survey of patient satisfaction with and subjective
experiences of treatment with antipsychotic medication. J Adv Nurs 52 (1):31-37

41. Weiden PJ, Buckley PF (2007) Reducing the burden of side effects during long-term antipsychotic
therapy: the role of "switching" medications. J Clin Psychiatry 68 (6):14-23

42. Eysenbach G, Diepgen TL (1999) Patients looking for information on the internet and seeking
teleadvice: Motivation, expectations, and misconceptions as expressed in e-mails sent to physicians.
Arch Dermatol 135 (2):151-156

43. Umefjord G, Petersson G, Hamberg K (2003) Reasons for consulting a doctor on the Internet:
Web survey of users of an Ask the Doctor service. J Med Internet Res 5 (4):e26

- 1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
44. Newby DA, Hill SR, Barker BJ, Drew AK, Henry DA (2001) Drug information for consumers: should it be disease or medication specific? Results of a community survey. *Aust N Z J Public Health* 25 (6):564-570
45. Grymonpre R, Steele J (1998) The medication information line for the elderly: an 8-year cumulative analysis. *Ann Pharmacother* 32 (7):743-748
46. Fleischhacker WW, Meise U, Günther V, Kurz M (1994) Compliance with antipsychotic drug treatment: influence of side effects. *Acta Psychiatr Scand* 89:11-15
47. Vermeire E, Hearnshaw H, Van Royen P, Denekens J (2001) Patient adherence to treatment: three decades of research. A comprehensive review. *J Clin Pharm Ther* 26 (5):331-342

Figures (created in Microsoft Excel)

Fig. 1 Comparison of the distribution of subgroups of antipsychotics (first-generation and second-generation) per year for calls received by *Medicines Line* and the sum of all prescriptions dispensed for these subgroups in Australia. Percentages of calls were calculated using the total number of calls per year where an antipsychotic was the primary drug. For prescription data, percentages were calculated using the annual total of defined daily dose (DDD) per 1000 population per day per the Australian Statistics on Medicines [20]. AP = antipsychotic

Fig. 2 Caller gender and age range compared for calls about first-generation antipsychotics, second-generation antipsychotics and rest of calls.

^a Missing data for gender and/or age is omitted from these total number of calls: $n=34$ (4.1% of 834) in first-generation antipsychotic-related calls, $n=104$ (2.9% of 3,564) in second-generation antipsychotic-related calls and $n=2,163$ (1.8% of 119,624) for rest of calls.

Tables

Table 1 Comparison of caller characteristics of antipsychotic-related and rest of calls

		Antipsychotic-related calls n=6,295		Rest of calls n=119,624		Statistics
		%	n	%	n	
Gender (%)	Female	66.1	4,159	77.4	92,452	$\chi^2=430.6$ df=1 $p<0.001$
	Male	33.9	2,136	22.6	27,009	
Total (missing omitted) ^a		100.0	6,295	100.0	119,461	
Missing ^b		0	0	0.1	163	
Age (years)	<5	0	0	0	0	
	5-14	0	0	0	27	
	15-24	4.2	257	4.1	4,834	
	25-34	20.4	1,252	19.4	22,835	
	35-44	23.9	1,465	18.0	21,127	
	45-54	24.1	1,477	15.5	18,262	
	55-64	16.6	1,016	16.4	19,255	
	65-74	6.8	418	14.7	17,252	
	≥75	4.1	252	11.8	13,891	
Total (missing omitted) ^a		100.0	6,137	100.0	117,483	
Missing ^b		2.5	158	1.8	2,141	
Mean age (SD)		46.0 (14.6)		50.8 (17.9)		$T=887.4$ df=123,618 $p<0.001$
Relationship (% self)		69.4	4,365/6,286	72.0	85,917/119,282	$\chi^2=19.8$ df=1 $p<0.001$
Missing		0.1	9	0.3	342	

^a Percentages are total of data available per variable. These may not add exactly to 100.0 due to rounding.

^b Number and percentage of missing data per variable for total antipsychotic-related and rest of calls. Total missing data in rest of calls for gender and/or age = 2163 with data about both gender and age missing for 141 callers.

Table 2 Comparison of the motivation for contacting the MCC and the enquiry type, classified by relationship of the caller to the patient

	Antipsychotic-related calls <i>n</i> =6,295				Rest of calls <i>n</i> =119,624			
	Self		Other		Self		Other	
	%	<i>n</i>	%	<i>n</i>	%	<i>n</i>	%	<i>n</i>
Prompt								
Inadequate information	40.1	1,746	46.4	890	44.8	38,421	50.9	16,956
Worrying symptom	24.3	1,058	22.6	433	18.3	15,663	14.3	4,779
Second opinion	22.4	975	22.3	427	23.0	19,720	23.7	7,904
Conflicting information	3.9	169	4.5	87	5.4	4,600	6.1	2,040
Other	3.8	166	0.5	10	3.0	2,595	0.6	199
Information overload	1.9	82	1.1	22	1.3	1,122	1.0	335
Forgot information	1.6	68	0.8	15	1.5	1,301	1.0	346
Professional information source	1.1	47	0.8	15	1.2	995	1.0	327
Consumer information source	1.1	46	1.0	19	1.6	1,405	1.3	439
Total (<i>missing omitted</i>) ^a	100.0	4,357	100.0	1,918	100.0	85,822	100.0	33,325
Missing ^b	0.3% (<i>n</i> =20)				0.4% (<i>n</i> =477)			
Enquiry Type								
Safety	58.5	2,551	53.5	1,027	53.6	46,003	54.6	18,206
Efficacy	35.9	1,564	36.7	705	33.6	28,787	32.4	10,791
Judicious use	4.4	191	9.2	177	7.9	6,756	9.9	3,289
Other	1.2	52	0.6	11	4.9	4,214	3.1	1,049
Total (<i>missing omitted</i>) ^a	100.0	4,358	100.0	1,920	100.0	85,760	100.0	33,335
Missing ^b	0.3% (<i>n</i> =17)				0.4% (<i>n</i> =529)			

^a Percentages are total of data available per variable. These may not add exactly to 100.0 due to rounding.

^b Number and percentage of calls where the relationship of the caller to the patient (self/other) and/or the prompt/enquiry type of the call is missing.