

Immediate and delayed antibiotic prescribing strategies used by Australian early-career GP: A cross-sectional analysis

Davey, Andrew; Tapley, Amanda; Mulquiney, Katie June; van Driel, Mieke; Fielding, Alison; Holliday, Elizabeth; Davis, Joshua S; Glasziou, Paul; Dallas, Anthea; Ball, Jean; Spike, Neil; FitzGerald, Kristen; Magin, Parker

Published in:

The British journal of general practice : the journal of the Royal College of General Practitioners

DOI:

[10.3399/BJGP.2021.0026](https://doi.org/10.3399/BJGP.2021.0026)

Licence:

CC BY

[Link to output in Bond University research repository.](#)

Recommended citation(APA):

Davey, A., Tapley, A., Mulquiney, K. J., van Driel, M., Fielding, A., Holliday, E., Davis, J. S., Glasziou, P., Dallas, A., Ball, J., Spike, N., FitzGerald, K., & Magin, P. (2021). Immediate and delayed antibiotic prescribing strategies used by Australian early-career GP: A cross-sectional analysis. *The British journal of general practice : the journal of the Royal College of General Practitioners*, 71(713), E895-E903. Advance online publication. <https://doi.org/10.3399/BJGP.2021.0026>

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

For more information, or if you believe that this document breaches copyright, please contact the Bond University research repository coordinator.

Andrew Davey, Amanda Tapley, Katie J Mulquiney, Mieke van Driel, Alison Fielding, Elizabeth Holliday, Joshua S Davis, Paul Glasziou, Anthea Dallas, Jean Ball, Neil Spike, Kristen FitzGerald and Parker Magin

Immediate and delayed antibiotic prescribing strategies used by Australian early-career GPs:

a cross-sectional analysis

Abstract

Background

Antibiotics are overused for non-pneumonia acute respiratory tract infections (ARTIs).

Aim

To establish prevalence and explore associations of delayed and immediate antibiotic prescribing strategies of Australian early-career GPs (specialist GP vocational trainees, also known as GP registrars) for non-pneumonia ARTIs.

Design and setting

Cross-sectional analysis of data collected between September 2016 and December 2017 from the Registrar Clinical Encounters in Training cohort (ReCEnt) study, an ongoing cohort study of GP registrars' in-practice clinical experiences in four Australian states and territories.

Method

Multinomial logistic regression with outcome antibiotic prescribing (no prescribing, immediate prescribing, and delayed prescribing).

Results

Of 7156 new ARTI diagnoses, no antibiotics were prescribed for 4892 (68%); antibiotics were prescribed for immediate use for 1614 diagnoses (23%) and delayed antibiotics were used for 650 diagnoses (9%). Delayed prescribing was used in 22% of otitis media, 16% of sinusitis, 13% of sore throat, 11% of acute bronchitis/bronchiolitis, and 5% of upper respiratory tract infection (URTI) diagnoses. Delayed prescribing was used for 29% of all prescriptions written. Delayed prescribing and immediate prescribing were associated with markers of clinical concern. Delayed prescribing was associated with longer duration of consultation and with fewer diagnoses/problems dealt with in the consultation.

Conclusion

Australian early-career GPs use no prescribing for ARTIs substantially more than established GPs; however, except where URIs are concerned, they still prescribe antibiotics in excess of validated benchmarks. Australian early-career GPs may use delayed prescribing more often than European established GPs, and may use it to manage diagnostic uncertainty and, possibly, conflicting influences on prescribing behaviour. The use of delayed prescribing may enable a transition to an environment of more-rational antibiotic prescribing for ARTIs.

Keywords

antibiotics; family practice; general practice; primary health care; respiratory tract infections.

INTRODUCTION

Antibiotic resistance and rational antibiotic use are national priorities in Australia, but antibiotics remain overused for non-pneumonia acute respiratory tract infections (ARTIs),^{1,2} which are common presentations to GPs.³ In 2017, 41.5% ($n = 10\,215\,109$) of the Australian population was dispensed at least one systemic antibiotic in primary care, and Australia is in the highest 25% of countries (European countries and Canada) in terms of rates of community antibiotic use.^{2,4} When managing ARTIs, antibiotics are mostly not indicated.¹

Doctors' use of antibiotics can be classified as: no prescribing, delayed prescribing, and immediate prescribing. Delayed prescribing is when the intention is to not use antibiotics unless certain criteria are met (for example, a deterioration in condition). A Cochrane review has shown that, when compared with immediate prescribing, no prescribing and delayed prescribing significantly reduce consumption of inappropriate antibiotics for ARTIs, with no difference in patient satisfaction between delayed prescribing and immediate prescribing.⁵

Some drivers of prescribing antibiotics for ARTIs are diagnostic uncertainty and concern about the consequences of potential serious illness.^{6,7} However, this concern may be misplaced because both delayed prescribing and no prescribing appear to be safe, with no statistically significant difference in hospital admissions or death when compared with immediate prescribing in a UK cohort of almost 29 000 people.⁸ Additionally, the study showed that delayed prescribing led to a significant reduction in re-consultation for new, worsening, or non-resolving symptoms, compared with both no prescribing and immediate prescribing.⁸ Although there is ample literature — including a Cochrane review⁵ — of trials of delayed prescribing, there is little evidence for the use of delayed prescribing in practice.⁹

General practice training is an important time for early-career GPs, during which they develop approaches to diagnosing and managing ARTIs, including antibiotic prescribing practices. There is evidence from a qualitative study that GPs' decision-making processes regarding antibiotic prescribing are consistent over time.¹⁰

A Davey, FRACGP, senior researcher; **A Tapley**, MMedStat, senior research officer and statistician; **KJ Mulquiney**, MCLinEpid(Distinct), research assistant; **A Fielding**, PhD, research manager, NSW & ACT Research and Evaluation Unit, GP Synergy, Regional Training Organisation, Australia. **M van Driel**, PhD, head of discipline of general practice and primary care clinical unit, Faculty of Medicine, University of Queensland, Brisbane, Australia. **E Holliday**, PhD, associate professor of biostatistics; **P Magin**, FRACGP, conjoint professor. School of Medicine and Public Health, University of Newcastle, Australia. **JS Davis**, FRACGP, senior principle research fellow, Menzies School of Health Research, Darwin, Australia. **P Glasziou**, PhD, director, Institute for Evidence-Based Healthcare, Bond University, Robina, Australia. **A Dallas**, MCLinEd, academic lead — community placements, Tasmanian School of Medicine, University of Tasmania, Hobart, Australia. **J Ball**, GradDipMedStats, statistician, Hunter Medical

Research Institute, New Lambton Heights, Australia. **N Spike**, FRACGP, director of medical education and training, Eastern Victoria General Practice Training, Regional Training Organisation, Hawthorn, Australia. **K FitzGerald**, MPH; director of education, General Practice Training Tasmania, Regional Training Organisation, Hobart, Australia.

Address for correspondence

Andrew Davey, NSW and ACT Research and Evaluation Unit, GP Synergy, Regional Training Organisation, Mayfield West 2304, NSW, Australia.

Email: andrew_davey@gpsynergy.com.au

Submitted: 11 January 2021; **Editor's response:** 7 April 2021; **final acceptance:** 21 May 2021.

©The Authors

This is the full-length article (published online 2 Nov 2021) of an abridged version published in print. Cite this version as: **Br J Gen Pract 2021**; DOI: <https://doi.org/10.3399/BJGP.2021.0026>

How this fits in

In Australia, GPs prescribe antibiotics for acute respiratory tract infections (ARTIs) well in excess of validated benchmarks. Australian early-career GPs (specialist GP vocational trainees, also known as GP registrars [13% of the GP workforce]) use no prescribing of antibiotics for ARTIs substantially more often than established Australian and European GPs, but still prescribe in excess of benchmarks, except for upper respiratory tract infections. They appear to use delayed prescribing of antibiotics to manage diagnostic uncertainty and, perhaps also, conflicting influences on antibiotic prescribing for ARTIs. Delayed prescribing may facilitate the transition to a culture of more-rational antibiotic prescribing for ARTIs in Australia.

In the study presented here, the authors aimed to:

- describe the prevalence of antibiotic prescribing strategies (immediate prescribing, delayed prescribing, and no prescribing) used by Australian early-career GPs (specialist GP vocational trainees, also known as GP registrars) for initial presentations of non-pneumonia ARTIs; and
- establish the determinants of registrars' choice of prescribing strategy.

METHOD

Participants

This study comprised a cross-sectional analysis of data from the Registrar Clinical Encounters in Training (ReCEnT) cohort study.¹¹ Data were collected from September 2016 until December 2017; the study methodology has been described in detail elsewhere.¹¹ Briefly, ReCEnT is an ongoing cohort study of GP registrars' in-practice clinical experiences undertaken in four of the eight Australian states and territories. ReCEnT is a component of GP registrars' educational programme, facilitating reflection on practice and learning.^{12,13} Registrars may consent to their data being used for research purposes.

Procedures

Characteristics of participating GP registrars and their practices are documented via a questionnaire at the beginning of each 6-monthly general practice-based training term (three compulsory terms). Registrars then record 60 consecutive office-based consultations (approximately 1 week of

consultations) at approximately the mid-point of each term.

Problems/diagnoses are coded according to the International Classification of Primary Care, second edition (ICPC-2).¹⁴ ARTI diagnoses were defined by the following codes: R74 (upper respiratory tract infection [URTI]); R78 (acute bronchitis/bronchiolitis); R75 (acute sinusitis); H71 (acute otitis media); R72 (strep throat); R76 (tonsillitis, acute); R74008 (pharyngitis, acute); R74006 (infection, throat); R74017 (pharyngitis); and R21005 (sore throat). R72, R76, R74008, R74006, and R74017 were grouped together as 'sore throat' for analysis.

Medications prescribed, provided in-consultation, or recommended are routinely recorded. Antibiotic medications were identified using the Anatomic Therapeutic Chemical (ATC) classification.¹⁵

GP registrars reported whether delayed or immediate prescribing was employed for diagnoses involving the prescription of antibiotics. No prescribing was defined as the absence of an antibiotic prescription for an ARTI diagnosis. GP registrars recorded a delayed prescription when they asked the patient not to start the antibiotic immediately.

Outcome factors

The primary outcome factor was whether a new ARTI diagnosis (that is, a diagnosis for a first presentation — re-consultations were excluded from analysis) involved delayed, immediate, or no prescribing.

Independent variables

After consideration of the potential relevance of individual variables to the research question/outcome factor (including considerations of the literature, the authors' clinical experience, and their experience with this study population and this dataset), the analysis plan, a priori, included 24 variables that were deemed relevant. These variables related to the patient, GP registrar, practice, and the consultation. Some require explanation:

- routinely bulk bills — there is no financial cost to the patient for the consultation;
- rurality — practice postcode was used to define the practice's Australian Standard Geographical Classification-Remoteness Area classification;¹⁶
- socioeconomic status of the practice location — classified according to the Socioeconomic Index for Areas Index of Relative Disadvantage¹⁷ of the practice location;
- sought clinical information or assistance

during the consultation — help was sought from the GP supervisor, a specialist, or electronic or hard-copy resources; and

- learning goals generated — clinical questions to be pursued after the consultation had finished.

Data analysis

The proportions of all ARTI diagnoses involving delayed, immediate, and no prescribing were calculated with 95% confidence intervals (CIs); they were also calculated by type of ARTI.

Multinomial logistic regression for a nominal outcome was used within the generalised estimating equations (GEE) framework to account for repeated measures within GP registrars. GEE was used for analyses, as the authors' interest was in effect estimates averaged across registrars, rather than registrar-specific effects (as produced with mixed [random-effects] models). To estimate the GEE, an exchangeable working correlation structure was assumed.

Multinomial logistic regression for a nominal outcome with three (non-ordered) levels produces two sets of statistics: one for each of the two test levels compared with the reference level. Delayed prescribing was used as the reference level. The model odds ratios (ORs) compared no prescribing with delayed prescribing, and immediate prescribing with delayed prescribing. The results are equivalent to conducting two binary logistic models.

Covariates with a univariate *P*-value of <0.20 were considered for inclusion in the multiple regression model. For each outcome, a model with all selected covariates was fitted, after which model reduction was assessed. Covariates that were no longer significant (at *P*<0.20) in the multivariable model were tested for removal from the model. If the covariate's removal did not substantively change the remaining coefficients in the model by >~10%, the covariate was removed from the final model.

In a post-hoc analysis, association of delayed prescribing, immediate prescribing, or no prescribing with seeking assistance from the registrar's supervisor was tested using a χ^2 statistic.

Analyses were programmed using Stata (version 14.0) and SAS (version 9.4).

RESULTS

A total of 631 GP registrars completed 1063 rounds of data collection (*N* = 1092, response rate 97.3%) at 454 practices contributed 101 019 patient problems/diagnoses from 63 628 consultations. The

characteristics of participating GP registrars and their practices are shown in Table 1.

There were 7156 new diagnoses of an ARTI; of these, no antibiotics were prescribed for 4892 (68% [95% CI = 67% to 69%]) diagnoses, antibiotics were prescribed for immediate use for 1614 (23% [95% CI = 22% to 24%]) diagnoses, and delayed antibiotics were used for 650 (9% [95% CI = 8% to 10%]) diagnoses (see Figure 1 and Supplementary Table S1 for details of each ARTI diagnosis). When prescribing antibiotics for ARTIs, GP registrars used delayed prescribing 29% of the time.

Characteristics associated with ARTI diagnoses by antibiotic prescribing strategy are presented in Table 2.

Univariate and multivariable associations of antibiotic prescribing strategies are shown in Table 3. For each association, the OR represents a comparison of immediate prescribing or no prescribing with delayed prescribing. When any in-consultation information or assistance was sought (12% of new ARTI diagnoses; Table 2), no prescribing was less likely (OR 0.41, 95% CI = 0.29 to 0.59) and immediate prescribing was more likely (OR 2.05, 95% CI = 1.48 to 2.84) than delayed prescribing (Table 3). When follow-up was arranged, immediate prescribing was more likely than delayed prescribing (OR 1.31, 95% CI = 1.01 to 1.70). When imaging was ordered, no prescribing was significantly more likely than delayed prescribing (OR 6.26, 95% CI = 1.42 to 27.7); there was some evidence (*P* = 0.068) for immediate prescribing being more likely than delayed prescribing (OR 3.90, 95% CI = 0.90 to 16.9).

Longer consultation duration was associated with use of delayed prescribing rather than no prescribing; for each additional minute of consultation duration, the odds of no prescribing decreased by ~2% (OR 0.98, 95% CI = 0.96 to 1.00). For each additional problem/diagnosis dealt with in the encounter, both no prescribing (OR 1.87, 95% CI = 1.55 to 2.25) and immediate prescribing (OR 1.32, 95% CI = 1.08 to 1.60) were more likely than delayed prescribing. Finally, both no prescribing (OR 1.05, 95% CI = 1.02 to 1.08) and immediate prescribing (OR 1.03, 95% CI = 1.00 to 1.06) were more likely than delayed prescribing with each additional year of age of the GP registrar.

In the post-hoc analyses, when information or assistance was sought from a supervisor (2% of new ARTI diagnoses), immediate prescribing was used significantly more than both delayed prescribing (*P* = 0.004) and no prescribing (*P* < 0.001).

Table 1. Characteristics of GP registrars and their teaching practices

Registrar variables, N= 631	n(%) ^a
Sex, Female	384 (61)
Qualified as doctor overseas	77 (12)
Years worked in hospital before entering GP training, mean (SD)	3.0 (2.6)
Registrar round/practice variables, N= 1070 ^b	
Worked at the practice in a previous term	224 (21)
Registrar age, years, mean (SD)	32.5 (6.3)
Registrar works part time	223 (22)
Registrar training term	
Term 1	414 (39)
Term 2	458 (43)
Term 3	198 (19)
Practice rurality/urbanicity	
Major city	728 (69)
Inner regional	237 (22)
Outer regional	94 (9)
Remote	0 (0)
Very remote	0 (0)
Practice location SES status (SEIFA index), mean (SD)	5.8 (2.9)
Practice routinely bulk bills	367 (36)
Full-time equivalent GPs working at the practice	
1-4	426 (41)
≥5	602 (59)

^aUnless otherwise stated. ^bNumbers may not add up to 1070 due to missing data. SD = standard deviation. SEIFA = Socioeconomic Index for Areas. SES = socioeconomic status.

DISCUSSION

Summary

Early-career GPs use no prescribing of antibiotics for 68% of ARTIs, substantially more than established Australian GPs (43%),¹ and delayed prescribing for 29% of antibiotic prescriptions for ARTIs. Except for URTIs, they prescribe antibiotics in excess of validated benchmarks.¹⁸ Prescribing antibiotics was associated with markers of diagnostic uncertainty such as seeking

in-consultation assistance and arranging follow-up.

Strengths and limitations

A high response rate, which is unusual for studies of GPs,¹⁹ along with little missing data on antibiotic prescribing (0.3%) are strengths of this study. GP registrar characteristics were reflective of the Australian GP registrar population overall,²⁰ and their practices were located in major cities, inner and outer regional areas, and remote areas. The participating Regional Training Organisations train 43% of Australian GP registrars.²⁰ That a large proportion of Australian registrars participated in the study, and their characteristics were similar to those of the wider registrar population, suggests generalisability to Australian specialist general practice vocational training.

A limitation is that contextual clinical details for the presenting conditions are not known, so it was not possible to judge how appropriate the prescription of antibiotics was for any individual problem/diagnosis; however, the finding that total prescribing is greatly in excess of benchmarks for all ARTIs other than URTI is robust because clinical judgement is implicit to the range of those benchmarks.¹⁸ A risk of misclassification bias and social desirability bias does exist, however, when a registrar does not accurately record an ARTI diagnosis because they prescribed antibiotics; the authors regard this risk as likely to be small — in ReCEnT, registrars record the broad range of their clinical activities over consecutive consultations (there is no focus on any single activity, including antibiotic prescribing).

The authors did not have data on whether patients filled the delayed prescriptions or actually took the dispensed antibiotics, nor did they have data on clinical outcomes; however, this is not considered a substantive limitation as the focus of the study was registrars' prescribing behaviour. To put the findings into some context, however, it has been found in randomised controlled trials of delayed prescribing that 31% of delayed scripts will be filled (with delayed scripts provided at consultation being more likely to be filled than those to be collected later).⁵

Comparison with existing literature

In the study presented here, GP registrars used no prescribing substantially more often than established Australian GPs in McCullough *et al's* modelling¹ for acute bronchitis/bronchiolitis (32% versus 15%), otitis media (26% versus 11%), and

Figure 1. Frequency of antibiotic prescribing strategies.




Table 2. Antibiotic prescribing for acute respiratory tract infections^a

Variable	Total ^b	No prescribing ^b	Immediate prescribing ^b	Delayed prescribing ^b	P-value
Age, years					
0–<5	1504 (21)	1119 (23)	258 (16)	127 (20)	<0.001
5–14	1067 (15)	722 (15)	234 (14)	111 (17)	
15–24	957 (13)	673 (14)	212 (13)	72 (11)	
25–44	1901 (27)	1296 (26)	437 (27)	168 (26)	
45–64	1156 (16)	756 (15)	275 (17)	125 (19)	
≥65	571 (8)	326 (7)	198 (12)	47 (7)	
Patient sex					
Male	2931 (42)	2036 (42)	648 (41)	247 (39)	0.094
Female	4086 (58)	2761 (58)	937 (59)	388 (61)	
Aboriginal and/or Torres Strait Islander					
No	6500 (98)	4446 (98)	1455 (98)	599 (99.3)	0.11
Yes	115 (2)	78 (2)	33 (2)	4 (0.7)	
NESB					
No	5974 (90)	4049 (89)	1374 (91)	551 (91)	0.079
Yes	690 (10)	505 (11)	129 (9)	56 (9)	
Registrar sex					
Male	3062 (43)	2126 (43)	681 (42)	255 (39)	0.28
Female	4094 (57)	2766 (57)	933 (58)	395 (61)	
Work hours					
Part time	1409 (20)	992 (21)	302 (19)	115 (18)	0.59
Full time	5520 (80)	3757 (79)	1249 (81)	514 (82)	
Registrar training term					
1	2749 (38)	1861 (38)	642 (40)	246 (38)	0.56
2	3227 (45)	2223 (45)	717 (44)	287 (44)	
3	1180 (16)	808 (17)	255 (16)	117 (18)	
Worked at practice previously					
No	5818 (82)	4012 (82)	1295 (81)	511 (80)	0.71
Yes	1277 (18)	853 (18)	300 (19)	124 (20)	
Qualified as doctor in Australia					
No	750 (11)	483 (10)	213 (13)	54 (8)	0.011
Yes	6390 (89)	4397 (90)	1397 (87)	596 (92)	
Practice size					
Small (1–4 FTE GPs)	2911 (42)	1976 (42)	668 (43)	267 (42)	0.90
Large (≥5 FTE GPs)	4009 (58)	2763 (58)	883 (57)	363 (58)	
Practice routinely bulk bills					
No	4235 (62)	2796 (60)	1017 (66)	422 (69)	<0.001
Yes	2609 (38)	1887 (40)	532 (34)	190 (31)	
Rurality					
Major city	5157 (73)	3597 (74)	1103 (69)	457 (71)	0.01
Inner regional	1444 (20)	914 (19)	383 (24)	147 (23)	
Outer regional remote	475 (7)	322 (7)	115 (7)	38 (6)	
Patient/practice status					
Existing patient	1530 (22)	1045 (22)	361 (23)	124 (19)	0.62
New to registrar	4783 (68)	3266 (68)	1070 (67)	447 (70)	
New to practice	712 (10)	484 (10)	157 (10)	71 (11)	
Sought help any source					
No	6284 (88)	4629 (95)	1125 (70)	530 (82)	<0.001
Yes	872 (12)	263 (5)	489 (30)	120 (18)	
Pathology ordered					
No	6713 (94)	4653 (95)	1455 (90)	605 (93)	<0.001
Yes	443 (6)	239 (5)	159 (10)	45 (7)	
Imaging ordered					
No	7050 (99)	4846 (99.1)	1557 (96)	647 (99.5)	<0.001
Yes	106 (1)	46 (0.9)	57 (4)	3 (0.5)	

... continued

Table 2 continued. Antibiotic prescribing for acute respiratory tract infections^a

Variable	Total ^b	No prescribing ^b	Immediate prescribing ^b	Delayed prescribing ^b	P-value
Referral ordered					
No	7084 (99)	4848 (99.1)	1593 (99)	643 (99)	0.86
Yes	72 (1)	44 (0.9)	21 (1)	7 (1)	
Follow-up ordered					
No	4849 (68)	3529 (72)	884 (55)	436 (67)	<0.001
Yes	2307 (32)	1363 (28)	730 (45)	214 (33)	
Learning goals generated					
No	6414 (94)	4487 (96)	1342 (87)	585 (94)	<0.001
Yes	393 (6)	165 (4)	192 (13)	36 (6)	
Respiratory disease					
Acute bronchitis	734 (10)	237 (5)	417 (26)	80 (12)	<0.001
Acute otitis media	578 (9)	148 (3)	303 (19)	127 (20)	
Acute URTI	4289 (60)	3887 (79)	179 (11)	223 (34)	
Strep/tonsillitis/pharyngitis	1002 (14)	412 (8)	458 (28)	132 (20)	
Acute sinusitis	553 (8)	208 (4)	257 (16)	88 (14)	
Registrar age, mean (SD)	32 (5)	31 (5)	32 (6)	31 (4)	0.005
SEIFA index, mean (SD)	6 (3)	6 (3)	6 (3)	6 (3)	0.25
Consultation duration, mean (SD)	15 (7)	15 (7)	16 (7)	16 (6)	<0.001
Number of problems/diagnoses, mean (SD)	2 (1)	2 (1)	1 (1)	1 (1)	<0.001

^aNumbers may not add up to 7156 due to missing data. ^bValues presented as n (%) unless otherwise stated. FTE = full-time equivalent. NESB = non-English-speaking background. SD = standard deviation. SEIFA = Socioeconomic Index for Areas. URTI = upper respiratory tract infection.

sore throat (41% versus 6%) (Figure 1, Supplementary Table S1). For URTI, GP registrars prescribed within the benchmark specified by the European Surveillance of Antimicrobial Consumption disease-specific benchmarks,¹⁸ which have been validated for use in Australian general practice,²¹ but this was not the case for the other ARTIs reported here.

When prescribing antibiotics, it appears Australian GP registrars are using delayed prescribing more often than European GPs: in a multi-country study, European GPs used delayed prescribing for 12% of prescriptions written for lower respiratory tract infections/ARTIs with cough as the dominant symptom.⁹ This percentage is most closely comparable with acute bronchitis/bronchiolitis in the study presented here — in which registrars used delayed prescribing for 16% of prescriptions written. For all cases (including where no antibiotic was prescribed), European GPs used delayed prescribing in 6.3%,⁹ compared with 11% of bronchitis/bronchiolitis in the study presented here.

Prescribing antibiotics was more likely when there were markers of clinical concern, such as seeking information or assistance during the consultation or arranging specific follow-up. This suggests GP registrars may be addressing their diagnostic uncertainty

regarding a more serious illness and the perceived consequences of not prescribing antibiotics. The authors have previously found that, when GP registrars seek help from their supervisor, they are significantly more likely to prescribe antibiotics for URTI and acute bronchitis²² — diagnoses for which authoritative Australian guidelines recommend not prescribing antibiotics.²³ Data for this study do not include clinical information to ascertain how appropriate antibiotic prescription was in other ARTIs, but it is noted that established GPs prescribe substantially more antibiotics for ARTIs in Australia than GP registrars¹ and also prescribe more than in similar medical systems in Europe and Canada.² Consequently, one possibility is that GP registrars may use immediate prescribing to be consistent with a supervisor's and/or a practice-wide approach to ARTIs; this is consistent with the authors' findings of qualitative research in this area²⁴ and may be supported by the finding of GP registrars being more likely to prescribe an antibiotic if they ask their supervisor for information or assistance.

Ordering imaging (for example, chest X-ray) can also be interpreted as a marker of clinical concern, but it is strongly associated with no prescribing. In Australia, GPs (including GP registrars) can order imaging, including

Table 3. Associations with acute respiratory tract infections^a

Variable	Outcome class	Univariate		Adjusted	
		OR (95% CI)	P-value	OR (95% CI)	P-value
Patient age group, years					
0-5	No antibiotic	1.19 (0.93 to 1.53)	0.1731	1.98 (1.39 to 2.83)	<0.001
	Immediate prescribing	0.78 (0.59 to 1.04)	0.0869	0.75 (0.51 to 1.10)	0.14
5-14	No antibiotic	0.87 (0.67 to 1.14)	0.3124	1.16 (0.82 to 1.66)	0.40
	Immediate prescribing	0.81 (0.60 to 1.08)	0.1549	0.84 (0.57 to 1.22)	0.36
15-24	No antibiotic	1.25 (0.93 to 1.68)	0.1422	1.35 (0.93 to 1.97)	0.12
	Immediate prescribing	1.17 (0.85 to 1.62)	0.3356	1.25 (0.84 to 1.86)	0.27
45-64	No antibiotic	0.78 (0.61 to 1.01)	0.0575	0.69 (0.50 to 0.95)	0.023
	Immediate prescribing	0.85 (0.64 to 1.12)	0.2515	0.81 (0.57 to 1.15)	0.23
≥65	No antibiotic	0.90 (0.63 to 1.28)	0.5656	0.75 (0.49 to 1.17)	0.21
	Immediate prescribing	1.62 (1.12 to 2.34)	0.0110	1.37 (0.86 to 2.17)	0.19
Aboriginal and/or Torres Strait Islander	No antibiotic	2.81 (1.01 to 7.80)	0.0472	1.73 (0.58 to 5.14)	0.32
	Immediate prescribing	3.11 (1.09 to 8.88)	0.0345	2.58 (0.86 to 7.69)	0.090
Practice routinely bulk bills	No antibiotic	1.54 (1.25 to 1.89)	<0.0001	1.34 (1.03 to 1.74)	0.030
	Immediate prescribing	1.17 (0.94 to 1.46)	0.1534	1.25 (0.95 to 1.64)	0.12
Sought help any source	No antibiotic	0.20 (0.16 to 0.26)	<0.0001	0.41 (0.29 to 0.59)	<0.001
	Immediate prescribing	2.13 (1.68 to 2.69)	<0.0001	2.05 (1.48 to 2.84)	<0.001
Pathology ordered	No antibiotic	0.71 (0.51 to 1.00)	0.0488	0.71 (0.46 to 1.09)	0.12
	Immediate prescribing	1.46 (1.03 to 2.08)	0.0341	1.02 (0.65 to 1.59)	0.93
Imaging ordered	No antibiotic	2.04 (0.63 to 6.63)	0.2380	6.26 (1.42 to 27.7)	0.016
	Immediate prescribing	7.63 (2.37 to 24.6)	0.0007	3.90 (0.90 to 16.9)	0.068
Follow-up ordered	No antibiotic	0.71 (0.59 to 0.86)	0.0003	0.88 (0.68 to 1.13)	0.32
	Immediate prescribing	1.76 (1.44 to 2.15)	<0.0001	1.31 (1.01 to 1.70)	0.044
Learning goals generated	No antibiotic	0.52 (0.35 to 0.77)	0.0010	1.35 (0.79 to 2.29)	0.27
	Immediate prescribing	2.39 (1.64 to 3.50)	<0.0001	1.36 (0.83 to 2.23)	0.23
Respiratory disease					
Acute otitis media	No antibiotic	0.06 (0.04 to 0.08)	<0.0001	0.05 (0.04 to 0.07)	<0.001
	Immediate prescribing	3.29 (2.45 to 4.42)	<0.0001	3.04 (2.09 to 4.43)	<0.001
Strep/tonsillitis/pharyngitis	No antibiotic	0.16 (0.13 to 0.21)	<0.0001	0.18 (0.13 to 0.24)	<0.001
	Immediate prescribing	4.85 (3.65 to 6.44)	<0.0001	4.31 (3.04 to 6.11)	<0.001
Acute sinusitis	No antibiotic	0.12 (0.09 to 0.16)	<0.0001	0.17 (0.12 to 0.25)	<0.001
	Immediate prescribing	4.05 (2.94 to 5.59)	<0.0001	3.68 (2.48 to 5.48)	<0.001
Acute bronchitis	No antibiotic	0.15 (0.11 to 0.20)	<0.0001	0.16 (0.11 to 0.24)	<0.001
	Immediate prescribing	6.97 (5.06 to 9.59)	<0.0001	5.93 (4.00 to 8.80)	<0.001
Registrar age, years	No antibiotic	1.02 (1.00 to 1.05)	0.0442	1.05 (1.02 to 1.08)	0.002
	Immediate prescribing	1.04 (1.01 to 1.06)	0.0012	1.03 (1.00 to 1.06)	0.026
Consultation duration, minutes	No antibiotic	0.99 (0.98 to 1.00)	0.0905	0.98 (0.96 to 1.00)	0.019
	Immediate prescribing	1.02 (1.00 to 1.03)	0.0294	0.99 (0.97 to 1.01)	0.19
Number of problems/diagnoses	No antibiotic	1.45 (1.28 to 1.65)	<0.0001	1.87 (1.55 to 2.25)	<0.001
	Immediate prescribing	1.21 (1.05 to 1.39)	0.0088	1.32 (1.08 to 1.60)	0.006

^aThe variables patient sex, non-English-speaking background, qualified as doctor in Australia, and rurality were not statistically significantly associated with the outcome and their removal did not alter other coefficients in the model; as such, these covariates were excluded from the multivariable model.

a radiologist report, and expect it to be performed the same day. One interpretation of this association could relate to advice to confirm a possible pneumonia diagnosis with a chest X-ray,²³ resulting in a delay in initiating antibiotics at first presentation; consequently, the GP registrar may diagnose an ARTI and investigate further before deciding whether it is pneumonia. The authors also found

some evidence for immediate prescribing also being associated with imaging; this is consistent with the GP registrar, in some cases, addressing their diagnostic uncertainty by taking a pre-emptive approach and deciding to prescribe antibiotics for a more-severe ARTI, whether or not a subsequent chest X-ray demonstrates pneumonia.²⁴

Delayed prescribing is associated with longer consultation duration; this may reflect more time being needed for complex or concerning presentations, with diagnostic uncertainty leading to the 'safety net' of delayed antibiotic prescribing. This is consistent with the authors' qualitative findings in this area,²⁴ but it may be that more time is needed to undertake an explanation of delayed prescribing rather than no prescribing.

The finding that with each additional problem/diagnosis dealt with in the consultation, delayed prescribing was increasingly less likely than both no prescribing and immediate prescribing could suggest that the GP registrar is choosing a strategy that requires less explanation and, consequently, less time in a busy consultation.

These associations with prescribing antibiotics — that is, markers of clinical concern, increased consultation duration, fewer problems/diagnoses managed per consultation — provide limited evidence that is congruent with qualitative findings. Some GP registrars use delayed prescribing, not only to address their own diagnostic uncertainty, but also to accommodate conflicting influences on prescribing for ARTIs such as: national guideline advice against use of antibiotics, desire to adhere to antimicrobial stewardship, knowledge that delayed prescribing reduces antibiotic consumption, both perceived and actual expectation from patients that antibiotics are necessary, and a supervisor approach or practice culture to prescribing for ARTIs.^{22,24}

The associations of the individual respiratory infective illness classifications used in this study, when compared with URTI, showed immediate prescribing was more likely than delayed prescribing, and delayed prescribing was more likely than no prescribing. The findings regarding otitis media, sore throat, and sinusitis may partly reflect Australian guideline recommendations²³ that antibiotics are indicated for these diagnoses in selected situations (and that delayed prescribing is an option in some circumstances). However, the associations of acute bronchitis (and the large effect sizes of these associations) cannot be reconciled with current Australian guideline recommendations that antibiotics are not indicated for acute bronchitis.²³

Implications for practice

Delayed prescribing represents a viable method for changing primary care practice to reduce the consumption of antibiotics in the community and is already being used quite frequently by early-career GPs in Australia. As GP registrars represent 13% of the Australian GP workforce by headcount,^{20,25} the findings have immediate, as well as medium- and long-term, implications for the quality of general practice care. Efforts to increase the use of both delayed prescribing and no prescribing for ARTIs are indicated, including in GP training programmes. In particular, encouraging the use of delayed prescribing may enable an acceptable transition to a future environment of more-rational antibiotic prescribing for ARTIs in primary care.

Funding

During the data collection period included in this study, funding of the Registrar Clinical Encounters in Training (ReCEnT) study was by an Australian Commonwealth Department of Health Research grant, and supported by GP Synergy, the general practice Regional Training Organisation for New South Wales and the Australian Capital Territory. GP Synergy is funded by the Australian Department of Health. The researchers gratefully acknowledge the Royal Australian College of General Practitioners Foundation for its support of this project (Family Medical Care and Research Grant 2016; reference: FMC16a 609796).

Ethical approval

The ReCEnT project has approval from the University of Newcastle Human Research Ethics Committee (reference: H-2009-0323).

Provenance

Freely submitted; externally peer reviewed.

Competing interests

The authors have declared no competing interests.

Acknowledgements

The authors would like to acknowledge the patients, GP registrars, GP supervisors, and practices that have participated in the ReCEnT project.

Open access

This article is Open Access: CC BY 4.0 licence (<http://creativecommons.org/licenses/by/4.0/>).

Discuss this article

Contribute and read comments about this article: bjgp.org.uk/letters

REFERENCES

1. McCullough AR, Pollack AJ, Plejdrup Hansen M, *et al*. Antibiotics for acute respiratory infections in general practice: comparison of prescribing rates with guideline recommendations. *Med J Aust* 2017; **207(2)**: 65–69.
2. Australian Commission on Safety and Quality in Health Care. *AURA 2019: third Australian report on antimicrobial use and resistance in human health*. 2019. <https://www.safetyandquality.gov.au/sites/default/files/2019-06/AURA-2019-Report.pdf> [accessed 21 Sep 2021].
3. Britt H, Miller GC, Henderson J, *et al*. *General practice activity in Australia 2015–16*. Sydney: Sydney University Press, 2016.
4. Van Boeckel TP, Gandra S, Ashok A, *et al*. Global antibiotic consumption 2000 to 2010: an analysis of national pharmaceutical sales data. *Lancet Infect Dis* 2014; **14(8)**: 742–750.
5. Spurling GK, Del Mar CB, Dooley L, *et al*. Delayed antibiotic prescriptions for respiratory infections. *Cochrane Database Syst Rev* 2017; **9(9)**: CD004417.
6. Tonkin-Crine S, Yardley L, Little P. Antibiotic prescribing for acute respiratory tract infections in primary care: a systematic review and meta-ethnography. *J Antimicrob Chemother* 2011; **66(10)**: 2215–2223.
7. Dallas A, van Driel M, van de Mortel T, Magin P. Antibiotic prescribing for the future: exploring the attitudes of trainees in general practice. *Br J Gen Pract* 2014; DOI: <https://doi.org/10.3399/bjgp14X681373>.
8. Little P, Stuart B, Smith S, *et al*. Antibiotic prescription strategies and adverse outcome for uncomplicated lower respiratory tract infections: prospective cough complication cohort (3C) study. *BMJ* 2017; **357**: j2148.
9. Francis NA, Gillespie D, Nuttall J, *et al*. Delayed antibiotic prescribing and associated antibiotic consumption in adults with acute cough. *Br J Gen Pract* 2012; DOI: <https://doi.org/10.3399/bjgp12X653561>.
10. Bjornsdottir I, Kristinsson KG, Holme Hansen E. Diagnosing infections: a qualitative view on prescription decisions in general practice over time. *Pharm World Sci* 2010; **32(6)**: 805–814.
11. Morgan S, Magin PJ, Henderson KM, *et al*. Study protocol: the Registrar Clinical Encounters in Training (ReCEnT) study. *BMC Fam Pract* 2012; **13**: 50.
12. Magin P, Morgan S, Henderson KM, *et al*. The Registrars' Clinical Encounters in Training (ReCEnT) project: educational and research aspects of documenting general practice trainees' clinical experience. *Australian Family Physician* 2015; **44(9)**: 681–684.
13. Morgan S, Henderson K, Tapley A, *et al*. How we use patient encounter data for reflective learning in family medicine training. *Med Teach* 2015; **37(10)**: 897–900.
14. World Health Organization. International classification of primary care, 2nd edition (ICPC-2). <https://www.who.int/standards/classifications/other-classifications/international-classification-of-primary-care> [accessed 21 Sep 2021].
15. WHO Collaborating Centre for Drug Statistics Methodology. Anatomical Therapeutic Chemical classification system: structure and principles. 2018. http://www.whocc.no/atc/structure_and_principles [accessed 21 Sep 2021].
16. Australian Bureau of Statistics. Australian Standard Geographical Classification (ASGC) — 2006, Jul 2006. 2007. <http://www.abs.gov.au/AUSSTATS/abs@.nsf/Latestproducts/1AE106C101420508CA2571A900170741> [accessed 21 Sep 2021].
17. Australian Bureau of Statistics. *An introduction to Socio-Economic Indexes for Areas (SEIFA)*. 2006. [https://www.ausstats.abs.gov.au/ausstats/subscriber.nsf/0/D729075E079F9FDECA2574170011B088/\\$File/20390_2006.pdf](https://www.ausstats.abs.gov.au/ausstats/subscriber.nsf/0/D729075E079F9FDECA2574170011B088/$File/20390_2006.pdf) [accessed 21 Sep 2021].
18. Adriaenssens N, Coenen S, Tonkin-Crine S, *et al*. European Surveillance of Antimicrobial Consumption (ESAC): disease-specific quality indicators for outpatient antibiotic prescribing. *BMJ Qual Saf* 2011; **20(9)**: 764–772.
19. Bonevski B, Magin P, Horton G, *et al*. Response rates in GP surveys — trialling two recruitment strategies. *Aust Fam Physician* 2011; **40(6)**: 427–430.
20. Taylor R, Clarke L, Edwards D. Australian General Practice Training Program: national report on the 2018 National Registrar Survey. 2018. https://research.acer.edu.au/cgi/viewcontent.cgi?article=1069&context=higher_education [accessed 21 Sep 2021].
21. Pont LG, Morgan TK, Williamson M, *et al*. Validity of prescribing indicators for assessing quality of antibiotic use in Australian general practice. *Int J Pharm Pract* 2017; **25(1)**: 66–74.
22. Dallas A, Magin P, Morgan S, *et al*. Antibiotic prescribing for respiratory infections: a cross-sectional analysis of the ReCEnT study exploring the habits of early-career doctors in primary care. *Fam Pract* 2015; **32(1)**: 49–55.
23. Therapeutic Guidelines. *Antibiotic – version 15*. West Melbourne: Therapeutic Guidelines Ltd, 2014.
24. Dallas A, Davey A, Mulquaney K, *et al*. Delayed prescribing of antibiotics for acute respiratory infections by GP registrars: a qualitative study. *Fam Pract* 2020; **37(3)**: 406–411.
25. Australian Government Department of Health. *Doctors in focus 2018*. 2018. <https://hwd.health.gov.au/resources/publications/factsheet-mdcl-2018-full.pdf> [accessed 13 Oct 2021].