

Bond University
Research Repository



Project Energise: Using participatory approaches and real time computer prompts to reduce occupational sitting and increase work time physical activity in office workers

Gilson, Nicholas D.; Ng, Norman; Pavey, Toby G.; Ryde, Gemma C.; Straker, Leon; Brown, Wendy J.

Published in:
Journal of Science and Medicine in Sport

DOI:
[10.1016/j.jsams.2016.01.009](https://doi.org/10.1016/j.jsams.2016.01.009)

Licence:
CC BY-NC-ND

[Link to output in Bond University research repository.](#)

Recommended citation(APA):
Gilson, N. D., Ng, N., Pavey, T. G., Ryde, G. C., Straker, L., & Brown, W. J. (2016). Project Energise: Using participatory approaches and real time computer prompts to reduce occupational sitting and increase work time physical activity in office workers. *Journal of Science and Medicine in Sport*, 19(11), 926-930.
<https://doi.org/10.1016/j.jsams.2016.01.009>

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

For more information, or if you believe that this document breaches copyright, please contact the Bond University research repository coordinator.

359 **Project Energise: Using participatory approaches and real time computer prompts to reduce**
360 **occupational sitting and increase work time physical activity in office workers**

361

362 Nicholas D. Gilson¹, Norman NG¹, Toby G Pavey¹, Gemma C. Ryde², Leon Straker³ and Wendy J.
363 Brown¹

364

365 Affiliations:

366 ¹ School of Human Movement and Nutrition Sciences, The University of Queensland, Brisbane, Australia.

367 ² School of Health Sciences, University of Stirling, Scotland.

368 ³ School of Physiotherapy and Exercise Science, Curtin University, Perth, Australia.

369

370 Corresponding author: Dr Nicholas Gilson, n.gilson1@uq.edu.au

371

372 Total word count: 2,479

373 Abstract word count: 250

374 Number of tables: 3

375

376

376 **Abstract**

377

378 **Objectives:** This efficacy study assessed the added impact real time computer prompts had on a
379 participatory approach to reduce occupational sedentary exposure and increase physical activity.

380 **Design:** Quasi-experimental.

381 **Methods:** 57 Australian office workers (mean [SD]; age=47 [11] years; BMI=28 [5] kg/m²; 46 men)
382 generated a menu of 20 occupational ‘sit less and move more’ strategies through participatory workshops,
383 and were then tasked with implementing strategies for five months (July-November 2014). During
384 implementation, a sub-sample of workers (n=24) used a chair sensor/software package (Sitting Pad) that
385 gave real time prompts to interrupt desk sitting. Baseline and intervention sedentary behaviour and
386 physical activity (GENEActiv accelerometer; mean work time percentages), and minutes spent sitting at
387 desks (Sitting Pad; mean total time and longest bout) were compared between non-prompt and prompt
388 workers using a two-way ANOVA.

389 **Results:** Workers spent close to three quarters of their work time sedentary, mostly sitting at desks (mean
390 [SD]; total desk sitting time=371 [71] mins/day; longest bout spent desk sitting=104 [43] mins/day).
391 Intervention effects were four times greater in workers who used real time computer prompts (8%
392 decrease in work time sedentary behaviour and increase in light intensity physical activity; p<0.01).
393 Respective mean differences between baseline and intervention total time spent sitting at desks, and the
394 longest bout spent desk sitting, were 23 and 32 mins/day lower in prompt than in non-prompt workers
395 (p<0.01).

396 **Conclusions:** In this sample of office workers, real time computer prompts facilitated the impact of a
397 participatory approach on reductions in occupational sedentary exposure, and increases in physical
398 activity.

399

400 Key words: Office workers, occupational sitting, physical activity, computer prompts

401

402 **1. Introduction**

403
404 High levels of total time spent sitting, and prolonged periods of time without interrupting sitting, have
405 been associated with increases in the prevalence of cardiovascular disease, type 2 diabetes and all-cause
406 mortality.^{1, 2} For many adults, high overall exposure and prolonged periods of sitting exposure occur at
407 work.³

408
409 In Australia for example, large-scale surveys suggest that 81% of workers are exposed to sitting,⁴ and that
410 workers employed in offices sit the most.^{5, 6} Smaller studies that have used accelerometers indicate that
411 Australian office workers spend between 4-7 hours/day sitting during work time.^{7, 8, 9}

412
413 While not as well advanced as the epidemiological and measurement based literature, evidence on the
414 effectiveness of various strategies for reducing and interrupting occupational sitting is beginning to build.
415 A recently published Cochrane review provides a comprehensive overview of studies up to February
416 2014, and highlights that there is a clear need to develop intervention strategies that target the physical
417 workplace, policy changes and information and counselling.¹⁰

418
419 Computer software that utilises screen prompts is a novel information based intervention strategy for
420 reducing and interrupting prolonged desk sitting. To date, four studies have investigated the impact
421 computer prompts have on occupational sitting, standing and moving.^{11, 12, 13, 14} Relative to a control or
422 comparative group that did not receive prompts, the findings from three of these studies showed
423 objectively measured changes in the duration of sitting bouts >30 minutes (n=14; significant decrease of
424 12%);¹¹ total sitting time at work (n=29; non-significant decrease of 7%);¹³ and standing time at work
425 (n=19; mean increase of 9%).¹⁴ One study found a significant pre-post increase in self-reported energy
426 expenditure (n=17; 188 calories) relative to a control group.¹² Intervention duration across all four studies

427 ranged from 3 days¹³ to 13 weeks,¹² and all four studies used passive prompts whereby a reminder was
428 given to break from sitting at a set time (e.g. 30 minutes) regardless of time spent sitting.

429
430 Use of passive prompts assumes that patterns of sitting in office workers are generic. Our research has
431 shown that these patterns are highly variable, suggesting that to be most effective, the receipt of prompts
432 for interrupting sitting should react to each worker's pattern of desk sitting throughout the day.⁹
433 Consequently, we have developed a low cost device called the Sitting Pad (SP), which fits easily onto
434 office chairs and has a real time, reactive feedback component that prompts employees to engage with
435 sitting reduction strategies.¹⁵

436
437 The aim of this efficacy study was to test the additional impact of real time feedback and prompts on a
438 participatory approach to reduce work time sedentary exposure and increase physical activity. In the
439 context of the present study, a participatory approach¹⁶ was used to engage two teams of Australian office
440 workers in the identification of 'sit less and move more' strategies, with one of these teams provided with
441 access to real time feedback.

442

443 **2. Methods**

444

445 The office teams were situated on different floors of a telecommunications worksite in the city centre of
446 Brisbane. Following ethics approval, the managers of the teams distributed recruitment emails to their
447 administrative and clerical workers (n=150). All members of each work team were eligible to participate,
448 although the total number of workers recruited was capped at 60 because of available resources. The
449 research targeted a relatively even number of workers within each team.

450

451 All recruited workers provided informed consent and met with researchers individually at work to
452 complete a demographic survey and basic physical measures (height, weight, waist circumference and

453 blood pressure; April 2014). Workers then wore a GENEActiv wrist accelerometer (Activinsights Ltd,
454 Cambridgeshire, UK) for one working week, 24 hours/day (30 Hz sampling rate). This device is a tri-
455 axial, ± 6 g seismic acceleration sensor, which is small (36x30x12 cm), lightweight (16 g), and
456 waterproof. GENEActiv validity studies have demonstrated strong correlations for criterion validity
457 (Pearson's $r = 0.79$ to 0.98) against indirect calorimetry for both sedentary behaviour and physical
458 activity.¹⁷

459
460 At baseline, the participants also had their office chairs fitted with the SP and this device remained on
461 chairs for the duration of the study. The SP was used as a measure of desk sitting in both teams of workers
462 and is described in detail elsewhere,¹⁵ but in brief consists of a cushion and medical grade pressure sensor
463 (43x32x2cm) which detects transitions of greater than 3 seconds to and from the seat. A microcontroller
464 linked to the pressure sensor records a time stamp for each transition and these data are downloaded using
465 a proprietary software package that summarises the total minutes spent sitting and the longest bout spent
466 sitting each day.

467
468 Following baseline measures, workers attended a one-hour workshop (n=10-15) held at the worksite (June
469 2014). During the workshop, researchers reviewed evidence on the benefits of reducing sitting and
470 increasing physical activity, and workers identified and discussed occupational strategies for 'sitting less
471 and moving more'. These strategies were collated, then thematically analysed separately by two
472 researchers, who then reached a consensus on the number of strategies, and the domains in which they
473 should be categorised. Finally, a list of strategies was distributed to workers at the start of the trial for use
474 during the five-month intervention period (July-November 2014).

475
476 The pilot study used a *posteriori*, quasi-experimental design, whereby between completion of the
477 workshops and distribution of the strategies, teams were allocated to either *Intervention Protocol 1* (IP1)
478 or *Intervention Protocol 2* (IP2). Both sets of workers were asked to implement 'sit less and move more'

479 strategies, but only workers in IP2 were provided with access to a software package that linked to the SP
480 and gave real time prompts to break from desk sitting, via a traffic light system displayed in the right hand
481 corner of computer screens.

482
483 A recently published expert statement from the United Kingdom has indicated that office workers should
484 aim to accumulate 2-4 hours/day of standing and light intensity activity time during work hours.¹⁸
485 However, the statement recognises the tentative nature of these recommendations and the need for more
486 evidence to better inform viability. In the absence of definitive health guidelines for the frequency and
487 duration of sitting breaks for adults, default settings for the SP software reflected pragmatic suggestions
488 from the literature that propose at least a five minute break from desk sitting every 30-60 minute block of
489 time.^{19,20} In this regard, prompts moved from green to amber, and then to red, after 30 and then 60
490 minutes of continuous desk sitting respectively, with the software re-setting to green after five continuous
491 minutes of the SP not being activated. User functions accessed through the software enabled workers to
492 reduce and tailor these thresholds to their own work time routine, and select an auditory as well as a visual
493 prompt.

494
495 GENEActivs were worn again for one week at the end of the intervention. Baseline and end-intervention
496 data were classified into mean percentages of waking time (work day) and work time spent in sedentary
497 behaviour, and in light and moderate-to-vigorous (moderate+) physical activity, using validated cut-point
498 thresholds. Monitors were initialised at 75Hz with cut-points of 362 (sedentary), 508 (light to moderate),
499 and 1698 (moderate+).¹⁷ Workers used the time stamp function on the wristwatch to denote when they
500 arrived and left work; this was cross-referenced with self-reported work times. SP data were isolated for
501 baseline and intervention periods, and the mean total sitting time and the longest bout spent sitting at
502 desks (minutes/day) calculated for these periods.

503

504 Minimum inclusion criteria for analyses of GENEActiv (24 hours) and SP data (≥ 6 hours) were three
505 separate workdays at baseline and (end) intervention. If intervention data were missing, baseline values
506 were carried forward, and a two-way repeated ANOVA was used to statistically compare between and
507 within group changes in sedentary behaviour, light and moderate+ physical activity and desk sitting. Main
508 outcome analyses focused on changes in these variables during work time; all statistical analyses were
509 conducted in SPSS version 22 (IBM Corporation); p-values were based on two-sided tests and were
510 considered statistically significant at $p < 0.05$.

511

512 **3. Results**

513

514 Table 1 summarises the demographic characteristics of the sample. Participants tended to be mainly men,
515 overweight and middle aged. At baseline, IP1 workers spent significantly more time in light intensity
516 physical activity on workdays than IP2 workers ($p = 0.024$), but no significant differences were observed
517 for GENEActiv and SP baseline data collected during work time.

518

519 Insert Table 1 here.

520

521 Analyses of data collected from workers during workshops identified a menu of 20 ‘sit less and move
522 more’ strategies. As Table 2 shows, these were themed into the four occupational contexts of ‘desk tasks’
523 (5 strategies; e.g. deliver some messages in person rather than always sending emails); ‘meetings’ (4
524 strategies; e.g. use walk-talk rather than sit-talk meetings); work breaks (7 strategies; e.g. take a standing
525 and stretching break); and travel (4 strategies; e.g. walk to and/or from work).

526

527 Insert Table 2 here.

528

529 Table 3 summarises work time GENEActiv and SP data, and describes changes relative to baseline and
530 (end) intervention values. With regard to the carry forward of GeneActiv and SP baseline data, seven and
531 three workers respectively did not meet minimal inclusion criteria for intervention data, while a further
532 four workers withdrew from the study prior to completion of final measures. Recorded GeneActiv and SP
533 work time for these phases ranged from 8.6 – 8.9 hours/day. The mean (SD) number of monitored days for
534 the GENEActiv was 4 (1) for both groups at baseline and end-intervention. For IP1, mean (SD) monitored
535 days for the SP at baseline and intervention were 10 (5) ranging from 4-17 days, and 15 (5) ranging from
536 3-24 days respectively. Equivalent SP baseline and intervention data for IP2 were 13 (5) ranging from 3-
537 24 days, and 20 (8) ranging from 5-32 days.

538

539 Insert Table 3 here.

540

541 At baseline, workers spent 68-74% of their work time sedentary, and 19-25% in light and 7% in
542 moderate+ physical activity. On average workers spent 370-372 mins/day sitting at a desk, with the
543 longest bouts of desk sitting averaging 100-111 minutes/day.

544

545 Relative to baseline, end-intervention GeneActiv data indicated that IP1 workers reduced sedentary work
546 time by an average of 2% (18 mins/day). Within group effects were significant and four times greater in
547 IP2 workers (decrease of 8%; 72 mins/day; $p=0.012$). IP2 workers replaced sedentary work time with light
548 intensity physical activity, with the proportional increase in light intensity work time significant for this
549 group ($p=0.018$).

550

551 For the SP, comparisons of baseline and intervention data indicated that the mean total time spent sitting
552 at desks across the intervention period increased for IP1 workers (10 mins/day) and decreased for IP2
553 workers (-13 mins/day; difference of 23 mins/day). This pattern was also observed for the longest bout

554 spent sitting at desks (IP1 = 17 mins/day; IP2 = -15 mins/day; significant difference of 32 mins/day;
555 p=0.018).

556

557 **4. Discussion**

558

559 In this efficacy study, Australian office workers attended workshops and were asked to identify ‘sit less
560 and move more’ workplace strategies through a participatory approach. In the ergonomics field,
561 participatory approaches have been used to successfully engage workers, managers and employers in the
562 development of strategies to address musculoskeletal issues.^{21, 22} More recently, studies that have focused
563 on sitting and chronic disease in office workers have advocated²³ and used¹⁶ participatory approaches as a
564 means of promoting ownership of occupational sitting reduction strategies and facilitating commitment to
565 sedentary behaviour change.

566

567 The Australian office workers involved in the participatory research undertaken by Parry et al¹⁶ identified
568 seven strategies within the overarching themes of ‘active office work, traditional physical activity and
569 office ergonomics’. In a more recent study, which used a prescriptive, top-down approach with Spanish
570 office workers, Bort-Roig et al²⁴ also described seven strategies focusing on reducing and interrupting
571 occupational sitting through incidental movement. A main finding of our study is that participants
572 identified a more comprehensive ‘menu’ of 20 strategies, themed into four specific occupational contexts.

573 This ‘menu’ is valuable for employers and practitioners interested in providing office workers with a
574 range of choices and opportunities for reducing and interrupting sitting in different situations, and with
575 occupational groups who have different job demands and daily routines. Ongoing testing now needs to
576 occur to assess if the ‘menu’ is exhaustive and replicable with other office-based samples.

577

578 Linked to the emergence of strategies from the workshops, a key aim of the present study was to assess
579 the added effect of using real time computer prompts to encourage positive changes in occupational

580 sedentary exposure and physical activity, over and above those seen when using a participatory approach
581 alone. The more meaningful reductions in both the proportion of sedentary work time, and patterns of
582 desk sitting found in IP2 compared to IP1 workers, together with the significant increase in work time
583 light intensity physical activity observed in the strategy and real time prompts group, highlighted the
584 efficacy of using this combined approach. The significant reduction in the mean longest bout of desk
585 sitting in IP2 workers is particularly noteworthy, given that most occupational sitting time occurs at desks,
586 ⁹ and frequent interruptions to sitting have been linked to a range of improved health indices. ²⁵
587
588 Technology based, real time feedback, provided through pedometers and smartphone applications for
589 example, has been recognised as playing a key role in facilitating engagement in physical activity. ²⁶ A
590 main finding of the present study indicates that this type of feedback may also be an important approach
591 for encouraging workers to engage in occupational sitting reduction strategies. Providing real time
592 prompts to interrupt occupational sedentary exposure allowed workers in IP2 to self-monitor and regulate
593 their desk sitting through a personalised feedback mechanism that reacted to variations in daily desk
594 sitting patterns. In this regard, the SP and its associated software alerted workers to prolonged desk sitting
595 as and when it occurred. Evidence concerning the impact individual level strategies such as education,
596 counseling and prompts have on reductions in occupational sitting time are inconsistent. ¹⁰ Our data adds
597 to the view that a participatory approach allied with real time prompts may be valuable. However, it is
598 important to consider that multi-level interventions, that target the individual, environmental,
599 organisational and policy domains of the office work system are more likely to have a comprehensive and
600 sustainable impact on occupational sitting, than interventions that target any one domain in isolation. ^{27, 28}
601
602 The study had a number of strengths. For example, the SP provided a highly accurate assessment of desk
603 sitting over multiple days, ¹⁵ and in conjunction with the accelerometers, enabled a comprehensive
604 assessment of work time sedentary and physical activity patterns. Other study strengths included engaging
605 workers in identifying strategies through participatory approaches, and the use of a quasi-experimental

606 design that allowed comparisons of work time changes during a five-month intervention period.

607
608 The study also had some limitations that future studies need to address. The research was undertaken at a
609 single worksite, with relatively small sample sizes in each group, which while in keeping with the remit of
610 efficacy research, presently restricts the applicability of the findings to office workers in general.
611 Furthermore, the lower and uneven number of participants in IP2 (N=24) relative to IP1 (N=33) impacted
612 the detection of statistically significant between group differences; for example based on an observed
613 group mean difference of 23 minutes/day for total time spent sitting at desks, analyses required at least 31
614 workers per group ($\alpha=0.05$; $\beta=0.80$; ± 46 minutes). Lastly, the intervention effects observed in our study
615 compare favourably to those reported in studies that have used passive computer prompts and objective
616 measures of sedentary behavior change,^{11, 13, 14} although comparisons are problematic, given the
617 differences in study design and measurement techniques. The next study in this area needs to include both
618 a passive and reactive prompts group, to concurrently appraise the extent to which each approach impacts
619 on strategy uptake and occupational sedentary exposure. Future study designs may also consider mapping
620 strategy use in comparative groups to quantify the extent to which prompts encourage the uptake of
621 different types of strategies.

622

623 **5. Conclusions**

624

625 This efficacy study is the first to evaluate the impact of real time computer prompts on work time
626 sedentary behavior, physical activity and desk sitting in office workers. The main study findings provide a
627 comprehensive ‘menu’ of ‘sit less and move more’ strategies for application and ongoing investigation,
628 and begin to provide insights into the potential benefits of using real time individualised computer
629 prompts as feedback to facilitate reductions in occupational sedentary exposure and increases in work time
630 physical activity.

631

632 **Practical Implications**

- 633
- 634 • Office workers can choose from a ‘menu’ of ‘sit less and move more’ strategies.

 - 635 • Real time computer prompts, which react to day-to-day variations in desk sitting patterns, may
636 facilitate the uptake of ‘sit less and move more’ strategies.

 - 637 • The next phase of studies in this area should concurrently compare the efficacy of passive and reactive
638 computer prompts.

639

640 **Acknowledgements**

641

642 This study was supported by an Australian Heart Foundation Vanguard Grant (no.X).

643

644 Authors X and X are supported by an Australian National Health and Medical Research Council Program
645 Grant (NHMRC no. X).

646

647 Author X is supported by an Australian National Health and Medical Research Council Senior Research
648 Fellowship (NHMRC no. X).

649

650

651

652

653

654

655

656

657 **References**

658

- 659 1. Chau JY, Grunseit AC, Chey T et al. Daily sitting time and all-cause mortality – a meta-
660 analysis. *PLoS One* 2013; 8: e80000. doi:10.1371/journal.pone.0080000.
- 661 2. Demsey PC, Owen N, Biddle SJH et al. Managing sedentary behaviour to reduce the risk
662 of diabetes and cardiovascular disease. *Curr Diab Rep* 2014; 14: 522 DOI
663 10.1007/s11892-014-0522-0.
- 664 3. Kazi A, Duncan M, Clemes S et al. A survey of sitting time among young adults. *Occup
665 Med* 2014; 64: 497-502.
- 666 4. Safe Work Australia. *National hazard exposure worker surveillance: Exposure to
667 biomechanical demands, pain and fatigue symptoms and the provision of controls in
668 Australian workplaces*. Safe Work Australia, 2011.
- 669 5. Australian Bureau of Statistics. *Australian Health Survey: Physical Activity*. Australian
670 Bureau of Statistics, 2013.
- 671 6. de Cocker K, Duncan MJ, Short C et al. Understanding occupational sitting: prevalence,
672 correlates and moderating effects in Australian employees. *Prev Med* 2014; 67: 288-294.
- 673 7. Brown HE, Ryde GC, Gilson ND et al. Objectively measured sedentary behavior and
674 physical activity in office employees: relationships with presenteeism. *J Occup Environ
675 Med* 2013; 55: 945-953.
- 676 8. Parry S, Straker L. The contribution of office work to sedentary behaviour associated risk.

- 677 *BMC Public Health*, 2013; 13, 296.
- 678 9. Ryde GC, Brown HE, Gilson ND et al. Are we chained to our desks? Describing desk
679 based sitting using a novel measure of occupational sitting. *J Phys Act Health* 2014; 11:
680 1318-1323.
- 681 10. Shrestha N, Ijaz S, Kukkonen-Harjula KT et al. Workplace interventions for reducing
682 sitting at work. *Cochrane Database Syst Rev* 2015; 1: CD010912.
- 683 11. Evans RE, Fawole HO, Sheffiff SA et al. Point-of-choice prompts to reduce sitting time at
684 work: a randomised trial. *Am J Prev Med* 2012; 43: 293-297.
- 685 12. Pedersen SJ, Cooley PD, Mainsbridge C. An e-health intervention designed to increase
686 workday energy expenditure by reducing prolonged occupational sitting habits. *Work*
687 2014; 49: 289-295.
- 688 13. Swartz AM, Rote AE, Welch WA et al. Prompts to disrupt sitting time and increase
689 physical activity at work. *Prev Chronic Dis* 2014; 1: E73.
- 690 14. Donath L, Faude O, Schefer Y et al. Repetitive daily point of choice prompts and
691 occupational sit-stand transfers, concentration and neuromuscular performance in office
692 workers: an RCT. *Int J Environ Res Public Health* 2015; 20: 4340-4353.
- 693 15. Ryde GC, Gilson ND, Suppini A et al. Validation of a novel, objective measure of
694 occupational sitting. *J Occup Health* 2012; 54: 383-386.
- 695 16. Parry S, Straker L, Gilson ND et al. Participatory workplace interventions can reduce
696 sedentary time for office workers – a randomized control trial. *PlosOne* 2013; 8: e78957.

- 697 17. Esliger DW, Rowlands AV, Hurst TL et al. Validation of the GENEActiv Accelerometer.
698 *Med Sci Sports Exerc* 2011; 43: 1085-1093.
- 699 18. Buckley JP, Hedge A, Yates T et al. The sedentary office: a growing case for change
700 towards better health and productivity. Expert statement commissioned by Public Health
701 England and the Active Working Community Interest Company. *Br J Sports Med* 2015;
702 doi: 10.1136/bjsports-2015-094618.
- 703 19. Owen N, Bauman A, Brown WJ. Too much sitting: a novel and important predictor of
704 chronic disease risk? *Br J Sports Med* 2009; 43: 81-3.
- 705 20. Straker L. Prevention needs to be a priority. *Journal of Physiother* 2012; 58: 5-7.
- 706 21. Rivlis I, Van Eerd D, Cullen K, et al. Effectiveness of participatory ergonomic
707 interventions on health outcomes: A systematic review. *Appl Ergon* 2008; 39: 342–358.
- 708 22. van Eerd D, Cole D, Irvin E et al. Process and implementation of participatory ergonomic
709 interventions: a systematic review. *Ergonomics* 2010; 53: 1153–1166.
- 710 23. Gilson ND, Straker L, Parry S. Occupational sitting: Practitioner perceptions of health
711 risks, intervention strategies and influences. *Health Prom J Austr* 2012; 23: 208-212.
- 712 24. Bort-Roig J, Martin M, Puig-Ribera A et al. Uptake and factors that influence the use of
713 'sit less, move more' occupational intervention strategies in Spanish office employees. *Int*
714 *J Behav Nutr Phys Act* 2014; 10: 152.
- 715 25. Healy GN, Dunstan DW, Salmon J et al. Breaks in sedentary time: Beneficial associations
716 with metabolic risk. *Diabetes Care* 2008; 31: 661-666.

- 717 26. Bort-Roig J, Gilson ND, Puig-Ribera A et al. Measuring and Influencing Physical
718 Activity with Smartphone Technology: A Systematic Review. *Sports Med* 2014; 44: 671-
719 686.
- 720 27. Healy GN, Eakin EG, Lamontagne AD et al. Reducing sitting time in office workers:
721 short-term efficacy of a multicomponent intervention. *Prev Med* 2013; 57: 43-48.
- 722 28. Neuhaus M, Healy GN, Dunstan DW et al. Workplace sitting and height-adjustable
723 workstations: a randomized controlled trial. *Am J Prev Med* 2014; 46: 30-40.
724

724 **Tables**

725

726 Table 1 Baseline sample characteristics (frequency or mean [SD]) relative to intervention protocol

	IP1	IP2	Total
Age (years)	47 (13)	48 (13)	47 (11)
Sex (n)			
Men	27	19	46
Women	6	5	11
BMI (kg/m ²)	28 (4)	29 (5)	28 (5)
Waist Circumference (cm)			
Men	95 (13)	100 (15)	98 (14)
Women	86 (15)	89 (14)	88 (13)
Blood Pressure (mm Hg)			
Systolic	127 (14)	130 (13)	128 (13)
Diastolic	86 (9)	87 (10)	86 (9)
GENEActiv (hrs/day) ¹			
Workdays			
Sedentary	9.4 (2.2)	10.1 (1.5)	9.7 (2.0)
Light ²	5.5 (1.7)	4.3 (1.4)	5.0 (1.7)
Moderate+	1.7 (0.8)	1.6 (0.6)	1.6 (0.7)

727

728 IP1=Intervention Protocol 1 (strategies and no prompts, N=33); IP2=Intervention Protocol 2 (strategies

729 and prompts, N=24).

730 ¹ Total time for the waking day; ² p<0.05 IP1 vs IP2.

731

732

733

734

735

736

737

738 Table 2 Sit less and move more strategies (x20) identified by workers

739

Theme 1: Standing and moving during desk tasks

Stand up and move while talking on the phone.

Stand up to read or review documents.

Use centralised office equipment (e.g. photocopier or printer) and spread these tasks across the day.

Deliver some messages in person, rather than always sending emails.

Drink more water at your desk and take frequent toilet breaks.

Theme 2: Standing and moving in or between meetings

Try to organise walk-talk, rather than sit-talk meetings.

Spend some of your time standing in meetings.

Schedule stand up meetings.

Schedule meetings away from your desk or use a meeting venue outside the office.

Theme 3: Standing and moving during work breaks

Schedule regular activity breaks in your weekly work calendar.

Use morning and afternoon work breaks to undertake a stand and stretch routine.

Don't eat lunch at your work desk.

Tea or coffee break away from the desk.

Plan lunchtime walks either on your own or with colleagues.

Include inclines in your walks or use the stairs between floors.

Take advantage of corporate gym memberships and work out at a facility close to the office.

Theme 4: Active travel to and from work

Aim to stand rather than sit on public transport.

Include a walk in your travel to and from work.

Park the car further away from the office, or use a public transport stop further away from home or the office to increase your walk time.

Cycle into work.

740

741

742

743

744

745

746 Table 3 Mean (SD) baseline, (end) intervention and changes ([end] intervention – baseline) in work time
 747 sedentary behaviour, physical activity and desk sitting relative to intervention protocol
 748

	Baseline		(End) Intervention		Change	
	IP1	IP2	IP1	IP2	IP1	IP2
GENEActiv (%)						
Sedentary ¹	68 (14)	74 (7)	66 (16)	66 (15)	-2	-8
Light ²	25 (11)	19 (7)	26 (12)	27 (14)	+1	+8
Moderate+	7 (4)	7 (3)	8 (5)	7 (3)	+1	0
Sitting Pad (mins/day)						
Total time sitting	370 (84)	372 (53)	380 (81)	359 (71)	+10	-13
Longest bout sitting ³	100 (42)	111 (45)	117 (49)	96 (30)	+17	-15

749
 750 IP1= Intervention Protocol 1 (strategies and no prompts, N=33); IP2=(Intervention Protocol 2 (strategies
 751 and prompts, N=24)

752 ¹ p<0.01 IP2 baseline vs end intervention; ² p<0.01 IP2 baseline vs end intervention; ³ p<0.05 IP1 vs IP2

753

754

755

756

757 29.