

Bond University  
Research Repository



## The use of a daily diary system to promote self-monitoring and improve health-related identity and self-efficacy

Urzi, Dino; Stapleton, Peta Berenice; Chatwin, Hannah

*Published in:*  
Graduate Journal of Sport, Exercise & Physical Education Research

*Licence:*  
CC BY-NC-ND

[Link to output in Bond University research repository.](#)

*Recommended citation(APA):*  
Urzi, D., Stapleton, P. B., & Chatwin, H. (2016). The use of a daily diary system to promote self-monitoring and improve health-related identity and self-efficacy. *Graduate Journal of Sport, Exercise & Physical Education Research*, 4, 14-28. <http://www.worc.ac.uk/gjseper/656.htm>

### General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

For more information, or if you believe that this document breaches copyright, please contact the Bond University research repository coordinator.

## **THE USE OF A DAILY DIARY SYSTEM TO PROMOTE SELF-MONITORING AND IMPROVE HEALTH-RELATED IDENTITY AND SELF-EFFICACY**

**Dino Urzi, Peta Stapleton  & Hannah Chatwin**

School of Psychology, Bond University, Gold Coast, Queensland, 4229, Australia

### **ABSTRACT**

Diet and exercise have been shown to have significant importance in predicting both physical and psychological well-being. However, there is little empirical evidence examining the effects of regular practices which encourage awareness on an individual's health-related identity and self-efficacy. This study examined the effects of a daily awareness-based intervention on health-related identity and self-efficacy. Fifty-four adults were allocated into either an 8-week control (n=25) or intervention group (n=29) and all completed a battery of measures designed to assess their health-related identity and self-efficacy. The intervention group were sent a daily email reminder for 8 weeks to log on to a computer website and answer a series of reflective questions. These questions related to the quality and volume of food and liquids consumed throughout the day, their mood, the type, intensity and duration of exercise which they engaged in throughout the day, and the amount of paid or unpaid work hours which they were engaged in throughout the day. Finally, participants were asked daily whether they aimed to achieve balance in any of their regular roles. The control group did not engage in any intervention during the 8-week phase. The 'daily diary' website and questions aimed to promote awareness of health behaviours, and results indicated the intervention group displayed a significant improvement in all variables, when compared with the control group.

**Keywords:** identity, self-efficacy, awareness, health

**Article citation:** Urzi, D., Stapleton, P., & Chatwin, H. (2016) The use of a daily diary system to promote self-monitoring and improve health-related identity and self-efficacy. *Graduate Journal of Sport and Exercise & Physical Education Research*, 4: 14-28.

### **INTRODUCTION**

The physical and psychological importance of exercise and diet has been well documented in research (Briefel & Johnson, 2004). More recently, research has focused on investigating the benefits of exercise and diet for mental health and well-being. According to the Australian National Health Survey in 2011 and 2012 (Australian Bureau of Statistics [ABS], 2012), 54% of Australians aged 30 to 65 years of age reported that they did not engage in enough physical activity as per national guidelines, with sedentary behaviour increasing with age. Participants reported that the main barriers to engagement in physical activity included perceived lack of time (44%) and a lack of enjoyment of physical exercise (39%; ABS, 2012). However, it is

clear that regular engagement in physical activity can prevent illnesses such as cardiovascular disease, cancer, diabetes and several mental health difficulties (Bauman et al., 2003).

Current research suggests one in every five Australians will also experience a mental illness in their lifetime, and mental illnesses are the third leading cause of disability burden. This accounts for an estimated 27% of the total years lost due to disability (Pirkis, Blood, Dare, & Holland, 2008). Men and women in Australia experience similar rates of mental illness, but rates are highest for men and women living alone, separated or divorced (ABS, 2012). We know mental illnesses can have many physical features, as well as psychological, and have profound physical effects such as insomnia, weight gain or loss, increase or loss of energy, chest pain and nausea (SANE Australia, 2012).

### **Diet**

It is well known that worldwide, life expectancy is increasing. In order to reach a longer and healthier life, we need to focus on lifestyle choices, importantly, diet and physical activity (Kennedy, 2006). Studies indicate the global obesity epidemic has been accelerating for four decades (Gortmaker et al., 2011). Recent research (WHO, 2014) reports that obesity now represents the fifth leading cause of global deaths, with at least 2.8 million adults dying each year as a result of being overweight or obese, largely due to unhealthy eating behaviours. Overeating or the excessive consumption of foods which are nutritionally poor and energy-dense has been shown to cause physical illness such as diabetes, asthma, irritable bowel syndrome, cardiovascular problems, pain disorders and sexual functioning disorders (Hunter, Goodie, Oordt, & Dobmeyer, 2009). However, little is known about the influence of people's diet on their psychological well-being. What we do know is that several mental health issues are exacerbated by, and can be rooted in, suboptimal nutritional intake (Leyse-Wallace, 2013).

Anxiety, depression, eating and substance use disorders in particular, have been linked to a depleted nutritional status (Bodnar & Wisner, 2005). More specifically, Leyse-Wallace (2013) explained that irregularities in nutritional intake (i.e., starvation and food restriction) cause chemical and neurotransmitter dysfunction, which may in turn contribute to changes in mood and general cognitive function. Overeating, or excessive consumption of nutritionally poor and energy-dense foods, is also associated with various physical ailments including diabetes, asthma, irritable bowel syndrome and cardiovascular problems (Hunter, Goodie, Oordt, & Dobmeyer, 2009). Additionally, poor dietary quality has been linked to the onset of psychiatric conditions such as depression, by means of biological (i.e., lack of nutritionally valuable vitamins) and psychological factors (i.e., body dissatisfaction, guilt, shame and decreased self-efficacy) (Ackard, Neumark-Sztainer, Story, & Perry, 2003; Bodnar & Wisner, 2005).

### **Exercise**

In terms of adequate intensity for health benefits, evidence has found that engaging in a moderate intensity exercise activity every day of the week has been linked with an increase in physical health (Bailis, Segall, & Chipperfield, 2003; Mustard, Vermeulen, & Lavis, 2003), an increase in physical functioning (Leino-Arjas, Solovieva, Riihimarki, Kirjonen, & Telama, 2004) and improved psychological well-being (Hassmen, Koivula,

& Uutela, 2000). Engagement in physical activity or exercise has been shown to be an effective primary and secondary treatment protocol for individuals with both physical and mental health concerns (Bauman et al., 2003; Sharma, Madaan, & Petty, 2006).

### **Understanding Health Choices through Theory**

An individual's health behaviour choices are often contextualised through identity theory, which can be summarised as the process whereby an individual endorses a specific identity, and aims to engage in behaviours that will enable them to maintain an identity-behaviour congruence (Martin, Leary, & O'Brien, 2001). Individuals with higher healthy eating identities score lower on affect measures, and greater intentions to regulate future behaviour towards identity-behaviour consistency (Strachan & Brawley, 2008).

Social cognitive theory (Bandura, 1997) also posits the importance of a personal sense of control and self-efficacy in an individual's pursuit of health-related behaviours and outcomes. Self-efficacy has been shown to impact the outcomes of various health behaviours, including the effort invested into the health behaviour and perseverance in the face of challenges (Schwarzer & Luszczynska, 2005). Adults with a greater sense of dietary self-efficacy report greater consumption of fruit and vegetables, lower consumption of energy-dense foods and generally healthier food choices, despite the presence of potential barriers (Bere & Klepp, 2004; Lubans et al., 2012; levers-Landis et al., 2003), when compared with individuals with a lower sense of dietary self-efficacy.

Results of Anderson and Cychosz's early research (1995) investigating the relationship between exercise identity and exercise commitment showed a positive correlation between time spent exercising and a participant's exercise identity. Authors further concluded that a reciprocal relationship may exist between exercise identity and exercise behaviour, which may be reinforced through validation and recognition of significant others. A positive correlation exists between physiological indicators of fitness (i.e., oxygen consumption levels, body fat percentages and cholesterol levels) and exercise identity, and a strong, positive correlation between exercise behaviour and exercise identity (Anderson, Cychosz, & Franke, 1998). Engaging in exercise may promote an exercise identity, and the physiological responses to exercise may be an important factor in determining exercise identity.

### **Self-awareness and Self-monitoring with Health Identity**

The act of learning to self-monitor and pay attention to one's own behaviour in the present moment can lead to greater self-awareness and insight, which may otherwise be unachievable in an environment rife with distraction and overstimulation (Bishop et al., 2004). Self-regulation theories, such as those mentioned, discuss that awareness and attention are vital in the maintenance and enhancement of psychological and behavioural functioning. Well-being may therefore be enhanced, through self-regulated activity whereby an individual chooses to fulfil a psychological need, making them more likely to keep engaging in that behaviour (e.g., exercise).

Self-monitoring has typically been taught as a skill in the weight loss and maintenance field. Monitoring – any food eaten, all foods eaten, time food was eaten, quantity of food eaten and grams of fat consumed – appears positively correlated with weight change (the recommendation to monitor on at least 75% of days). In addition, not monitoring at all appears negatively associated with weight change (Baker &

Kirschenbaum, 1993; Boutelle & Kirschenbaum, 1998). There is some suggestion too, that beyond dietary self-monitoring, the self-monitoring of exercise also contributes to successful weight loss. Greater self-monitoring of exercise has been significantly associated with fewer difficulties with exercise, higher levels of weekly exercise and greater weight loss (Carels et al., 2005).

Health identity and self-efficacy has continually been linked to the practice of being self-aware and learning to self-monitor (Saksvig et al., 2005). However, there is a lack of published empirical research investigating the potential reasons why some participants who hold healthy identities do not engage in health-related behaviours – is there a link between self-monitoring and self-awareness? The present study sought to examine this and assess whether engaging in a daily intervention based on increasing self-monitoring and self-awareness, would improve exercise adherence, and dietary intake and health-related identity and self-efficacy.

### **The Current Study**

The present study utilised a daily act of recording eating- and exercise-related information online via a daily reminder message service (for 8 weeks) to increase self-monitoring and self-awareness in participants, and hypothesised this would improve health-related identity and self-efficacy. One group would receive the daily reminder message to record their behaviour, and another group would not.

It was hypothesised that participants in the intervention group would report significantly greater exercise identity, healthy eating identity, exercise self-efficacy, and healthy eating self-efficacy at post-intervention, when compared with participants in the control group. It was anticipated that the daily prompting message service would enhance self-awareness and attention.

## **METHOD**

### **Participants**

A total of 54 participants were allocated into the intervention ( $n = 29$ ) and control groups ( $n = 25$ ). The age of participants in both the control and intervention groups ranged from 18 to 65 years and the total sample consisted of 16 males and 38 females. The treatment group consisted of 9 males and 20 females, and the control group 7 males and 18 females. The mean age of the intervention group was 32.2 years ( $SD = 1.35$ ), and the control group 35.7 years ( $SD = 1.07$ ). The upper limit of 65 was chosen because several questions for the intervention group related to employment and 65 is the current age of retirement. All participants within the intervention and control groups were required to complete an online questionnaire package, which served as baseline data. Each set of questionnaires included a unique participant code that was used to match individuals' responses across all measurement points, whilst maintaining confidentiality. The questionnaire package involved several outcome measures, including: the Exercise Identity Questionnaire (Anderson & Cychosz, 1994), the Healthy Eating Habits Questionnaire (Dillman, 2012), the Exercise Self-efficacy Scale (Schwarzer & Renner, 2000) and the Nutrition Self-efficacy Scale (Schwarzer & Renner, 2000). Intervention participants engaged in the 8-week programme (detailed below) and then both groups again completed the online questionnaire package, after this 8-week period.

## Design

The independent variable for this study was group, consisting of two levels including the intervention and control groups. The dependent variables (DVs) were: exercise identity (Exercise ID), healthy eating identity (Eating ID), exercise self-efficacy (Exercise SE) and healthy eating self-efficacy (Eating SE). The data was analysed through a series of repeated measures ANOVA. Prior to data analysis, the data was cleaned via standardised procedures for analysing missing data, patterned responses and unvarying data. There were no significant differences between the intervention and control group at pre-intervention on any of the DVs.

## Sample Characteristics

A series of chi-square analyses were performed to determine whether the groups differed based on demographic variables. Results revealed no significant differences between groups in relation to gender  $\chi^2(1) = 0.59$ ,  $p = 0.808$ ; age  $\chi^2(4) = 4.633$ ,  $p = 0.327$ ; Exercise ID  $\chi^2(33) = 34.56$ ,  $p = 0.393$ ; Exercise SE  $\chi^2(12) = 14.221$ ,  $p = 0.287$ ; Eating ID  $\chi^2(25) = 28.795$ ,  $p = 0.273$ ; Eating SE  $\chi^2(10) = 15.312$ ,  $p = 0.121$ . In summary, the two groups did not differ significantly at baseline.

## Measures

*Healthy-eater identity.* The Healthy Eater Identity Scale (Strachan & Brawley, 2009) was used to assess participants' identity with respect to several healthy eating behaviours. A sample item includes "When I describe myself to other people, I usually include my involvement in healthy eating". Participants were required to rate the accuracy of statements on a 7-point Likert-type scale ranging from 1 (strongly disagree) to 7 (strongly agree). This scale has demonstrated high internal consistency ( $\alpha = 0.89$ ; Dillman, 2012) and a strong criterion validity with existing indices of healthy eating behaviour (Cardinal, 1997). In the current study, this scale revealed high internal consistency ( $\alpha = 0.87$ ).

*Exercise identity.* The Exercise Identity Scale (Anderson & Cychosz, 1994) was used to assess whether participants possess identities consistent with that of a frequent exerciser. A sample item includes "Physical exercise is a central factor to my self-concept". Participants rated items on a 7-point Likert-type scale ranging from 1 (strongly disagree) to 7 (strongly agree). In previous studies, this scale has demonstrated sound internal consistency in the empirical literature, with a Cronbach's alpha ranging from 0.61 to 0.89 (Dillman, 2012; Strachan & Brawley, 2008). In the current study, this scale demonstrated high internal consistency ( $\alpha = 0.93$ ).

*Healthy eating self-efficacy.* The Nutrition Self-efficacy Scale (Schwarzer & Renner, 2000) was used to assess participants' beliefs regarding their ability to eat healthily. A sample item includes "I can manage to stick to healthy foods even if I do not receive a great deal of support from others when making my first attempts". Participants were required to rate the accuracy of statements on a 4-point scale ranging from 1 (definitely not true) and 4 (exactly true). This scale has been shown to possess sound internal consistency ( $\alpha = 0.77$ ; Chang, Nitzke, & Brown, 2003), and strong construct validity (Kroll, Kehn, Ho, & Groah, 2007). In the current study, this scale revealed high internal consistency ( $\alpha = 0.87$ ).

*Exercise self-efficacy.* The Exercise Self-efficacy Scale (Schwarzer & Renner, 2000) was used to assess participants' beliefs regarding their ability to initiate and pursue

active engagement in physical activity. A sample item includes “I can manage to carry out my exercise intentions even when I have worries and problems”. Participants were required to rate the accuracy of statements on a 4-point scale ranging from 1 (definitely not true) and 4 (exactly true). Previous studies have evidenced the internal consistency ( $\alpha = 0.84$ ; Motl et al., 2002) and construct validity (Shin, Yang, & Pender, 2001) of this measure. In the current study, this scale demonstrated high internal consistency,  $\alpha = 0.86$ .

### **Procedure**

Ethical approval was obtained from the Bond University Human Research Ethics Committee (BUHREC) prior to the commencement of intervention. Participants were recruited using an online advertisement through a variety of multimedia platforms including Facebook, LinkedIn and online community noticeboards. Participants were provided with an explanatory statement, which outlined the purposes and procedures of the research project, in order to obtain informed consent. Following completion of baseline questionnaires, participants were randomly allocated to either the intervention or control group via a computer-generated randomization. The control group consisted of participants who were not required to engage in any intervention programme that could potentially alter behavioural outcomes, therefore they did not receive the daily reminder to complete. The purpose of incorporating a control group was to make comparisons to the intervention participants who were taught the self-monitoring strategies.

During the treatment period, intervention participants were given free access to a digital database, in which participants were required to log-in daily and respond to a battery of questions related to the quality and volume of food and liquids consumed throughout the day (e.g., “How much water have you had today?”). Participants were also encouraged to document the type, intensity and duration of exercise which they engaged in throughout the day (e.g., “How much exercise have you done today?”). In the current study, exercise was defined as a measure of intensity and physical exertion, with levels of intensity ranging from strenuous to mild, where strenuous activities involved a significant increase in the participant’s heartbeat, moderate activities involved exercise that did not lead to participants feeling exhausted or fatigued upon completion, and mild activities involved minimal effort (Anderson & Cychosz, 1994). In order to ensure adherence, participants were sent a daily reminder via email to complete the daily diary. Diary entries were de-identified for the purposes of data analysis and interpretation. A unique participant code was used to match each participant’s diary entries. At the end of the 8-week intervention participants again completed the baselines measures, as did the control group who did not receive any intervention.

## **RESULTS**

### **Data Cleaning**

Prior to the above-mentioned statistical analysis being run, the data was analysed for missing data, patterned responses and unvarying data. This initial analysis revealed 22 cases with extensive missing data. These cases were removed from the data set. Furthermore, 12 cases showed patterned and unvarying data and were removed.

Therefore, the dataset was reduced from 88 to 54 cases (i.e., 29 cases from the intervention group and 25 cases from the control group). These cases were reviewed using a scatter matrix across each variable of interest. Upon further inspection of the scatter matrix, it appeared that three cases were identified as possible outliers. They did not affect the results and therefore were included in the data set, resulting in a final sample of 54 cases.

### **Descriptive Statistics**

Descriptive statistics for the intervention and control groups on each of the dependent variables are displayed in Table 1. Variables with moderate skewness ( $\pm 1.96$ ) were transformed using a square root transformation. Variables with severe skewness ( $\pm 2.58$ ) were transformed using a log transformation. Data was analysed using both the original and transformed data, and showed the log transformed data followed a normal or near normal distribution therefore, all further data analysis utilised original data. Descriptive statistics for the intervention group reveal several noteworthy findings that occurred from pre- to post-treatment, including an increase in duration of exercise.



**Table 1: Means and standard deviations for the dependent variables from the intervention and control groups at pre- and post-intervention**

| Variables                       | Pre           |        |       |                   | Post          |        |       |                   |
|---------------------------------|---------------|--------|-------|-------------------|---------------|--------|-------|-------------------|
|                                 | Mean (SD)     | 95% CI |       | Standardised Skew | Mean (SD)     | 95% CI |       | Standardised Skew |
|                                 |               | Lower  | Upper |                   |               | Lower  | Upper |                   |
| Exercise Identity Tx            | 42.87 (9.91)  | 39.10  | 46.62 | 42.87 (9.91)      | 49.34 (9.88)  | 45.58  | 53.10 | -1.04             |
| Exercise Self-efficacy Tx       | 12.10 (3.75)  | 10.68  | 13.53 | 12.10 (3.75)      | 18.07 (5.95)  | 15.81  | 20.33 | -0.09             |
| Healthy Eating Identity Tx      | 43.63 (8.19)  | 40.51  | 46.73 | 43.63 (8.19)      | 49.93 (7.22)  | 47.19  | 52.67 | -0.82             |
| Healthy Eating Self-efficacy Tx | 12.38 (2.65)  | 11.37  | 13.39 | 12.38 (2.65)      | 18.82 (4.94)  | 16.95  | 20.71 | 0.10              |
| Exercise Identity Control Tx    | 36.00 (11.98) | 31.05  | 40.94 | 0.35              | 33.92 (11.98) | 28.24  | 39.59 | 0.33              |
| Exercise Self-efficacy Control  | 12.68 (2.91)  | 11.47  | 13.88 | 0.20              | 13.04 (3.96)  | 11.41  | 14.67 | -0.16             |
| Healthy Eating Identity Control | 41.00 (9.93)  | 36.89  | 45.10 | 0.38              | 39.88 (11.71) | 35.05  | 44.71 | 0.03              |
| Healthy Eating SControl         | 14.48 (2.86)  | 13.29  | 15.66 | -0.03             | 13.44 (2.90)  | 12.24  | 14.63 | -0.45             |

Tx = Treatment

### Repeated Measures ANOVA

*Exercise identity.* Before conducting analyses, the appropriate assumptions were checked and confirmed for all variables. A 2 x 2 repeated measures ANOVA was used to investigate the impact of time on exercise identity. A significant main effect of time was obtained,  $F(1, 52) = 7.42$ ,  $p = 0.009$ , partial  $\eta^2 = 0.12$ , with exercise-identity levels at pre-intervention ( $M = 39.43$ ) being significantly lower than at post-intervention ( $M = 41.63$ ). A significant main effect of group was indicated,  $F(1, 52) = 13.82$ ,  $p < 0.001$ , partial  $\eta^2 = 0.21$ , with exercise-identity levels of intervention group participants ( $M = 46.10$ ) being significantly higher than control group participants ( $M = 34.96$ ).

A significant interaction between time and group was revealed,  $F(1, 52) = 28.08$ ,  $p < 0.001$ , partial  $\eta^2 = 0.35$ . Pairwise comparisons indicated that a significant difference ( $p < 0.001$ ) was found between the intervention and control groups ( $M = 11.14$ ). Post hoc tests revealed that exercise-identity scores for both the intervention group ( $p < 0.001$ ) and the control group ( $p < 0.001$ ) significantly increased from pre- to post-intervention but that individuals within the intervention group reported a greater sense of exercise identity than control group participants.

*Healthy eating identity.* A 2 x 2 repeated measures ANOVA was used to investigate the impact of time on healthy eating identity. A significant main effect of time was obtained,  $F(1, 52) = 12.02$ ,  $p = 0.001$ , partial  $\eta^2 = 0.19$ , with healthy eating identity at pre-intervention ( $M = 42.31$ ) being significantly lower than healthy eating identity levels at post-intervention ( $M = 44.91$ ). A significant main effect of group was indicated,  $F(1, 52) = 6.82$ ,  $p = 0.012$ , partial  $\eta^2 = 0.12$ , with healthy eating identity levels of intervention group participants ( $M = 46.78$ ) being significantly higher than control group participants ( $M = 40.44$ ).

A significant interaction between time and group was revealed,  $F(1, 52) = 24.63$ ,  $p < 0.001$ , partial  $\eta^2 = 0.32$ . Pairwise comparisons indicated that a significant difference ( $p = 0.012$ ) was found between the intervention and control groups ( $M = 6.34$ ). Post hoc tests revealed that healthy eating identity scores for the intervention group ( $p < 0.001$ ) significantly increased from pre- to post-intervention, while there was no statistically significant change in scores for the control group from pre- to post-intervention ( $p = 0.263$ ). The intervention group participants reported greater healthy eating identity than control group participants.

*Exercise self-efficacy.* A 2 x 2 repeated measures ANOVA was used to investigate the impact of time on exercise self-efficacy. A significant main effect of time was obtained,  $F(1, 52) = 28.48$ ,  $p < 0.001$ , partial  $\eta^2 = 0.35$ , with exercise self-efficacy at pre-intervention ( $M = 12.39$ ) being significantly lower than exercise self-efficacy at post-intervention ( $M = 15.54$ ). A significant main effect of group was indicated,  $F(1, 52) = 4.70$ ,  $p = 0.035$ , partial  $\eta^2 = 0.08$ , with exercise self-efficacy levels of intervention group participants ( $M = 15.09$ ) being significantly higher than control group participants ( $M = 12.86$ ).

A significant interaction between time and group was revealed,  $F(1, 52) = 22.37$ ,  $p < 0.001$ , partial  $\eta^2 = 0.30$ . Pairwise comparisons indicated that a significant difference ( $p = 0.035$ ) was found between the intervention and control groups ( $M = 2.23$ ). Post hoc tests revealed that exercise self-efficacy scores for both the intervention group ( $p < 0.001$ ) and the control group ( $p = 0.023$ ) significantly increased from pre- to post-

intervention, however, individuals within the intervention group reported greater exercise self-efficacy than control group participants.

*Healthy eating self-efficacy.* A 2 x 2 repeated measures ANOVA was used to investigate the impact of time on healthy eating self-efficacy. A significant main effect of time was obtained,  $F(1, 52) = 18.53, p < 0.001$ , partial  $\eta^2 = 0.26$ , with healthy eating self-efficacy levels pre-intervention ( $M = 13.43$ ) being significantly lower than healthy eating self-efficacy levels at post-intervention ( $M = 16.13$ ). A significant main effect of group was indicated,  $F(1, 52) = 5.19, p = 0.027$ , partial  $\eta^2 = 0.09$ , with healthy eating self-efficacy levels of intervention group participants ( $M = 15.60$ ) being significantly higher than control group participants ( $M = 13.96$ ).

A significant interaction between time and group was also found,  $F(1, 52) = 35.52, p < 0.001$ , partial  $\eta^2 = 0.40$ . Pairwise comparisons indicated that a significant difference ( $p = 0.027$ ) was found between the intervention and control groups ( $M = 1.64$ ). Post hoc tests revealed that healthy eating self-efficacy scores for the intervention group ( $p < 0.001$ ) significantly increased from pre- to post-intervention, but there was no statistically significant change in scores for the control group from pre- to post-intervention ( $p = 0.263$ ). The intervention group participants reported greater healthy eating self-efficacy than control group participants.

## DISCUSSION

This study sought to investigate whether a self-monitoring practice (i.e., daily diary entries) enhanced individuals' exercise identity, exercise self-efficacy, healthy eating identity and healthy eating self-efficacy. As expected, participants within the intervention programme displayed a significant improvement in all variables when compared with the control group.

### Identity Theory in Exercise and Healthy Eating

The intervention group participants exhibited significantly greater exercise identities when compared with the control group, and this finding is consistent with empirical studies that indicate identity can be improved through the introduction of self-awareness practices (Chang et al., 2004; Kristeller & Wolever, 2010). Given the significance of identity and self-efficacy in improving health behaviours, the findings in the present study are particularly important. The intervention group also reported greater healthy eating identity compared with the control group, and this may have implications for treatment planning, in that targeting and improving an individual's belief in his or her ability to maintain a healthy diet may lead to additional improvements in other health-related areas. Overall, health-related identity appeared to significantly improve following the intervention participants' completion of the daily diary over the 8-week period, which further implies that health-related identity may be shifted or modified using self-awareness and monitoring practices.

### Self-efficacy in Exercise and Healthy Eating

Exercise self-efficacy was significantly greater for the intervention group and this finding suggests that self-awareness and monitoring practices may be effective in improving individuals' belief in their ability to successfully achieve their exercise goals

despite the presence of potential obstacles. The intervention participants also reported greater healthy eating self-efficacy compared with the control group, and this suggests that healthy eating self-efficacy is amenable to change and improvement through self-awareness and monitoring practices, in line with established literature (e.g., Schnoll & Zimmerman, 2001).

Overall, health-related self-efficacy appeared to significantly improve following the intervention participants' completion of self-awareness and monitoring practices. Given the strong association between self-efficacy and behavioural performance as demonstrated in previous studies (see Schwarzer & Luszczynska, 2000), these findings have significant implications for improving health-related behaviours. The practical implication of this finding is that greater exercise self-efficacy is associated with increased engagement in behavioural performance, which is useful to consider when tailoring exercise and diet programmes to improving health behaviours.

### **Strengths and Limitations**

A particular strength of the current research is that, despite the smaller sample size, the study was successful in recruiting individuals across different age groups. This is likely to improve the generalisability of the study to the general population. Another strength of the current study is the nature of the study design, which allowed greater adherence to completion of daily diaries due to the convenience of the online function.

Nonetheless, the current study was not without limitations, primarily the fact that data was self-reported and thus susceptible to various effects including social desirability and impression management. This study only examined increasing attention to diet and exercise behaviours, not actually improving those behaviours if they were lacking in quality (e.g., making better dietary choices). Future research could collect this data and investigate the changes that may occur in actual behaviour alongside increasing attentional awareness.

The current study could have also been improved by collecting more longitudinal data, such as a 3- or 6-month follow-up. This may have allowed further examination of stability of change over time, and thus more powerful conclusions regarding self-awareness and regulation with behavioural change. Finally, the participant group was small, thus reliability and generalisation to a wider sample is difficult to conclude, and the inclusion of a social desirability measure may have been useful to investigate any impact of positive impression management.

## **IMPLICATIONS AND CONCLUSIONS**

Results of the current study have provided preliminary evidence for the benefits associated with self-awareness and monitoring practices in improving health-related outcomes across more than one area. The 8-week daily diary intervention was associated with improved exercise and healthy eating identity, and simultaneous improved exercise and healthy eating self-efficacy in the intervention group when compared with the control group.

There is little empirical literature examining the effect of self-awareness and monitoring strategies upon health-related identity and self-efficacy. As such, the current study sought to address limitations in previous research by investigating the impact of a daily reminder service for a website-based self-awareness strategy in improving identity and self-efficacy, and thus behavioural performance. Results may be useful in informing clinicians designing and tailoring treatment programmes for improving specific health behaviours and outcomes, and including a consistent reminder service for engaging in such behaviours.

## FIRST AUTHOR BIOGRAPHY

This project represented research conducted as part of the Masters of Clinical Psychology program for dissertation requirements. The research interest centred on whether logging daily habits would change health behaviour and identity.

## REFERENCES

- Ackard, D.M., Neumark-Sztainer, D., Story, M., & Perry, C. (2003) Overeating among adolescents: prevalence and associations with weight-related characteristics and psychological health. *Paediatrics*, 111(1): 67–74. doi:10.1542/peds.111.1.67
- Anderson, D.F., & Cychosz, C.M. (1994) Development of an exercise identity scale. *Perceptual and Motor Skills*, 78(3): 747–751. doi: 10.2466/pms.1994.78.3.747
- Anderson, D.F., & Cychosz, C.M. (1995) Exploration of the relationship between exercise behavior and exercise identity. *Journal of Sport Behaviour*, 18: 159–159. doi:10.10320091/j.1
- Anderson, D.F., Cychosz, C.M., & Franke, W.D. (1998) Association of exercise identity with measures of exercise commitment. *Journal of Sport Behavior*, 21(3): 233–241. doi:10.106.30179014.27/10
- Australian Bureau of Statistics (2012) *Australian Health Survey: Nutrition Results: Foods And Nutrients, 2011–2012*. ABS: Canberra.
- Bailis, D.S., Segall, A., & Chipperfield, J.G. (2003) Two views of self-rated general health status. *Social Science & Medicine*, 56(2): 203–217. doi:10.1016/S02779536(02)00020-5
- Bandura, A. (1997) *Self-efficacy: The Exercise of Control*. New York: Freeman.
- Baer, R.A. (2003) Mindfulness training as a clinical intervention: A conceptual and empirical review. *Clinical Psychology: Science and Practice*, 10(2): 125–143. doi:10.1093/clipsy.bpg015
- Bailis, D.S., Segall, A., & Chipperfield, J.G. (2003) Two views of self-rated general health status. *Social Science and Medicine*, 56(2): 203–217. doi: 10.1016/S02779536(02)00020-5
- Baker, R.C., & Kirschenbaum, D.S. (1993) Self-monitoring may be necessary for successful weight control. *Behavior Therapy*, 24: 377–394
- Bauman, A., Armstrong, T., Davies, J., Owen, N., Brown, W., Bellew, B., & Vita, P. (2003) Trends in physical activity participation and the impact of integrated campaigns among Australian adults, 1997–99. *Australian and New Zealand Journal of Public Health*, 27(1): 76–79. doi:10.1111/j.1467-842X.2003.tb00384
- Bere, E., & Klepp, K.I. (2004) Correlates of fruit and vegetable intake among Norwegian school children. Parental and self-reports. *Public Health Nutrition*, 7: 991–998. doi:10.1079/PHN2004619
- Bishop, S.R., Lau, M., Shapiro, S., Carlson, L., Anderson, N.D., Carmody, J., & Devins, G. (2004). Mindfulness: A proposed operational definition. *Clinical Psychology: Science and Practice*, 11(3): 230–241. doi: 10.1093/clipsy.bph077
- Blair, S.N., Kohl, H.W., Paffenbarger, R.S., Clark, D.G., Cooper, K.H., & Gibbons, L.W. (1989) Physical fitness and all-cause mortality: A prospective study of healthy men and women. *Journal of the American Medical Association*, 262(17): 2395–2401. doi:10.1001/jama.1989.03430170057028.

- Blumenthal, J.A., Babyak, M.A., Doraiswamy, P.M., Watkins, L., Hoffman, B.M., Barbour, K.A., & Sherwood, A. (2007). Exercise and pharmacotherapy in the treatment of major depressive disorder. *Psychosomatic Medicine*, 69(7): 587. doi:10.1097/PSY.0b013e318148c19a
- Bodnar, L.M., & Wisner, K.L. (2005) Nutrition and depression: Implications for improving mental health among childbearing-aged women. *Biological Psychiatry*, 58(9): 679–685. doi:10.1016/j.biopsych.2005.05.009
- Boutelle, K.N., & Kirschenbaum D.S.S. (1998) Further support for consistent self-monitoring as a vital component of successful weight control. *Obesity Research*, 6(3): 219–224.
- Briefel, R.R., & Johnson, C.L. (2004) Secular trends in dietary intake in the United States. *Journal of Nutrition*, 24: 401–431. doi:10.1146/23.011702.073349
- Brown, D.R., Wang, Y., Ward, A.N.N., Ebbeling, C B., Fortlage, L., Puleo, E., & Rippe, J.M. (1995) Chronic psychological effects of exercise and exercise plus cognitive strategies. *Medicine & Science in Sports & Exercise*, 27(5): 765–775. doi:10.1249/00005768-199505000-00021
- Cardinal, B.J., & Cardinal, M.K. (1997) Changes in exercise behavior and exercise identity associated with a 14-week aerobic exercise class. *Journal of Sport Behavior*, 20(4): 377–386. doi:10.01/2729010.13
- Carels, R.A., Darby, L.A., Rydin, S., Douglass, O.M., Cacciapaglia, H.M., & O'Brien, W.H. (2005) The relationship between self-monitoring, outcome expectancies, difficulties with eating and exercise, and physical activity and weight loss treatment outcomes. *Ann Behav Med*, 30(3):182–190.
- Chan, D.W. (2002) Stress, self-efficacy, social support, and psychological distress among prospective Chinese teachers in Hong Kong. *Educational Psychology*, 22(5): 557–569. doi: 10.1080/0144341022000023635
- Chang, M.W., Brown, R., & Nitzke, S. (2003) Participant recruitment and retention in a pilot program to prevent weight gain in low-income overweight and obese mothers. *Journal of Public Health*, 9(1): 424–439. doi:10.1186/1471-2458-9-424
- Chang, V.Y., Palesh, O., Caldwell, R., Glasgow, N., Abramson, M., Luskin, F., & Koopman, C. (2004) The effects of a mindfulness based stress reduction program on stress, mindfulness self efficacy, and positive states of mind. *Stress and Health*, 20(3): 141–147. doi: 10.1002/smi.1011
- Cohen, S., Kamarck, T., & Mermelstein, R. (1983) A global measure of perceived stress. *Journal of Health and Social Behavior*, 385–396. doi:10.180281/1968/jt
- DeVellis, B.M., & DeVellis, R.F. (2000) *Handbook of health psychology*. Mahwah, NJ: Erlbaum.
- Dillman, H. (2012) Stability of exercise identity and healthy-eater identity over six months among rural hospital employees. *Journal of Kinesiology*, 12: 104–113. doi:10.19030.1902
- Feltz, D.L., & Riessinger, C.A. (1990) Effects of *in vivo* emotive imagery and performance feedback on self-efficacy and muscular endurance. *Journal of Sport & Exercise Psychology*, 12: 132–143. doi:10.1905/01851
- Gauvin, L, & Spence, J.C. (1996) Physical activity and psychological well-being: Knowledge base, current issues, and caveats. *Nutrition Reviews*, 54(4): 53–65. doi:10.1111/j.1753-4887.1996.tb03899.x
- Gortmaker, S.L., Swinburn, B.A., Levy, D., Carter, R., Mabry, P.L., Finegood, D.T., & Moodie, M.L. (2011) Changing the future of obesity: Science, policy, and action. *The Lancet*, 378(9793): 838–847. doi:10.1016/S0140-6736(11)60815-5
- Hassmén, P., Koivula, N., & Uutela, A. (2000) Physical exercise and psychological well-being: A population study in Finland. *Preventive Medicine*, 30(1): 17–25. doi: 10.1006/pmed.1999.0597
- House, J.S., Wells, J.A., Landerman, L.R., McMichael, A.J. & Kaplan, B.H. (1979) Occupational stress and health among factory workers. *Journal of Health and Social Behaviour*, 20(2): 139–160. doi: 10.1036/29.2/820n
- Hunter, C.L., Goodie, J.L., Oordt, M.S., & Dobmeyer, A.C. (2009) *Integrated Behavioural Health In Primary Care: Step-by-step Guidance for Assessment and Intervention*. Washington: American Psychological Association.
- Ievers-Landis, C.E., Burant, C., Drotar, D., Morgan, L., Trapl, E.S., & Kwoh, C.K. (2003) Social support, knowledge, and self-efficacy as correlates of osteoporosis preventive behaviours among preadolescent females. *Journal of Pediatric Psychology*, 28: 335–345. doi: 10.1093/jpepsy/jsg023
- Johansson, G., Johnson, J.V., & Halls, E.M. (1991) Smoking and the sedentary behaviour as related to work organization. *Social Science and Medicine*, 32(7): 837–846. doi:10.1016/0277-9536(91)90310-9

- Kennedy, E. (2006) Evidence for nutritional benefits in prolonging wellness. *American J. Clinical Nutrition*, 83(2), 410S-414S.
- Kristeller, J.L., & Wolever, R.Q. (2010) Mindfulness-based eating awareness training for treating binge eating disorder: The conceptual foundation. *Eating Disorders*, 19(1): 4961. doi:10.1080/10640266.2011.533605
- Kroll, T., Kehn, M., Ho, P.S., & Groah, S. (2007) The exercise self-efficacy scale (ESES): development and psychometric properties. *International Journal of Behavioral Nutrition and Physical Activity*, 4(1): 34. doi:10.1186/1479-5868-4-34.
- Lawlor, D.A., & Hopker, S.W. (2001) The effectiveness of exercise as an intervention in the management of depression: Systematic review and meta-regression analysis of randomised controlled trials. *British Medical Journal*, 322(72): 763–771. doi:10.1136/bmj.322.7289.763
- Leino-Arjos, P., Solovieva, S., Riihimaki, H., Kirjonen, J., & Telema, R. (2004) Leisure time physical activity and strenuousness of work as predictors of physical functioning: A 28-year follow up of a cohort of industrial employees. *Occupational and Environmental Medicine*, 61: 1032–1038. doi:10.1136/oem.2003.012054
- Leyse-Wallace, R. (2013) *Nutrition and Mental Health*. Florida: CRC Press. Taylor & Francis Group.
- Lubans, D.R., Plotnikoff, R.C., Morgan, P.J., Dewar, D., Costigan, S., & Collins, C.E. (2012) Explaining dietary intake in adolescent girls from disadvantaged secondary schools. A test of Social Cognitive Theory. *Appetite*, 58: 517–524. doi:10.1016/j.appet.2011.12.012
- Martin, K.A., Leary, M.R., & O'Brien, J. (2001) Role of self-presentation in the health practices of a sample of Irish adolescents. *Journal of Adolescent Health*, 28(4): 259–262. doi:10.1016/S1054-139X(00)00209-3
- Martinsen, E.W. (2008) Physical activity in the prevention and treatment of anxiety and depression. *Nordic Journal of Psychiatry*, 62(47): 25–29. doi:10.1080/08039480802315640
- McAuley, E., Lox, C., & Duncan, T.E. (1993) Long-term maintenance of exercise, self-efficacy, and physiological change in older adults. *Journal of Gerontology*, 48(4): 218–224. doi:10.1093/geronj/48.4
- Merom, D., Phongsavan, P., Wagner, R., Chey, T., Marnane, C., Steel, Z., & Bauman, A. (2008) Promoting walking as an adjunct intervention to group cognitive behavioural therapy for anxiety disorders—a pilot group randomized trial. *Journal of Anxiety Disorders*, 22(6): 959–968. doi: 10.1016/j.janxdis.2007.09.010.
- Miller, K.H., Ogletree, R.J., & Welshimer, K. (2002) Impact of activity behaviors on physical activity identity and self-efficacy. *American Journal of Health Behavior*, 26(5): 323–330. doi: 10.5993/AJHB.26.5.1
- Miller, C.K., Kristeller, J.L., Headings, A., Nagaraja, H., & Miser, W.F. (2012) Comparative effectiveness of a mindful eating intervention to a diabetes self management intervention among adults with type 2 diabetes: A pilot study. *Journal of the Academy of Nutrition and Dietetics*, 112(11): 1835–1842. doi:10.1080/00223980309600624
- Motl, R.W., Dishman, R.K., Saunders, R.P., Dowda, M., Felton, G., Ward, D.S., & Pate, R.R. (2002) Examining social-cognitive determinants of intention and physical activity among black and white adolescent girls using structural equation modelling. *Health Psychology*, 21(5): 459–471. doi: 10.1037/0278-6133.21.5.459
- Mustard, C.A., Vermeulen, M., & Lavis, J.N. (2003) Is position in the occupational hierarchy a determinant of decline in perceived health status? *Social Science and Medicine*, 57(12): 2291–2303. doi:10.1016/j.socscimed.2003.08.001
- Oliver, G., Wardle, J., & Gibson, E.L. (2000) Stress and food choice: A laboratory study. *Psychosomatic Medicine*, 62(6): 853–865. doi:10.1022/0893-3200.62.6.853
- Peluso, M.A., & Andrade, L.H. (2005) Physical activity and mental health: the association between exercise and mood. *Clinics*, 60(1): 61–70. DOI:10.1590/S180759322005000100012
- Polman, R., Borkoles, E., & Nicholls, A.R. (2010) Type D personality, stress, and symptoms of burnout: The influence of avoidance coping and social support. *British Journal of Health Psychology*, 15(3): 681–696. doi:10.1348/135910709X479069
- Pirkis, J., Blood, R.W., Dare, A., & Holland, K. (2008) The Media Monitoring Project: Changes in media reporting of suicide and mental health and illness in Australia: 2000/01–2006/07. Canberra, ACT: Commonwealth Department of Health and Aged Care.
- Raglin, J.S. (1990). Exercise and mental health. *Sports Medicine*, 9(6): 323–329. doi: 10.2165/00007256-199009060-00001
- Remor, E. (2006). Psychometric properties of a European Spanish version of the Perceived Stress Scale (PSS). *The Spanish Journal of Psychology*, 9(1): 86–93. doi:10.1017/S1138741600006004

- Saksvig, B.I., Gittelsohn, J., Harris, S.B., Hanley, A.J., Valente, T.W., & Zinman, B. (2005) A pilot school-based healthy eating and physical activity intervention improves diet, food knowledge, and self-efficacy for native Canadian children. *The Journal of Nutrition*, 135(10): 2392–2398. doi: 10.1016/j.jneb.2011.03.143
- SANE Australia. (2012) Fact Sheet 5: Mental Illness and Violence. South Melbourne, VIC: SANE Australia <http://www.mindframe-media.info/for-media/reporting-mental-illness/facts-and-stats#sthash.190nxgST.dpuf>
- Schnoll, R., & Zimmerman, B.J. (2001) Self-regulation training enhances dietary self-efficacy and dietary fiber consumption. *Journal of the American Dietetic Association*, 101: 1006–1011. doi: 10.1016/S0002-8223(01)00249-8
- Schwarzer, R. & Luszczynska, A. (2000) The self-efficacy scale: Multicultural validation studies. *The Journal of Psychology*, 139(5): 439–457. doi:10.3200/JRLP.139.5.439
- Schwarzer, R., & Renner, B. (2000) Social-cognitive predictors of health behavior: Action self-efficacy and coping self-efficacy. *Health Psychology*, 19(5): 487–495. doi:10.1037/0278-6133.19.5.487
- Schwarzer, R., & Luszczynska, A. (2005) Social cognitive theory. *Predicting Health Behaviour*, 2: 127–169. doi: 10.1016/j.04.12.016
- Shin, Y., Jang, H., & Pender, N.J. (2001). Psychometric evaluation of the exercise self-efficacy scale among Korean adults with chronic diseases. *Research in Nursing & Health*, 24(1): 68–76. doi: 10.1002/1098-240X(200102)
- Smith, L.L. (2003). Overtraining, excessive exercise, and altered immunity. *Sports Medicine*, 33(5): 347–364. doi: 10.2165/00007256-200333050-00002
- Strachan, S.M., & Brawley, L.R. (2008). Reactions to a perceived challenge to identity: A focus on exercise and healthy eating. *Journal of Health Psychology*, 13(5): 575–588. doi: 10.1177/1359105308090930
- Strachan, S.M., & Brawley, L.R. (2009). Healthy-eater identity and self-efficacy predict healthy eating behavior: A prospective view. *Journal of Health Psychology*, 14(5): 684–695. doi: 10.1177/1359105309104915
- Thoits, P.A. (2010). Stress and health major findings and policy implications. *Journal of Health and Social Behavior*, 51(1): 41–53. doi:10.1177/0022146510383499
- Timmerman, G.M., & Brown, A. (2012). The effect of a mindful restaurant eating intervention on weight management in women. *Journal of Nutrition Education and Behavior*, 44(1), 22–28. doi: 10.1016/j.jneb.2011.03.143
- Wardle, J., Steptoe, A., Oliver, G., & Lipsey, Z. (2000). Stress, dietary restraint and food intake. *Journal of Psychosomatic Research*, 48(2): 195–202. doi: 10.1016/S00223999(00)00076-3
- Williams, K.E., & Bond, M.J. (2002). The roles of self-efficacy, outcome expectancies and social support in the self-care behaviors of diabetics. *Psychology, Health & Medicine*, 7, 127–141. doi: 10.1080/13548500120116076
- World Health Organisation (2014). Obesity and overweight fact sheets, 2008-12. Retrieved from <http://www.who.int/en/>