

Bond University
Research Repository



Australian Army recruit training: course length and recruit injury rates

Dawson, Georgina M E; Broad, Ryan; Orr, Rob Marc

Licence:
CC BY-ND

[Link to output in Bond University research repository.](#)

Recommended citation(APA):
Dawson, G. M. E., Broad, R., & Orr, R. M. (2015). *Australian Army recruit training: course length and recruit injury rates*. CONNECT 2015 Physiotherapy Conference, Gold Coast, Queensland, Australia.

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.


For more information, or if you believe that this document breaches copyright, please contact the Bond University research repository coordinator.

Australian Army recruit training course length and recruit injury rates



Dawson, G.¹, Broad, R.² & Orr, R.¹

- 1. Bond University**
- 2. Kapooka Military Area**



Dawson, G., Broad, R. & Orr, R. M. (2014). Australian Army recruit training course length and recruit injury rates. *Journal of Military and Veteran's Health*, 23(2) 14-19.



Background


- Military recruits are at a greater risk of injury when compared to qualified soldiers

(Orr & Pope, 2015; Booth et al., 2006; Kaufman et al., 2000)

- The sudden increase in load may lead to over training and eventual injury

(Prigg et al. 2000)

- Would decreasing load but retaining training requirements decrease risk of injury?



Aims and Hypothesis

- Aims:
 - Investigate the influence of lengthening a recruit training course from 80 days to 100 days
 - Profile injuries that occur
- Hypothesis:
 - The longer the training period, the greater the risk of injury.





Participants

- Australian Regular Army recruits attending Basic Recruit training at Kapooka
- Recruits were randomly selected for each course

Course	Number of Platoons	Number of Recruits	Male Recruits	Female Recruits
ASC	2	73	56	17
ARC	4	194	152	42
Total	6	267	208	59



Methods

- Data recorded during two different Army recruit training courses over 1 year period (2013)
 - *ASC (100 d) / ARC (80 d)*
- The ASC contained all aspects of the ARC with a more gradual increase in load over the first four weeks
....also contained additional military field training and an extended field phase





Methods

- Injury Prevalence

- *Number of reported injuries / number of recruits completing the respective course x 100*

- Injury Incidence

- *(Number of reported injuries / number of recruits completing the respective course x 100 (soldiers)) / (course length in days / 100 days)*



Methods

- Ethics approval from BUHREC & ADHREC





Results


- **ASC (100 days):**
 - 73 recruits, nil excluded
 - 13 recruits injured (17.8%)

- **ARC (80 days):**
 - 194 recruits (23 excluded from original data set)
 - 27 recruits injured (13.9%)





Results

- Injury prevalence:
 - ASC: 17.8%
 - ARC: 13.9%
- Injury incidence:
 - ASC: 17.8 / 100 soldiers / 100 days
 - ARC: 17.4 / 100 soldiers / 100 days




Results





Results





Results

- Collectively the highest anatomical injury sites:
 - Ankle/foot: 20%
 - Back/torso: 12.5%
 - Lower leg: 12.5%
- ASC (100 day):
 - Back/torso: 30.7 %
 - Ankle/foot 15.4 %
 - Shoulder 15.4 %
- ARC (80 day):
 - Ankle/foot 22.2 %
 - Lower leg 14.8 %



Discussion

- ASC had a notably higher prevalence of injuries compared to ARC
- However when looking at cohort size and exposure to training, both courses had similar incidence rates
- In contrast to previous studies, the current study revealed much lower prevalence and incidence rates



Discussion

- Prevalence

- Current study: ASC: 17.8 % & ARC: 13.9% prevalence
- Havenetidis et al. (2011): 233 male Greek army recruits, 7 week course - 28.3% prevalence

- Incidence

- Current study: ASC: 17.8 % & ARC: 17.4% incidence
- O'Connor et al. (2000): 480 Marine Corp officers, 6 week course - 60.7% incidence



Discussion

- **Most common anatomical sites of injuries:**
 - Current study: Ankle and foot= 20 %
 - Similar to Havenetidis et al. (2011), O'Connor et al. (2000) and the Australian Department of Defence (2000) the most common injury sites were to the ankle and foot.
 - These sites of injury were also in the top three injury sites in studies by Ross & Allsopp (2002) and Knapik et al. (2001)




Discussion

- Most common type of injuries:
 - Current study:
 - Sprains and strains
 - Stress fractures
 - These injury types were also found to be the most common type in studies conducted by Havenetidis et al. (2011) and O'Connor et al. (2000)



Conclusion / Take Home Message


- While the ASC had a higher prevalence of injury when injuries took into account exposure, incidence rates were virtually identical
- Lengthening a recruit training program (or period of training) with the aim of making it less intensive may not reduce the proportion of recruits injured - in fact, a higher proportion may be injured due to the longer period of exposure to training.



References



- Booth CK, Probert B, Forbes-Ewan C, Coad RA. Australian army recruits in training display symptoms of overtraining. *Military medicine*. 2006;171(11):1059-1064.
- Dawson, G., Broad, R. & Orr, R. M. (2014). Australian Army recruit training course length and recruit injury rates. *Journal of Military and Veteran's Health*, 23(2) 14-19.
- Havenetidis K, Kardaris D, Paxinos T. Profiles of musculoskeletal injuries among Greek Army officer cadets during basic combat training. *Military medicine*. 2011;176(3):297-303.





References

- Kaufman KR, Brodine S, Shaffer R. Military training-related injuries: surveillance, research, and prevention. *American journal of preventive medicine*. 2000;18(3):54-63.
- Knapik JJ, Sharp MA, Canham-Chervak M et al. Risk factors for training-related injuries among men and women in basic combat training. *Journal of the American College of Sports Medicine*. 2001; 33(6): 946-954.
- O'Connor F. Injuries during Marine Corps officer basic training. *Mil Med*. 2000;165(7):515-520.



References

- Orr RM, Pope R. Optimizing the Physical Training of Military Trainees. *Strength & Conditioning Journal*. 2015;37(4):53-59.
- Ross RA, Allsopp A. Stress fractures in Royal Marines Recruits. *Journal of Military Medicine*. 2002;167(7):560-565.