

Bond University
Research Repository



Profiling a workplace physiotherapy and rehabilitation program within a police force

Hua, M; Orr, Rob Marc; Stierli, Michael

Licence:
CC BY-ND

[Link to output in Bond University research repository.](#)

Recommended citation(APA):
Hua, M., Orr, R. M., & Stierli, M. (2015). *Profiling a workplace physiotherapy and rehabilitation program within a police force*. CONNECT 2015 Physiotherapy Conference, Gold Coast, Queensland, Australia.

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

For more information, or if you believe that this document breaches copyright, please contact the Bond University research repository coordinator.

Profiling a Workplace Physiotherapy and Rehabilitation Program within a Police Force



Hua M¹, Orr R¹, Stierli, M².

1 Bond University, Gold Coast

2 New South Wales Police, Sydney





Introduction

- Initial pilot study research with NSW Police found that injured officers treated within the workplace had notably better Physical and Mental RTW scores (FMS and SF-36) and even RTW prospects

(Orr, et al., 2013)

- NSW Police implementing RECOND program



Introduction

- Aim of research:
 - Profile police attendees of a workplace physiotherapy and reconditioning program in order to guide future treatment strategies





Participants

- Injured police officers attending workplace rehabilitation (August to December 2014)
 - Inclusion Criteria: Musculoskeletal injury
 - Exclusion Criteria: Illness or diagnosed mental health injury
- 30 Male / 12 Female NSW Police Force Officers
 - Male n=30: mean age 43.3 ± 9.56 years:
 - Female n=12, mean age 38.2 ± 6.39 years



Methods

1) Primary Interview

- Type of injury (classified according to body region)
- Gender, DOB, rank, work status & years of service
- Use of load bearing vest (LBV), hip or thigh holster

2) Height & Weight

- to calculate BMI

3) Nominal Roll & Attendance (Aug – Dec 14)

- Number of treatment sessions booked/ attended





Results

- Number of Treatments attended = 296/340 (87%)
- Lumbar spine injury = 40.5% (n=17); 119 Rx attended
 - Mean number of Rx attended = 7.0 ± 3.71
- Lumbar spine injury occurred across all ranks & groups for years of service
 - Highest prevalence among SGTs (29% of all Lx Sp inj.)
 - More frequent in 0-10 yrs and 21-30 yrs of service



Results

- BMI:
 - 57.5% (n=23) → 25.0 – 29.9 (overweight)
- Males vs. Females:
 - ♀ > mean number of Rx attended all injuries than ♂
(8.25 ± 5.12 vs. 6.57 ± 4.03)





Results

- **LBV:**
 - 31% (n=13); No LBV: 69% (n=29)
 - Low back injury > when no LBV used vs. when LBV used (44.8% vs. 30.8%)
- **Thigh Holster:**
 - 35.7% (n=15); Hip Holster: 64.3% (n=27)
 - Low back injury > when hip holster used vs. when thigh holster used (44.4% vs. 33.3%)



Discussion

- **Lx Sp injury** = most commonly reported WMSD in NSW Police officers attending a workplace-led physiotherapy & rehabilitation program
 - Anderson et al. (2011) - high incidence of lower back pain associated with many occupational stressors or lifestyle related issues
 - Burton et al. (1998) - chronic low back pain (CLBP) was associated with length of service (due to reoccurrence of previous injury)



Discussion

- Female officers showed higher average for number of Rx attended
 - Feuerstein, et al. (1997) - higher overall and musculoskeletal-related disability risk in women US Army Personnel



Discussion

- Research on military populations indicates that WMSD represent a prevalent source of outpatient visits, **lost work time**, hospitalisation & disability (Feuerstein et al., 1997)
 - The findings of this study support the use of an in-house physiotherapy & rehabilitation program – eliminates cost of travel to external physio services
 - Mean cost per person = \$711.81 , Revised mean cost per person = \$394.67; Savings =\$317.14 per person



Conclusions and Practical Applications

- Injuries to the lumbar spine were the most common presentation in a police workplace rehabilitation service
 - more often associated with wearing a hip holster than wearing a thigh holster
- Workplace rehabilitation services for injured police officers can limit lost productivity and travel costs associated with travel to external services during work time



References

- Anderson, G., Zutz, A., & Plecas, D. Police officer back health. *The Journal of Criminal Justice Research*. 2 (1): 1-17. 2011.
- Burton, A. K., Tillotson, K. M., & Troup, J. D. Prediction of low back trouble frequency in a working population. *Spine*. 14 (9): 939-946. 1998.
- Feuerstein, M., Berkowitz, S. M., & Peck, C. A. Jr. Musculoskeletal-related disability in US Army personnel: prevalence, gender, and military occupational specialties. *Journal of Occupational and Environmental Medicine*. 39 (1): 60-78. 1997.
- Orr R, Stierli M, Amabile ML, Wilkes B. The impact of a structured reconditioning program on the physical attributes and attitudes of injured police officers: A pilot study. *Journal of Australian Strength and Conditioning*. 2013;21(4):42.