

Bond University
Research Repository



Cost-effectiveness of food, supplement and environmental interventions to address malnutrition in residential aged care: A systematic review

Hugo, Cherie; Isenring, Elisabeth; Miller, Michelle; Marshall, Skye

Published in:
Age and Ageing

DOI:
[10.1093/ageing/afx187](https://doi.org/10.1093/ageing/afx187)

Licence:
Other

[Link to output in Bond University research repository.](#)

Recommended citation(APA):
Hugo, C., Isenring, E., Miller, M., & Marshall, S. (2018). Cost-effectiveness of food, supplement and environmental interventions to address malnutrition in residential aged care: A systematic review. *Age and Ageing*, 47(3), 356-366. <https://doi.org/10.1093/ageing/afx187>

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

For more information, or if you believe that this document breaches copyright, please contact the Bond University research repository coordinator.

1 **Actual title:** Cost-effectiveness of food, supplement and environmental interventions to
2 address malnutrition in residential aged care: a systematic review

3

4 **Authors:** Cherie Hugo^{ab}, Elisabeth Isenring^{ac}, Michelle Miller^d and Skye Marshall^{ae}

5 **Journal:** Age and Ageing Journal

6 ^a Faculty of Health Sciences and Medicine, Bond University, Robina, Queensland, 4226, Australia.

7 ^b BHSc, MasterNutrDiet, PhD Scholar, Accredited Practising Dietitian. chugo@bond.edu.au

8 ^c Professor of Nutrition & Dietetics, PhD, Advanced Accredited Practising Dietitian

9 ^d Professor of Nutrition & Dietetics, Dean of People and Resources at Flinders University, PhD,
10 Advanced Accredited Practising Dietitian

11 ^e BNutr&Diet(Hons), PhD, Accredited Practising Dietitian

12

13 **Contributors and their role in the paper**

14 This study was conceptualised by CH, LI, MM and SM.

15 CH and SM developed and ran search strategy.

16 CH and SM independently screened, extracted and synthesised data before meeting to compare
17 results and resolve minor conflicts in interpretation of results.

18 CH and SM compiled the original manuscript. LI and MM contributed towards the manuscript
19 revision. All authors approved the final manuscript.

20 **Conflicts of Interest** – The authors have no conflict of interests to declare.

21 **Funding Information** - This research did not receive any specific grant from funding agencies
22 in the public, commercial, or not-for-profit sectors.

1 **Word Count** - 2772 words (excl abstract, highlights, title page, references), 250 words

2 (abstract)

3 **Tables and Figures** – 2 Tables, 2 Figures, 1 Supplementary File.

4

5

1 **Abstract**

2 **Background:**Observational studies have shown that nutritional strategies to manage
3 malnutrition may be cost-effective in aged care; but more robust economic data is needed to
4 support and encourage translation to practice. Therefore, the aim of this systematic review is to
5 compare the cost-effectiveness of implementing nutrition interventions targeting malnutrition
6 in aged care homes versus usual care.

7 **Setting:**Residential aged care homes.

8 **Methods:**Systematic literature review of studies published between January 2000 - August
9 2017 across 10 electronic databases. Cochrane Risk of Bias tool and GRADE were used to
10 evaluate the quality of the studies.

11 **Results:**Eight included studies (3098 studies initially screened) reported on 11 intervention
12 groups, evaluating the effect of modifications to dining environment (n=1), supplements (n=5)
13 and food-based interventions (n=5). Interventions had a low cost of implementation
14 (<£2.30/resident/day) and provided clinical improvement for a range of outcomes including
15 weight, nutritional status and dietary intake. Supplements and food-based interventions further
16 demonstrated a low cost per quality adjusted life year or unit of physical function improvement.
17 GRADE assessment revealed the quality of the body of evidence that introducing malnutrition
18 interventions, whether they be environmental, supplements or food-based, are cost-effective in
19 aged care homes was low.

20 **Conclusion:**This review suggests supplements and food-based nutrition interventions in the
21 aged care setting are clinically effective, have a low cost of implementation and may be cost-
22 effective at improving clinical outcomes associated with malnutrition. More studies using well-
23 defined frameworks for economic analysis, stronger study designs with improved quality,
24 along with validated malnutrition measures are needed to confirm and increase confidence with
25 these findings.

26

27

28 **Keywords** – Malnutrition; Systematic Review; Cost; Aged Care; Economic

29 **Key Points** –

- 1 - Malnutrition is a significant economic burden on society.
- 2 - Nutrition offers opportunity to improve the quality of life of residents and the economic
- 3 position of aged care homes.
- 4 - Quality economic studies evaluating malnutrition interventions in the aged care setting are
- 5 lacking.
- 6 - More robust malnutrition economic evaluation intervention studies in aged care are needed to
- 7 support research translation.
- 8
- 9

1 **Introduction**

2 The financial cost of residential aged care, accommodation and care support for frail and aged
3 residents, is high and increasing [1-3]. Whilst significant resources go towards supporting the
4 health of older residents, outcomes are often suboptimal and associated with malnutrition
5 (undernutrition). Malnutrition is a wasting syndrome which presents most commonly in older
6 adults, and occurs when lean body mass, with or without fat mass, is unintentionally lost due to
7 inadequate bioavailability of energy and protein [4]. Cost-of-illness studies indicate that the
8 annual direct cost of malnutrition in residential aged care ranges from €107 million to €1.7
9 billion (£98.4 million to £1.56 billion) for the Netherlands and the UK respectively [5-8].

10 Higher food budgets (>£4.20 per resident per day) in aged care homes decrease the risk of a
11 resident becoming malnourished by 66% (OR: 0.66 [95%CI: 0.46-0.95] P=0.023) [9]. Recent
12 research in developed countries demonstrate a downward trend in the amounts spent on the
13 food budget in aged care homes [10]. There is also an increase in spending on oral nutrition
14 supplements (“supplements”) which is believed to be in response to high malnutrition rates
15 [10]. There is evidence that interventions such as supplements, food-first approaches
16 (prioritising food over supplements) and environmental changes improve clinical outcomes for
17 residents in resident aged care homes [11]. In acute care, these malnutrition interventions are
18 ranked as one of the top strategies to produce health care cost savings by the National Institute
19 for Health and Care Excellence (NICE) [12]. Observational studies have shown that nutritional
20 strategies to manage malnutrition may be cost-effective in the aged care setting; but more
21 robust economic data is needed to support and encourage translation to practice [13-16].
22 Therefore, the aim of this systematic review is to compare the cost-effectiveness of
23 implementing nutrition interventions (including food fortification, supplements, menu changes
24 and dining environment changes) targeting malnutrition in aged care homes versus no
25 intervention or usual care for older residents.

26 **Methods**

27 A systematic review was planned and reported according to the PRISMA guidelines [17]. The
28 protocol for this review was developed in consultation with topic experts and the search
29 strategy was developed in consultation with an information specialist. The protocol was
30 registered with PROSPERO (<http://www.crd.york.ac.uk/PROSPERO>) (registration number -
31 CRD42016048175).

32 *Search strategy*

1 Published studies were searched for in the following electronic databases: MEDLINE
2 (PubMed), Cochrane, CINAHL, EMBASE, EBSCO Megafire Complete, Business Source
3 Complete, EconLit, NHS EED and Web of Science from January 2000 to 24 August 2017
4 Publications predating 1 January 2000 were excluded as health inflation analysis has shown
5 that the health sector prices have grown much faster than inflation, the population, population
6 ageing and the broader economy in the past 15 years [18]. As a result of the documented year-
7 on-year health cost increases, comparison of data prior to the 2000 would be difficult. No
8 language restrictions were used.

9 The search strategy used keywords and each database's controlled vocabulary (Online
10 Supplementary Material 1). The search strategy was complemented by a "snowball" search
11 which involved pursuing article references of identified studies in addition to electronic citation
12 tracking and brief Google Scholar searches. For this review, nutrition interventions to prevent
13 and/or treat malnutrition in older residents (mean age of sample ≥ 65 years) dwelling in a
14 residential aged care home were included. Eligibility criteria included studies that had original
15 financial data related to the intervention and/or outcomes. Specifically, studies were included
16 which reported data related to the direct cost, cost-effectiveness and/or cost-benefit of the
17 interventions. Due to differences in economies, studies implemented in developing countries
18 were excluded. Reviews, observational studies, abstracts and conference papers were also
19 excluded from the review.

20 *Selection of studies and data synthesis*

21 After citations were identified from all databases, duplicates were isolated and removed. A
22 two-step screening process was employed. In step 1, two researchers (CH and SM) scanned the
23 titles and abstracts of studies identified by the search for their potential eligibility. At step 2,
24 full-text articles relating to the inclusion and exclusion criteria were screened by two
25 researchers for eligibility (CH and SM). Conflicts between the two screening authors were
26 resolved through consensus.

27 A list of outcomes meaningful to the research aim was developed to identify the relevant
28 effects of the interventions. The primary outcomes were financial and economic data relating to
29 the interventions, including the direct cost of implementing the intervention, the cost of usual
30 care/no intervention, the mean difference between intervention and control, the cost associated
31 with negative patient outcomes related to malnutrition, the cost-savings relating to malnutrition
32 outcomes, the cost per quality adjusted life years (QALYs) and disability adjusted life years


1 (DALYs) associated with the intervention. Secondary outcomes included patient, health and
2 aged care related outcomes associated with malnutrition, including nutrition status, weight
3 change, BMI, energy and/or protein intake, plate wastage, resident satisfaction, staff
4 satisfaction, acute and sub-acute hospital admissions, a change in the level of aged care
5 provided, quality of life, physical function, mental health, self-efficacy, mortality, and
6 malnutrition-related complications such as pressure ulcers, poor wound healing, oedema and
7 falls.

8 Data related to the primary and secondary outcome measures, the study populations and the
9 intervention details were extracted from the published papers into standardised tables by one
10 researcher (CH) and checked for accuracy by a second researcher (SM).

11 *Review of study strength and quality*

12 Risk of bias of individual studies was assessed using the Cochrane Risk of Bias tool [19]
13 covering six domains of bias: selection, performance, detection, attrition, reporting, and other
14 bias (e.g. funding sources, conflicts of interest). The quality of the body of evidence for each
15 type of intervention and outcome was determined using the Grading of Recommendations,
16 Assessment, Development and Evaluation (GRADE) system rated from very low to high
17 quality based on study design, reporting risk of bias, consistency, directness, effect size, and
18 precision [20]. The GRADE system is a formal process to rate the quality of scientific evidence
19 in systematic reviews [20].

20



15 **Figure 1:** Prisma flow diagram of records identified, screened and included in this systematic
 16 literature review.

18 **Results**

19 The search identified 3,098 records (Figure 1). Of these, 87 were considered suitable for full
 20 text review following removal of duplicates and initial screening of title/abstracts. From these
 21 papers, eight intervention studies met eligibility criteria. Due to inconsistent intervention
 22 approaches and methods of reporting cost-related outcomes, data could not be pooled.
 23 Interventions ranged in duration from 6-weeks to 6-months with follow-up ranging from 10 to
 24 29 weeks (Table 1). The seven intervention studies were from USA (n=3), Taiwan (n=1),
 25 Sweden (n=2), Netherlands (n=1) and United Kingdom (n=1) with a total of 774 enrolled older
 26 adults.

27 *Study Quality (Risk of Bias)*

1 Of the studies reviewed, four were RCTs [21-24], three were non-randomised controlled trials
 2 [25-27] and one was a 2-armed non-controlled intervention trial [28]. There was a high risk of
 3 bias across studies, particularly with lack of, and poor description of, randomisation and
 4 blinded allocation, intervention and assessment of outcomes (Figure 2; justifications in Online
 5 Supplementary Material 2). There was also a high risk of bias regarding outcomes (detection
 6 bias), as several studies did not use systematic or validated methods to measure and report
 7 financial data. Other bias considered included funding sources and conflicts of interest.

	Random sequence generation (selection bias)	Allocation concealment (selection bias)	Blinding of participants and personnel (performance bias)	Blinding of outcome assessment (detection bias)	Incomplete outcome data (attrition bias)	Selective reporting (reporting bias)	Other bias
Edwards & Beck 2002	-	-	?	?	?	+	?
Elia et al 2017	+	+	-	-	+	-	-
Lee et al 2013	+	?	+	+	+	+	+
Lorefalt et al 2011	-	?	?	?	+	+	?
Neelemaat et al 2012	+	+	-	-	+	+	+
Odlund Olin et al 2003	-	?	?	?	+	+	?
Simmons et al 2010	?	?	-	?	+	+	+
Simmons et al 2015	+	?	?	?	+	+	+

8

9 **Figure 2:** Risk of bias of included studies

10

11

12 *Types of interventions*

1 The eight studies included 11 intervention groups – supplements (n=5), food-based
2 interventions (n=5) and dining environment changes (n=1).

3 The study incorporating dining environment changes involved the addition of fish aquariums
4 into the dining area of three dementia units, with no other intervention factors.

5 There were five studies which used ONS, one combined with a high protein-high energy diet
6 and post-hospital discharge telehealth (Table 1). This study was primarily implemented in the
7 hospital setting with post-discharge ONS and fortnightly telehealth consultations from a
8 dietitian to participants, but it is unclear if the telehealth was provided to aged care home-
9 dwelling residents or only those in their own homes [22]. The supplements used had 9.5g to
10 12g of protein and 250-330kcal; however, two studies did not specify the nutritional content of
11 the ONS used. Timing and dosage of ONS interventions varied from one to two a day, and
12 from weekends only to daily.

13 Food-based interventions were simple, and included offering additional appetisers and snacks,
14 providing advice to eat high protein-high energy foods, and fortifying usual meals with cream
15 and butter. However, one food-based study implemented three 2-hour education sessions to
16 staff promoting nourishing snacks for residents [27] to support the provision of additional
17 foods; and the high protein-high energy intervention received the advice from a dietitian at two
18 time-points over 3-months [28]. The group which received food fortification received an
19 additional 2100kcal/day; however, no other study reported the additional protein or energy
20 provided.

21 *Financial outcomes by intervention type*

22 Cost data was largely heterogeneous in terms of costs measured, analysis method and style of
23 reporting which prevented synthesis or identification of a consistent finding across studies
24 (Table 1).

25 The one environmental study reported a cost saving of \$11.44 (assumed USD; £8.93) in
26 decreased ONS use; however, this was measured in one third of the group only [25]. The
27 quality of the evidence that the true financial effect of environmental interventions to improve
28 nutrition was assessed as very low, downgraded due to uncertainty across most domains
29 assessed by GRADE (Table 2).

30 Three of the studies which used an ONS intervention reported direct cost of the intervention,
31 with a difference of USD\$0.40 (£0.10) to USD\$2.54 (£1.99) per resident per day between

1 intervention and control groups [24, 29, 30]. Elia et al [28] reported a direct cost of £162.30 per
2 resident across 12-weeks (estimated as £1.93 per resident per day); but did not compare this
3 with a control. Four ONS intervention studies also included cost-effectiveness analyses. The
4 study by Neelemaat et al [22] reveals that the study may be cost-effective in improving
5 functional limitations (€618/functional limitation improvement) but not for improving QALYs
6 (£24,798/QALY); but the cost is not reflective of savings only to aged care homes but rather to
7 the health and aged care sector combined. The other two studies reported by Simmons et al [24,
8 30] compare ONS with food-based interventions, with conflicting results; both interventions
9 may be considered to have good probability of cost-efficacy (Table 1). The study by Elia et al
10 [28] reported £9857/QALY; however, this reflects the cost of ONS minus cost of the high
11 protein-high energy group; and actual cost/QALY was not reported for either intervention.
12 Certainty in the body of evidence that ONS is cost-effective to improve malnutrition in aged
13 care homes was assessed as very low; primarily due to high risk of bias and heterogeneity
14 across studies (Table 2).

15 Three of the five studies which used food-based interventions reported the direct cost of food-
16 related interventions had a difference of £0.10 to USD\$3.85 (£3.01) per resident per day [24,
17 26, 30]. Lorefalt et al [31] also reported a difference between groups of direct cost of £77.26
18 per year; however, this included staff training as well as additional food items [27]. There was
19 low confidence in the body of evidence that food-based interventions are cost-effective in aged
20 care homes; due to a risk of bias and heterogeneity across studies (Table 2).

21
22

23 *Clinical outcomes*

24 Regarding clinical outcomes, two of the studies [21, 27] used the Mini Nutritional Assessment
25 (MNA) in addition to other measures; however, most did not use validated malnutrition
26 assessments [22-26] (Table 1). Body weight was reported in all of the studies [21-27, 32], and
27 BMI in six. The next most reported outcomes were energy intake (n=4 studies) and physical
28 function (n=3 studies). Some of the studies reported gender differences between malnutrition,
29 however this was not listed in most of the studies.

30 All studies showed significant clinical improvement in the intervention groups; excepting the
31 high protein-high energy advice group which was reported by Elia et. al. 2017 [28] and also
32 included in Parsons et. al. 2017 [32] (results reported across two papers). Seven of the eight

- 1 studies showed increases in weight and six interventions (reported in n=4 studies) reported
- 2 improvements in energy intake compared with control and/or baseline.

Table 1: Study design, characteristics and outcomes of intervention studies with financial outcome data which aim to improve malnutrition in residential aged care

Citation	Setting and population	Study design and economic methods	Intervention and comparator conditions	Summary of findings
Interventions modifying the dining environment				
Edwards and Beck. (2002) [25]	<ul style="list-style-type: none"> USA Mean age 80.1yrs N=62 participants with Alzheimer's Disease Females=61% N=3 clusters (aged care homes) 	<ul style="list-style-type: none"> Cluster non-randomised controlled cross-over trial. Intervention: 8 weeks. Follow-up: 10 weeks. Economic method: Basic economic figures. No analysis. Economic cost vs. benefit/effect measured: Cost =none reported; Benefit/effect = Financial benefit (cost saving) reported for one aged care home related to supplement use. 	<ul style="list-style-type: none"> IG: 8 weeks with fish aquarium in the dining room. IG did not cross-over to CG. CG: 2 weeks with scenic ocean picture introduced to dining room followed by a 2-week washout period (no picture and no aquarium) followed then by 8 weeks with aquarium. 	<ul style="list-style-type: none"> IG: Food intake increased significantly (27.1% increased compared with baseline; p<0.000). Mean weight increase (1.65lbs; p<0.000) compared with baseline. CG: No significant changes observed in food intake or body weight observed. Between groups: not compared. Economic Findings: \$11.44 decreased daily cost of ONS in n=1/3 facilities. Currency unclear; assumed to be USD.
Interventions providing oral nutritional supplementation				
Lee et. al. (2013)[21]	<ul style="list-style-type: none"> Taiwan Mean age 79-80±8yrs N=92. Females=58%. N=1 aged care homes 	<ul style="list-style-type: none"> Double-blind RCT. Intervention: minimum of 12-24 weeks depending on needs of participant. Follow-up: 24 weeks + 12months for mortality Economic method: Cost of Intervention/supplement reported. No analysis. Economic cost versus benefit/effect measured: cost = direct cost of supplement. Benefit/Effect = none included in economic analysis. 	<ul style="list-style-type: none"> IG: If BMI <24 km/m2 and MNA score <24 were provided a 50g/day soy protein-based supplement (9.5g protein, 250kcal, all essential micronutrients) as a warm drink at AT until MNA or BMI improved to >24 and >24kg/m2 respectively + encouragement to consume by staff. CG: Including non-eligible persons for supplement in IG received normal meals including warm soup at AT. 	<ul style="list-style-type: none"> Between groups: Accounting for group allocation and time, at 24 weeks follow-up, IG participants increased body weight (β1.62[95%CI: 0.21-3.03], P<0.05), BMI (β0.57[95%CI: 0.05-1.09], P<0.05), MAC (β0.91[95%CI: 0.40-1.41], P<0.001) and CC (β1.00[95%CI: 0.43-1.80], P<0.001). No improvement in albumin, cholesterol. Mortality not reported. Economic Findings: \$0.40 (£0.24 per resident per day. Analysis by review authors estimates approximately \$2,024 for the cost of supplementation for the entire study period. Assumed dollar is USD.

<p>Neelemaat et. al. (2012) [22]</p>	<ul style="list-style-type: none"> Netherlands Mean age 74.6±9.5yrs. N=210. Female: 55%. N=0 aged care homes sampled. Sample is a hospitalised population; approx. 10% of which were dwelling in an Aged care home 	<ul style="list-style-type: none"> RCT. Intervention: hospital admission period + 3-months post discharge follow-up. Follow-up: 3-months after hospital discharge Economic method: CEA and CUA. Economic cost vs. benefit/effect measured: Cost = Direct costs were supplement costs, telehealth cost, hospital admission costs, specialist visits. Non-direct health costs were included using a diary e.g complementary medicine, informal care, and other indirect costs were absenteeism paid, unpaid labour. Costs were Dutch standard costs. Effect/Benefit: CEA = nutritional status and physical function. CUA: QALY generated by the EQ-D instrument. 	<ul style="list-style-type: none"> IG: In hospital nutrition support: HPHE diet + 2 ONS (330kcal; 12g protein per supplement) + 1 vitamin/mineral supplement (400IE Vit D3 + 500mg Ca/day); post-hospital nutrition support: 2 ONS continued, 1 vitamin/mineral supplement continued + 6 weeks of fortnightly telehealth (6 sessions total) by dietitian until 3/12 post hospital discharge. CG: Usual care with ONS/other supplements only if physician prescribed. No post-hospital support. 	<ul style="list-style-type: none"> IG: Functional limitation change μ-0.24±S.E.0.15; hospital LOS - μ13±16.8; QALYs μ0.15±0.01; physical activities μ0.52±0.17. Significance of change not reported. CG: Functional limitation change μ-0.47±0.15; hospital LOS μ14±12.5; QALYs μ0.13±S.E.0.01; physical activities μ0.42±0.26. Significance of change not reported. Between groups: No significant difference in hospital LOS, QALYs at 3-months follow-up or physical function. IG improved in functional limitations (CG change: μ-0.24±S.E.0.15 vs IG change μ-0.47±0.15; difference -0.72 [95%CI: -1.15 to -0.28; P-value not reported]). Economic Findings: Overall results (not aged care home specific) £24,798/QALY. £4.111/physical activity scale improvement. €618/functional limitation improvement. Probability that intervention is cost-effective for improvement in QALYs and physical activity are low (0.5 and 0.6 respectively). £5978 (below £18395 maximum) investment from Netherlands society, 0.95 probability the intervention is cost effective for improvement in functional limitations.
<p>Simmons et. al. (2010) [23]</p>	<ul style="list-style-type: none"> USA Mean age 86.9±11.3yrs. N=86. Female=62%. N=3 aged care homes 	<ul style="list-style-type: none"> Three-armed RCT. Intervention: 6 weeks Follow-up: 6 weeks Economic method: CEA. Economic cost vs. benefit/effect measured: cost = additional daily food, fluid or supplement spending and salary for staff time for nutritional care delivery. Benefit/Effect = between meal and total daily energy intake. 	<ul style="list-style-type: none"> IG: ONS [not further described] offered twice daily at 10am and 2pm. Second intervention arm reported below. Second IG was food based (see below). CG: no foods or ONS offered, only usual provided food and beverages (not further described). 	<ul style="list-style-type: none"> IG: Compared with baseline, the mean difference of energy intake was -125kCal (P<0.05), Increased energy intake in mid-meals (151kcal; P<0.05) but this caused an overall ↓ in total energy intake. No significant change in body weight. CG: Compared with baseline, the mean difference of energy intake was 5kcal. No significant change in body weight. Between groups: not compared. Economic Findings: Mean difference of direct costs of intervention from baseline to 6-weeks were USD\$2.10 per resident per day for the supplement group and USD\$-0.03 for the control group per resident per day. CEA analysis shows supplement group more likely to result in a decrease in total calories relative to the snack intervention (see below). CEA acceptability curves show snack intervention consistently exceeds supplement intervention for net benefit (e.g. USD\$0.04 value of one-unit caloric gain, probability of net benefit is 65% for supplement group and 80% for snack group).

<p>Simmons et al.(2015) [24]</p>	<ul style="list-style-type: none"> • USA • Mean age 83.1±11.9yrs. • N=175. • Female = 81%. • N=5 aged care homes 	<ul style="list-style-type: none"> • Three-armed RCT • Intervention: 6-months • Follow-up: 6-months • Economic method: CEA. • Economic cost vs. benefit/effect measured: cost = additional daily food, fluid or supplement spending and salary for staff time for nutritional care delivery. Benefits/effects = between meal and total daily energy intake. 	<ul style="list-style-type: none"> • IG ONS [not further described] offered twice daily in the morning and afternoon for five days per week. Second IG was food based (see below). • CG: no foods or ONS offered, only usual provided food and beverages (not further described). 	<ul style="list-style-type: none"> • IG: Average of 1.8kg weight gain, the mean difference of total energy intake was 253kcal (95%CI: 109-397). Mid-meal energy intake increased (151kcal; P<0.05) but this caused an overall decrease in total energy intake. • CG: Average loss of 0.5kg body weight in control group. • Between groups: not compared. • Economic Findings: Mean difference of direct costs of intervention at 6-months compared with the control group was USD\$2.54 per resident per day. Incremental cost-effectiveness ratios 103kcal/USD\$. CEA acceptability curves show supplement intervention consistently exceeds snack intervention (see below) for net benefit (e.g. USD\$0.01 value of one-unit caloric gain, probability of net benefit is 57%.)
<p>Elia et al 2017. [28] Data also reported in Parsons et al. [32]</p>	<ul style="list-style-type: none"> • UK • Mean age 88.8±8yrs. • N=104 (incl 57 aged care home residents) • Female=86%. • N=53 aged care homes 	<ul style="list-style-type: none"> • Two-armed, non-controlled, intervention trial. • Intervention: 12 weeks • Follow-up: 12 weeks • Economic method: CEA • Economic cost vs. benefit/effect measured: cost = direct costs of intervention, unit costs of health care utilisation. Benefits/effects = QALYs adjusted for malnutrition and other factors. 	<ul style="list-style-type: none"> • IG: ONS (1.5-2.4kCal/ml) aiming to increase intake by at least 600kCal/day and 16g protein a day. Saw dietitian at baseline and 6 weeks to receive advice relating to ONS. • CG: none. Compared to 12-week baseline observation period. 	<ul style="list-style-type: none"> • IG: Quality of life (EQ-5D-TTO) decreased (μ change: -0.02) (not tested statistically). Body weight improved (μ change: 1.22±0.45kg; P=0.010). Energy increased (μ change: 286kcal) (not tested statistically). QALY gained μ 0.1302±0.0084. • CG: N/A. • Economic Findings: Direct cost of intervention: £162.30 per resident. Direct unit cost of health care utilisation: £376±34. Significantly higher than HPHE group (see below). Cost/QALY: £9857 (ONS group minus HPHE group; actual cost/QALY not reported for each group).
<p>Interventions providing food-based modifications</p>				
<p>Simmons et. al. (2010) [23]</p>	<ul style="list-style-type: none"> • As per above. 	<ul style="list-style-type: none"> • As per above. 	<ul style="list-style-type: none"> • IG: Variety of snacks (yoghurt, pudding, fruit, juices) offered twice daily at 10am and 2pm. • CG: As per above. 	<ul style="list-style-type: none"> • IG: Compared with baseline, the mean difference of energy intake was 163kcal (P<0.001) for the snack group. No change in body weight. • CG: as per above. • Between groups: not compared. • Economic Findings: Mean difference of direct costs of intervention from baseline to 6-weeks were USD\$2.06 per resident per day for the snack group, and USD\$-0.03 for the control group per resident per day.

<p>Simmons et al.(2015) [24]</p>	<ul style="list-style-type: none"> As per above. 	<ul style="list-style-type: none"> As per above. 	<ul style="list-style-type: none"> IG: Variety of snacks (yoghurt, pudding, juices, liquid supplements) offered twice daily in the morning and afternoon. CG: As per above. 	<ul style="list-style-type: none"> IG: Compared with the control group, the mean difference of total energy intake was 288kcal (95%CI: 144-432). No change in body weight. CG: as per above. Between groups: not compared. Economic Findings: Mean difference of direct costs of intervention at 6-months compared with the control group was USD\$3.85 per resident per day. Incremental cost-effectiveness ratios 79kcal/USD\$ for the snack group. CEA acceptability curves show supplement intervention consistently exceeds snack intervention for net benefit (e.g. USD\$0.01 value of one unit caloric gain, probability of net benefit is 18%).
<p>Elia et al 2017. [28] Data also reported in Parsons et al. [32]</p>	<ul style="list-style-type: none"> As per above. 	<ul style="list-style-type: none"> As per above. 	<ul style="list-style-type: none"> IG: Dietary advice for HPHE snacks and drinks with aid of a diet sheet. Saw dietitian at baseline and 6 weeks to receive advice about HPHE diet. Dietitian discussed plan with care home. CG: none. Compared to 12-week baseline observation period. 	<ul style="list-style-type: none"> IG: Quality of life (EQ-5D-TTO) decreased (μ change: -0.159) (not tested statistically). No change in body weight. kCal decreased (μ change: -93kcal) (not tested statistically). QALY gained μ 0.1128\pm0.0086. CG: N/A. Economic Findings: Direct cost of intervention: not reported. Direct unit cost of health care utilisation: £186\pm38. Significantly lower than ONS group (see above).
<p>Lorefalt et. al. (2011) [27]</p>	<ul style="list-style-type: none"> Sweden Mean age 83-86\pm8-9yrs N=109 Females=50%. N= 6 aged care homes 	<ul style="list-style-type: none"> Non-randomised controlled trial Intervention: 3 months Follow-up: 3 months for clinical data, 1 year for cost data. Economic method: Health care unit cost comparison on direct healthcare costs. Economic cost vs. benefit/effect measured: cost = Cost for each health care contact; Benefit/Effect = none included in economic analysis. 	<ul style="list-style-type: none"> IG: aged care home staff provided with 3x2hr education program by project leader - a nurse with nutrition background. MNA >24 (well nourished) offered snack (e.g. fruit, yoghurt) at midmeals. MNA <24 (risk of malnutrition/ malnourished) had modified food choices within existing food availability and costs: offered appetizer at lunch (e.g. soup, egg, herring), additional snacks (e.g. smoothies, bread and butter, milk and yoghurt) distributed throughout the day 	<ul style="list-style-type: none"> IG: MN prevalence 26% at baseline and 12% at follow-up; body weight change at 3/12 follow-up 2.7\pm3.9kg; BMI at 3/12 follow-up 25.6\pm4.9kg/m². CG: Malnutrition prevalence 18% at baseline and 28% at follow-up; Body weight change at 3/12 follow-up -0.6\pm4.9kg; BMI at 3/12 follow-up 23.7\pm4.9 kg/m². Between groups: Body weight $p=0.0001$; BMI $p=0.05$. Economic Findings: Direct health care cost in IG: median £924, CG: £847 per year. Not compared statistically.

			according to needs and preference. <ul style="list-style-type: none"> • CG: No change to routine meals. 	
Odlund Olin et. al. (2003) [26]	<ul style="list-style-type: none"> • Sweden • Median age 80-83yrs (IQR 71-89) • N=40. • Female: 52%. • N=1 aged care home recruited (N=2 clusters [wards]). 	<ul style="list-style-type: none"> • Non-randomised clustered controlled intervention trial. • Intervention: 15 weeks • Follow-up: 29 weeks post-baseline/27 weeks post intervention commencement. • Economic method: Cost of Intervention. No analysis. • Economic cost vs. benefit/effect measured: cost = Cost of additional butter and cream; Benefit/Effect = none included in economic analysis. 	<ul style="list-style-type: none"> • IG: Served regular hospital diet fortified with butter and cream (2100kCal/day). • CG: Served regular hospital diet (1600kCal/day). 	<ul style="list-style-type: none"> • IG: Compared with baseline, IG increased protein intake (median 48.3 [IQR: 41.8-54.3g] vs median 57.9 [IQR: 46.2-61.2g], P<0.001). ADL remained unchanged. • CG: worsened in ADL during the intervention (median score 15.5 [IQR: 10.0-17.0] increased to 16.0 [IQR: 15.0-18.0], P<0.001). • Between groups: No difference for number of infections. IG increased energy intake (median 1437 [IQR: 1252-1617kcal] vs median 1840 [IQR: 1497-2012kcal], P<0.01). • Economic Findings: £0.10 per resident per day

AT: Afternoon Tea, BMI: Body Mass Index, CC: Calf Circumference, CEA: Cost Effectiveness Analysis, CUA: Cost Utility Analysis, CG: control group, IG: Intervention group, kCal: kilocalorie, kg: kilogram, MAC: Mid Arm Circumference, ONS: Oral Nutrition Supplements, QALY: Quality Adjusted Life Year, RCT: Randomised Control Trial, USD: United States Dollar, vs.: versus, yrs: years.

Table 2: GRADE assessment of the cost-effectiveness environmental, oral nutrition support or food-based malnutrition interventions

OUTCOME: Cost-effectiveness (assessed with: direct cost, cost utility analysis or cost effectiveness analysis)										
Quality assessment							Nº of patients		Quality	Importance
Nº of studies	Study design	Risk of bias	Inconsistency	Indirectness	Imprecision	Other considerations (e.g. publication bias, large effect, plausible confounding and dose response gradient)	Intervention group	Control group (no intervention)		
Question: Environmental changes compared to any other method of nutrition intervention or provision for cost-effectively preventing and/or managing malnutrition in residential aged care										
1	Randomised trials ^a	Very serious ^b	Very serious ^c	Very serious ^d	Very serious ^e	None	45	17	⊕○○○ VERY LOW	IMPORTANT
Question: Supplements compared to usual are for cost-effectively preventing and/or managing malnutrition in residential aged care										
4	Randomised trials	Very serious ^b	Serious ^f	Not serious	Not serious	None	275	218	⊕○○○ VERY LOW	IMPORTANT
Question: Food modifications compared to any other method of nutrition intervention or provision for cost-effectively preventing and/or managing malnutrition in residential aged care										
5	Randomised and non-randomised controlled trials ^g	Serious ^b	Serious ^h	Not serious	Not serious	None	292	258	⊕⊕○○ LOW	IMPORTANT

a. Cluster non-randomised controlled trial

b. Risk of bias assessed by the Cochrane risk of bias tool (Figure 2)

- c. Data relating to cost outcomes were reported for only one of the groups. No data provided for consistency within study groups (three groups received intervention in cross-over design). Could not be compared to any other studies.
- d. Cost-saving data was reported for the decreased spend of supplements; which is not a direct measure of the environmental intervention. There is no confidence that the cost-related outcome is due to the intervention.
- e. No measure of variability reported.
- f. Some inconsistency between the two most clinically homogenous studies (Simmons 2010 and 2015). Combined studies all show cost of ONS is low and results in clinical benefit.
- g. Three studies were non-randomised controlled trials; two were RCTs.
- h. Interventions included significant clinical heterogeneity; however, cost-related results were reported differently between studies making comparison of consistency difficult.

Discussion

There is good evidence that malnutrition places a significant financial burden on our healthcare system [33-35] as well as good evidence that supplements and other nutrition interventions improve intake and nutritional status [36, 37]. This review, however, revealed there is a lack of confidence in the body of economic evidence that introducing malnutrition interventions, whether they be environmental, supplements or food-based, are cost-effective in residential aged care. This lack of confidence is due to the small number and poor-quality of studies economically evaluating nutritional interventions in aged care; particularly for environmental interventions. Despite this, the review showed that included interventions had a low direct cost of implementation (less than £2.30 per resident per day) and provided clinical improvement in patients. Supplements and food-based interventions further demonstrated a low cost per QALY or unit of physical function improvement.

There is great variation in the scope of economic reviews on the topic of malnutrition. One large nutrition and health economics review looked at malnutrition across all ages and settings and concluded nutrition to be a powerful force improving both the health and economic status of society [16]. However, in agreement with the current review, the study found large variations in the approach to economic modelling of malnutrition interventions, and highlighted the need for a well-defined framework for economic analysis on nutrition interventions [16].

Although this current review found insufficient evidence supporting the cost-effectiveness of malnutrition interventions in aged care homes, evidence in the acute setting is stronger as evaluated by three recent systematic reviews [13, 37, 38]. Mitchell et al [13] in a systematic review concluded that malnutrition interventions in the hospital setting showed positive cost-effectiveness for improving outcomes, informed by intervention studies from 2003 to 2013. Although Mitchell et al. only identified three studies for inclusion, they were comprehensive and of a high quality. In 2017, the systematic review by Muscatoli et al [38] found that there was insufficient evidence as to whether supplements significantly reduced hospital readmissions when given to malnourished hospitalised patients and outpatients. However, Muscatoli [38] found supplements resulted in cost savings with a return of investment of \$52.63 in net savings for every dollar spend on supplements in terms of reduced episode cost amongst young patients. The systematic review by Elia et al [37] also examined the cost-effectiveness of using supplements in hospitals, and subgroup analysis found supplements to be cost-effective with a mean net cost saving of £746 per patient. In this same review, the mean

cost saving across 12 of the 14 cost analysis studies comparing supplements with routine care found 12.2% mean cost saving with supplements use [37]. Further hospital-based economic modelling by Banks et al [39] showed cost-effective reduction in risk of developing pressure ulcers with the use of nutritional intervention (including costs of additional food, supplements and additional nutrition/nursing support staff time). This strong and consistent evidence in support of nutrition interventions to cost-effectively improve malnutrition in the acute care setting suggests that similar conclusions may be found in the aged care setting once further well-conducted studies including economic data are undertaken.

Limitations and implications for future research

This systematic literature review focussed on interventional studies only, as these studies provide a higher quality of evidence to evaluate the research question. However, it is acknowledged that excluding observational studies may limit potential learnings [40], particularly regarding external validity. All but two included studies did not sufficiently evaluate the impact of interventions on malnutrition, and none used malnutrition in the cost-utility analysis. Instead, the outcomes of weight, BMI, energy intake and physical function were most frequently used. Although these are important components of malnutrition assessment, they do not reflect malnutrition risk or status alone.

Future research on cost-effectiveness of nutrition-related interventions in the aged care setting need to accurately measure malnutrition, clearly describe interventions and economic methods and provide a detailed description of research design. Rigorous intervention and economic study designs, such as RCTs and cost-utility analyses in future malnutrition studies in the aged care setting may further strengthen and increase confidence in the cost-effective treatment of malnutrition. Although research has demonstrated nutrition interventions are low risk and effective in improving clinical outcomes, stronger evidence regarding cost-effectiveness will support aged care funders and governance to select the most cost-effective treatment options.

Conclusion interventions

Malnutrition places significant economic burden upon the aged care sector and nutrition may be a powerful force for improving both the health and economic status of aged care homes. While there is good evidence that nutrition improves clinical outcomes, the limited and poor-quality studies including economic data in this review indicate evidence of cost-effectiveness in the aged care setting is still limited. This systematic review suggests that supplements and food-based nutrition interventions in the aged care setting have a low cost of implementation, low risk of harm, and may be cost-effective. More studies using well-defined frameworks for

economic analysis, stronger study designs such as double-blinded RCTs, improved quality (reduced risk of bias), along with validated malnutrition measures are needed.

References -

1. Milte, R.K., et al., *Economic evaluation for protein and energy supplementation in adults: opportunities to strengthen the evidence*. European Journal of Clinical Nutrition, 2013. **67**(12): p. 1243-1250.
2. Commission, P., *Economic implications of an ageing Australia*. Productivity Commission, Government of Australia Research Reports, 2005.
3. Ratcliffe, J., et al., *Health economics and geriatrics: challenges and opportunities*, in *Geriatrics*. 2012, InTech.
4. Marshall, S., *Protein-energy malnutrition in the rehabilitation setting: Evidence to improve identification*. Maturitas, 2016. **86**: p. 77-85.
5. Meijers, J.M.M., et al., *Estimating the costs associated with malnutrition in Dutch nursing homes*. Clinical Nutrition, 2012. **31**(1): p. 65-68.
6. Freijer, K., et al., *The economic costs of disease related malnutrition*. Clinical Nutrition, 2013. **32**(1): p. 136-141.
7. Russell, C.A., *The impact of malnutrition on healthcare costs and economic considerations for the use of oral nutritional supplements*. Clinical Nutrition Supplements, 2007. **2**(1): p. 25-32.
8. Rice, N. and C. Normand, *The cost associated with disease-related malnutrition in Ireland*. Public Health Nutrition, 2012. **15**(10): p. 1966-1972.
9. Strathmann, S., et al., *Institutional factors associated with the nutritional status of residents from 10 german nursing homes (ernstes study)*. The Journal of Nutrition, Health & Aging, 2013. **17**(3): p. 271-6.
10. Hugo, C., et al., *What does it cost to feed aged care residents in Australia?* Nutrition & Dietetics, 2017.
11. Agarwal, E., et al., *Optimising nutrition in residential aged care: A narrative review*. Maturitas, 2016. **92**: p. 70-78.
12. *Cost saving guidance*. 2016, National Institute for Health and Clinical Excellence (NICE): <https://www.nice.org.uk/about/what-we-do/into-practice/cost-saving-guidance>.
13. Mitchell, H. and J. Porter, *The cost-effectiveness of identifying and treating malnutrition in hospitals: a systematic review*. Journal of Human Nutrition and Dietetics, 2016. **29**(2): p. 156-164.
14. Weekes, C., et al., *A review of the evidence for the impact of improving nutritional care on nutritional and clinical outcomes and cost*. Journal of human nutrition and dietetics, 2009. **22**(4): p. 324-335.
15. Walzer, S., et al., *Health economics evidence for medical nutrition: are these interventions value for money in integrated care?* ClinicoEconomics and outcomes research: CEOR, 2014. **6**: p. 241.
16. Gyles, C.L., et al., *Health economics and nutrition: a review of published evidence*. Nutrition Reviews, 2012. **70**(12): p. 693-708.

17. Liberati, A., et al., *The PRISMA statement for reporting systematic reviews and meta-analyses of studies that evaluate healthcare interventions: explanation and elaboration*. BMJ : British Medical Journal, 2009. **339**.
18. Welfare, A.I.o.H.a., *25 years of health expenditure in Australia 1989-90 to 2013-14*. 2015(HEALTH AND WELFARE EXPENDITURE SERIES NO. 56).
19. Oxford, U.o., *Critical Appraisal Tools - Centre for Evidence-Based Medicine*. . Website.
20. Guyatt, G.H., et al., *GRADE: an emerging consensus on rating quality of evidence and strength of recommendations*. BMJ, 2008. **336**(7650): p. 924.
21. Lee, L.-C., et al., *Need-based intervention is an effective strategy for improving the nutritional status of older people living in a nursing home: A randomized controlled trial*. International Journal of Nursing Studies, 2013. **50**(12): p. 1580-1588.
22. Neelemaat, F., et al., *Oral nutritional support in malnourished elderly decreases functional limitations with no extra costs*. Clinical Nutrition, 2012. **31**(2): p. 183-190.
23. Simmons, S., X. Zhuo, and E. Keeler, *Cost- effectiveness of nutrition interventions in nursing home residents: A pilot intervention*. The journal of nutrition, health & aging, 2010. **14**(5): p. 367-372.
24. Simmons, S.F., et al., *Cost - Effectiveness of Nutrition Intervention in Long - Term Care*. Journal of the American Geriatrics Society, 2015. **63**(11): p. 2308-2316.
25. Edwards, N.E. and A.M. Beck, *Animal-Assisted Therapy and Nutrition in Alzheimer's Disease*. Western Journal of Nursing Research, 2002. **24**(6): p. 697-712.
26. Ödlund Olin, A., et al., *Energy-dense meals improve energy intake in elderly residents in a nursing home*. Clinical Nutrition, 2003. **22**(2): p. 125-131.
27. Lorefält, B., et al., *Nutritional status and health care costs for the elderly living in municipal residential homes — An intervention study*. The journal of nutrition, health & aging, 2011. **15**(2): p. 92-97.
28. Elia, M., et al., *Cost-effectiveness of oral nutritional supplements in older malnourished care home residents*. Clinical Nutrition, 2017.
29. Lee, L.C., et al., *Need-based intervention is an effective strategy for improving the nutritional status of older people living in a nursing home: a randomized controlled trial*. Int J Nurs Stud, 2013. **50**(12): p. 1580-8.
30. Simmons, S., X. Zhuo, and E. Keeler, *Cost- effectiveness of nutrition interventions in nursing home residents: A pilot intervention*. The journal of nutrition, health & aging, 2010. **14**(5): p. 367-372.
31. Lorefält, B., et al., *Nutritional status and health care costs for the elderly living in municipal residential homes - An intervention study*. Journal of Nutrition, Health and Aging, 2011. **15**(2): p. 92-97.
32. Parsons, E.L., et al., *Oral nutritional supplements in a randomised trial are more effective than dietary advice at improving quality of life in malnourished care home residents*. Clinical Nutrition, 2017. **36**(1): p. 134-142.
33. Elia, M., *The cost of malnutrition in England and potential cost savings from nutritional interventions (full report)*. BAPEN report, 2015.
34. Snider, J.T., et al., *Economic Burden of Community-Based Disease-Associated Malnutrition in the United States*. Journal of Parenteral and Enteral Nutrition, 2014. **38**(2 suppl): p. 77S-85S.
35. Meijers, J.M.M., et al., *Estimating the costs associated with malnutrition in Dutch nursing homes*. Clinical Nutrition, 2011.
36. Baldwin C, K.K., Gibbs M, Weekes C, *Supportive interventions for enhancing dietary intake in malnourished or nutritionally at-risk adults*. Cochrane Database of Systematic Reviews, 2016.
37. Elia, M., et al., *A systematic review of the cost and cost effectiveness of using standard oral nutritional supplements in the hospital setting*. Clinical Nutrition, 2016. **35**(2): p. 370-380.
38. Muscaritoli, M., et al., *Effectiveness and efficacy of nutritional therapy: A systematic review following Cochrane methodology*. Clinical Nutrition, 2017. **36**(4): p. 939-957.

39. Banks, M.D., et al., *Cost effectiveness of nutrition support in the prevention of pressure ulcer in hospitals*. European Journal of Clinical Nutrition, 2013. **67**(1): p. 42-6.
40. Kovesdy, C.P. and K. Kalantar-Zadeh, *Observational studies versus randomized controlled trials: avenues to causal inference in nephrology*. Advances in chronic kidney disease, 2012. **19**(1): p. 11-18.

ONLINE SUPPLEMENTARY MATERIAL 1 – Search Strategy

This table has been provided by the authors to give readers additional information about this work.

Database	Search terms
Medline	1. "nursing home*" [keyword] OR "aged care" [keyword] OR "residential care" [keyword] OR "care home*" [keyword] OR "residential facilit*" [keyword] OR "assisted living" [keyword]
	2. nursing home patient [exp][Emtree term] OR senior center [exp] [Emtree term] OR home for the aged [exp] [Emtree term] OR elderly care [exp] [Emtree term]) OR institutional care [exp] [Emtree term] OR residential care [exp] [emtree term] OR long term care [exp] [Emtree term]
	3. "malnutrition*" [keyword] OR "nutrition*" [keyword] OR "undernutrition" [keyword] OR "under nutrition" [keyword] OR "emaciation" [keyword] OR "undernourish*" [keyword] OR "under nourish*" [keyword] OR "malnourish*" [keyword]
	4. Nutrition [exp] [Emtree term] OR malnutrition [exp] [Emtree term] OR protein calorie <i>malnutrition</i> [exp] [Emtree term]) OR nutritional status [exp] [Emtree term]
	5. "Cost Benefit" [keyword] OR "Economic Benefit" [keyword] OR "Cost Savings" [keyword] OR "Cost Effectiveness" [keyword] OR "healthcare cost" [keyword] OR "health care cost" [keyword] OR "Economic" [keyword] OR "Financial" [keyword]
	6. <i>cost benefit analysis</i> [exp] [Emtree term]) OR <i>economic evaluation</i> [exp] [Emtree term] OR <i>cost effectiveness analysis</i> [exp] [Emtree term])
	(1 OR 2) AND (3 OR 4) AND 6
PubMed & Cochrane	1. nursing home* [keyword] OR "aged care" [keyword] OR "residential care" [keyword] OR care home* [keyword] OR residential facilit* [keyword] OR "assisted living" [keyword]
	2. residential facilities [MeSH term] OR assisted living facility [MeSH term] OR group homes [MeSH term] OR homes for the aged [MeSH term] OR nursing homes [MeSH term] OR health services for the aged [MeSH term]
	3. malnutrition* [keyword] OR nutrition* [keyword] OR undernutrition [keyword] OR "under nutrition" [keyword] OR emaciation [keyword] OR undernourish* [keyword] OR under nourish* [keyword] OR malnourish* [keyword]
	4. Diet, Food and Nutrition [MeSH] OR <i>protein energy malnutrition</i> [MeSH term] OR <i>malnutrition</i> [MeSH term] OR <i>nutritional status</i> [MeSH term] OR <i>undernutrition</i> [MeSH term] OR <i>nutritional deficiency</i> [MeSH term] OR <i>protein calorie malnutrition</i> [MeSH term] OR

	<p><i>emaciation</i> [MeSH term] OR <i>nutrition status</i> [MeSH term] OR <i>protein deficiency</i> [MeSH term]</p>
	<p>5. Cost Benefit [keyword] OR Economic Benefit [keyword] OR Cost Savings [keyword] OR Cost Effectiveness [keyword] OR healthcare cost [keyword] OR health care cost [keyword] OR Economic [keyword] OR Financial [keyword]</p>
	<p>6. (1 OR 2) AND (3 OR 4) AND 5</p>
Embase using Emtree terms	<p>1. "nursing home*" [keyword] OR "aged care" [keyword] OR "residential care" [keyword] OR "care home*" [keyword] OR "residential facilit*" [keyword] OR "assisted living" [keyword]</p>
	<p>2. nursing home patient [exp][Emtree term] OR senior center [exp] [Emtree term] OR home for the aged [exp] [Emtree term] OR elderly care [exp] [Emtree term]) OR institutional care [exp] [Emtree term] OR residential care [exp] [emtree term] OR long term care [exp] [Emtree term]</p>
	<p>3. "malnutrition*" [keyword] OR "nutrition*" [keyword] OR "undernutrition" [keyword] OR "under nutrition" [keyword] OR "emaciation" [keyword] OR "undernourish*" [keyword] OR "under nourish*" [keyword] OR "malnourish*" [keyword]</p>
	<p>4. Nutrition [exp] [Emtree term] OR malnutrition [exp] [Emtree term] OR protein calorie <i>malnutrition</i> [exp] [Emtree term]) OR nutritional status [exp] [Emtree term]</p>
	<p>5. "Cost Benefit" [keyword] OR "Economic Benefit" [keyword] OR "Cost Savings" [keyword] OR "Cost Effectiveness" [keyword] OR "healthcare cost" [keyword] OR "health care cost" [keyword] OR "Economic" [keyword] OR "Financial" [keyword]</p>
	<p>6. <i>cost benefit analysis</i> [exp] [Emtree term]) OR <i>economic evaluation</i> [exp] [Emtree term] OR <i>cost effectiveness analysis</i> [exp] [Emtree term])</p>
	<p>7. (1 OR 2) AND (3 OR 4) AND 6</p>
Econlit	<p>1. ("aged care" [keyword] OR "nursing home*" [keyword] OR "residential care" [keyword] OR "care home*" [keyword] OR "residential facilit*" [keyword] OR "assisted living" [keyword]) OR ("elderly" [keyword] OR "older adults" [keyword] OR "residents" [keyword] OR "aged" [keyword]) AND</p>
	<p>1. "malnutrition*" [keyword] OR "nutrition*" [keyword] OR "undernutrition" [keyword] OR "under nutrition" [keyword] OR "emaciation" [keyword] OR "undernourish*" [keyword] OR "under nourish*" [keyword] OR "malnourish*" [keyword] AND</p>

	<p>3. <i>“Cost Benefit”</i> [keyword] OR <i>“Economic Benefit”</i> [keyword] OR <i>“Cost Savings”</i> [keyword] OR <i>“Cost Effectiveness”</i> [keyword] OR <i>“healthcare cost”</i> [keyword] OR <i>“health care cost”</i> [keyword] OR <i>“Economic”</i> [keyword] OR <i>“Financial”</i> [keyword]</p>
	<p>4. 1 AND 2 AND 3</p>
Web of Science	<p>1. (aged care [keyword] OR aged care facility [keyword] OR nursing home** [keyword] OR “residential care” [keyword] OR “care home*” [keyword] OR “residential facilit*” [keyword] OR “assisted living” [keyword]) OR (“elderly” [keyword] OR “older adults” [keyword] OR “residents” [keyword] OR “aged” [keyword])</p>
	<p>2. “malnutrition*” [keyword] OR “nutrition*” [keyword] OR “undernutrition” [keyword] OR “under nutrition” [keyword] OR “emaciation” [keyword] OR “undernourish*” [keyword] OR “under nourish*” [keyword] OR “malnourish*” [keyword] AND</p>
	<p>3. <i>“Cost Benefit”</i> [keyword] OR <i>“Economic Benefit”</i> [keyword] OR <i>“Cost Savings”</i> [keyword] OR <i>“Cost Effectiveness”</i> [keyword] OR <i>“healthcare cost”</i> [keyword] OR <i>“health care cost”</i> [keyword] OR <i>“Economic”</i> [keyword] OR <i>“Financial”</i> [keyword]</p>
	<p>4. 1 AND 2 AND 3</p>
CINAHL (via EBSCO HOST)	<p>1. “nursing home*” [keyword] OR “aged care” [keyword] OR “residential care” [keyword] OR care home* [keyword] OR residential facilit* [keyword] OR “assisted living” [keyword]</p>
	<p>2. Nursing Homes [exp] [CINAHL heading] OR Gerontologic Nursing [exp] [CINAHL heading] OR Gerontologic Care [exp] [CINAHL heading] OR Housing for the Elderly [exp] [CINAHL heading] OR Nursing Home Patients [exp] [CINAHL heading] OR Residential Care [exp] [CINAHL heading]</p>
	<p>3. “malnutrition*” [keyword] OR “nutrition*” [keyword] OR “undernutrition” [keyword] OR “under nutrition” [keyword] OR “emaciation” [keyword] OR “undernourish*” [keyword] OR “under nourish*” [keyword] OR “malnourish*” [keyword]</p>
	<p>4. Nutrition [exp] [CINAHL heading] OR Nutrition Services [exp] [CINAHL heading] OR Malnutrition [exp] [CINAHL heading] OR Protein-Energy Malnutrition [exp] [CINAHL heading] OR Nutritional Status [exp] [CINAHL heading]</p>
	<p>5. <i>“Cost Benefit”</i> [keyword] OR <i>“Economic Benefit”</i> [keyword] OR <i>“Cost</i></p>

	<p><i>Savings</i> [keyword] OR <i>Cost Effectiveness</i> [keyword] OR <i>healthcare cost</i> [keyword] OR <i>health care cost</i> [keyword] OR <i>Economic</i> [keyword] OR <i>Financial</i> [keyword]</p>
	<p>6. <i>Cost Savings</i> [CINAHL heading] OR <i>Cost and Cost Analysis</i> [CINAHL heading] OR <i>Health Care Costs</i> [CINAHL heading] OR <i>Cost Benefit Analysis</i> [CINAHL heading]</p>
	<p>7. (1 OR 2) AND (3 OR 4) AND (5 OR 6)</p>
<p>NHS Economic Evaluation Database</p>	<p>1. "nursing home*" [keyword] OR "aged care" [keyword] OR "residential care" [keyword] OR "care home*" [keyword] OR "residential facilit*" [keyword] OR "assisted living" [keyword]</p>
	<p>2. residential facilities [MeSH term] OR assisted living facility [MeSH term] OR group homes [MeSH term] OR homes for the aged [MeSH term] OR nursing homes [MeSH term] OR health services for the aged [MeSH term]</p>
	<p>3. malnutrition* [keyword] OR nutrition* [keyword] OR undernutrition [keyword] OR "under nutrition" [keyword] OR emaciation [keyword] OR undernourish* [keyword] OR under nourish* [keyword] OR malnourish* [keyword]</p>
	<p>4. Diet, Food and Nutrition [MeSH] OR <i>protein energy malnutrition</i> [MeSH term] OR <i>malnutrition</i> [MeSH term] OR <i>nutritional status</i> [MeSH term] OR <i>undernutrition</i> [MeSH term] OR <i>nutritional deficiency</i> [MeSH term] OR <i>protein calorie malnutrition</i> [MeSH term] OR <i>emaciation</i> [MeSH term] OR <i>nutrition status</i> [MeSH term] OR <i>protein deficiency</i> [MeSH term]</p>
	<p>5. Cost Benefit [keyword] OR Economic Benefit [keyword] OR Cost Savings [keyword] OR Cost Effectiveness [keyword] OR healthcare cost [keyword] OR health care cost [keyword] OR Economic [keyword] OR Financial [keyword]</p>
	<p>6. (1 OR 2) AND (3 OR 4) AND 5</p>

Online Supplementary Material 2

Supplementary Table 1: Study design, characteristics and outcomes of intervention studies with financial outcome data which aim to improve malnutrition in residential aged care

Citation	Setting and population	Study design and economic methods	Study purpose	Intervention and comparator conditions	Summary of clinical findings	Summary of economic findings
Interventions modifying the dining environment						
Edwards and Beck. (2002) [26]	<ul style="list-style-type: none"> USA Mean age 80.1yrs N=62 participants with Alzheimer's Disease Females=61% N=3 clusters (aged care homes) 	<ul style="list-style-type: none"> Cluster non-randomised controlled cross-over trial. Intervention: 8 weeks. Follow-up: 10 weeks. Economic method: Basic economic figures. No analysis. Economic cost vs. benefit/effect measured: Cost =none reported; Benefit/effect = Financial benefit (cost saving) reported for one aged care home related to supplement use. 	Examining the influence of animal assisted therapy, specifically fish aquariums, on nutritional intake in individuals with Alzheimer's Disease.	<ul style="list-style-type: none"> IG: 8 weeks with fish aquarium in the dining room. IG did not cross-over to CG. CG: 2 weeks with scenic ocean picture introduced to dining room followed by a 2-week washout period (no picture and no aquarium) followed then by 8 weeks with aquarium. 	<ul style="list-style-type: none"> IG: Food intake increased significantly (27.1% increase compared with baseline; $p<0.000$). Mean weight increase (1.65lbs; $p<0.000$) compared with baseline. CG: No significant changes observed in food intake or body weight observed. Between groups: not compared. 	\$11.44 decreased daily cost of ONS in n=1/3 facilities. Currency unclear; assumed to be USD.
Interventions providing oral nutritional supplementation						
Lee et. al. (2013)[22]	<ul style="list-style-type: none"> Taiwan Mean age 79-80±8yrs N=92. Females=58%. N=1 aged care homes 	<ul style="list-style-type: none"> Double-blind RCT. Intervention: minimum of 12-24 weeks depending on needs of participant. Follow-up: 24 weeks + 12months for mortality Economic method: Cost of Intervention/supplement reported. No analysis. Economic cost versus benefit/effect measured: cost = direct cost of 	Examining the effectiveness of routine screening and nutrition supplementation in improving the nutritional status of persons living in NH's.	<ul style="list-style-type: none"> IG: If BMI <24 kg/m² and MNA score <24 were provided a 50g/day soy protein-based supplement (9.5g protein, 250kcal, all essential micronutrients) as a warm drink at AT until MNA or BMI improved to >24 and >24kg/m² respectively + encouragement to 	<ul style="list-style-type: none"> Between groups: Accounting for group allocation and time, at 24 weeks follow-up, IG participants increased body weight (β1.62[95%CI: 0.21-3.03], $P<0.05$), BMI (β0.57[95%CI: 0.05-1.09], $P<0.05$), MAC (β0.91[95%CI: 0.40-141], $P<0.001$) and CC 	\$0.40 (£0.24 per resident per day. Analysis by review authors estimates approximately \$2,024 for the cost of supplementation for the entire study period. Assumed dollar is USD.

		supplement. Benefit/Effect = none included in economic analysis.		consume by staff. <ul style="list-style-type: none"> CG: Including non-eligible persons for supplement in IG received normal meals including warm soup at AT. 	(β 1.00[95%CI: 0.43-1.80], $P < 0.001$). No improvement in albumin, cholesterol. Mortality not reported.	
Neelemaat et. al. (2012) [23]	<ul style="list-style-type: none"> Netherlands Mean age 74.6\pm9.5yrs. N=210. Female: 55%. N=0 aged care homes sampled. Sample is a hospitalised population; approx. 10% of which were dwelling in an Aged care home 	<ul style="list-style-type: none"> RCT. Intervention: hospital admission period + 3-months post discharge follow-up. Follow-up: 3-months after hospital discharge Economic method: CEA and CUA. Economic cost vs. benefit/effect measured: Cost = Direct costs were supplement costs, telehealth cost, hospital admission costs, specialist visits. Non-direct health costs were included using a diary e.g complementary medicine, informal care, and other indirect costs were absenteeism paid, unpaid labour. Costs were Dutch standard costs. Effect/Benefit: CEA = nutritional status and physical function. CUA: QALY generated by the EQ-D instrument. 	Cost-effectiveness of ONS in the community following hospital discharge.	<ul style="list-style-type: none"> IG: In hospital nutrition support: HPHE diet + 2 ONS (330kcal; 12g protein per supplement) + 1 vitamin/mineral supplement (400IE Vit D3 + 500mg Ca/day); post-hospital nutrition support: 2 ONS continued, 1 vitamin/mineral supplement continued + 6 weeks of fortnightly telehealth (6 sessions total) by dietitian until 3/12 post hospital discharge. CG: Usual care with ONS/other supplements only if physician prescribed. No post-hospital support. 	<ul style="list-style-type: none"> IG: Functional limitation change μ-0.24\pmS.E.0.15; hospital LOS - μ13\pm16.8; QALYs μ0.15\pm0.01; physical activities μ0.52\pm0.17. Significance of change not reported. CG: Functional limitation change μ-0.47\pm0.15; hospital LOS μ14\pm12.5; QALYs μ0.13\pmS.E.0.01; physical activities μ0.42\pm0.26. Significance of change not reported. Between groups: No significant difference in hospital LOS, QALYs at 3-months follow-up or physical function. IG improved in functional limitations (CG change: μ-0.24\pmS.E.0.15 vs IG change μ-0.47\pm0.15; difference -0.72 [95%CI: -1.15 to -0.28; P-value not reported]). 	Overall results (not aged care home specific) £24,798/QALY. £4.111/physical activity scale improvement. €618/functional limitation improvement. Probability that intervention is cost-effective for improvement in QALYs and physical activity are low (0.5 and 0.6 respectively). £5978 (below £18395 maximum) investment from Netherlands society, 0.95 probability the intervention is cost effective for improvement in functional limitations.

<p>Simmons et. al. (2010) [24]</p>	<ul style="list-style-type: none"> • USA • Mean age 86.9±11.3yrs. • N=86. • Female=62%. • N=3 aged care homes 	<ul style="list-style-type: none"> • Three-armed RCT. • Intervention: 6 weeks • Follow-up: 6 weeks • Economic method: CEA. • Economic cost vs. benefit/effect measured: cost = additional daily food, fluid or supplement spending and salary for staff time for nutritional care delivery. Benefit/Effect = between meal and total daily energy intake. 	<p>Aim was to determine the cost-effectiveness of supplements or snacks foods between meals to increase caloric intake compared to usual care.</p>	<ul style="list-style-type: none"> • IG: ONS [not further described] offered twice daily at 10am and 2pm. Second intervention arm reported below. Second IG was food based (see below). • CG: no foods or ONS offered, only usual provided food and beverages (not further described). 	<ul style="list-style-type: none"> • IG: Compared with baseline, the mean difference of energy intake was -125kCal (P<0.05), Increased energy intake in mid-meals (151kcal; P<0.05) but this caused an overall ↓ in total energy intake. No significant change in body weight. • CG: Compared with baseline, the mean difference of energy intake was 5kcal. No significant change in body weight. • Between groups: not compared. 	<p>Mean difference of direct costs of intervention from baseline to 6-weeks were USD\$2.10 per resident per day for the supplement group and USD\$-0.03 for the control group per resident per day. CEA analysis shows supplement group more likely to result in a decrease in total calories relative to the snack intervention (see below). CEA acceptability curves show snack intervention consistently exceeds supplement intervention for net benefit (e.g. USD\$0.04 value of one-unit caloric gain, probability of net benefit is 65% for supplement group and 80% for snack group).</p>
------------------------------------	------------------------------------------------------------------------------------------------------------------------------------------------------------------	-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------	----------------------------------------------------------------------------------------------------------------------------------------------------	------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------	------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------	--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------

Simmons et. al.(2015) [25]	<ul style="list-style-type: none"> USA Mean age 83.1±11.9yrs. N=175. Female = 81%. N=5 aged care homes 	<ul style="list-style-type: none"> Three-armed RCT Intervention: 6-months Follow-up: 6-months Economic method: CEA. Economic cost vs. benefit/effect measured: cost = additional daily food, fluid or supplement spending and salary for staff time for nutritional care delivery. Benefits/effects = between meal and total daily energy intake. 	Aim was to determine the cost-effectiveness of supplements or snacks foods between meals to increase caloric intake compared to usual care.	<ul style="list-style-type: none"> IG ONS [not further described] offered twice daily in the morning and afternoon for five days per week. Second IG was food based (see below). CG: no foods or ONS offered, only usual provided food and beverages (not further described). 	<ul style="list-style-type: none"> IG: Average of 1.8kg weight gain, the mean difference of total energy intake was 253kcal (95%CI: 109-397). Mid-meal energy intake increased (151kcal; P<0.05) but this caused an overall decrease in total energy intake. CG: Average loss of 0.5kg body weight in control group. Between groups: not compared. 	Mean difference of direct costs of intervention at 6-months compared with the control group was USD\$2.54 per resident per day. Incremental cost-effectiveness ratios 103kcal/USD\$. CEA acceptability curves show supplement intervention consistently exceeds snack intervention (see below) for net benefit (e.g. USD\$0.01 value of one-unit caloric gain, probability of net benefit is 57%.)
Elia et al 2017. [29] Data also reported in Parsons et al. [32]	<ul style="list-style-type: none"> UK Mean age 88.8±8yrs. N=104 (incl 57 aged care home residents) Female=86%. N=53 aged care homes 	<ul style="list-style-type: none"> Two-armed, non-controlled, intervention trial. Intervention: 12 weeks Follow-up: 12 weeks Economic method: CEA Economic cost vs. benefit/effect measured: cost = direct costs of intervention, unit costs of health care utilisation. Benefits/effects = QALYs adjusted for malnutrition and other factors. 	Aim was to complete a cost-utility analysis to specifically examine whether the use of ONS in care home residents, with a wide variety of diseases and clinical conditions is cost effective relative to dietary advice.	<ul style="list-style-type: none"> IG: ONS (1.5-2.4kCal/ml) aiming to increase intake by at least 600kCal/day and 16g protein a day. Saw dietitian at baseline and 6 weeks to receive advice relating to ONS. CG: none. Compared to 12-week baseline observation period. 	<ul style="list-style-type: none"> IG: Quality of life (EQ-5D-TTO) decreased (μ change: -0.02) (not tested statistically). Body weight improved (μ change: 1.22±0.45kg; P=0.010). Energy increased (μ change: 286kcal) (not tested statistically). QALY gained μ 0.1302±0.0084. CG: N/A. 	Direct cost of intervention: £162.30 per resident. Direct unit cost of health care utilisation: £376±34. Significantly higher than HPHE group (see below). Cost/QALY: £9857 (ONS group minus HPHE group; actual cost/QALYnot reported for each group).
Interventions providing food-based modifications						
Simmons et. al. (2010) [24]	<ul style="list-style-type: none"> As per above. 	<ul style="list-style-type: none"> As per above. 	<ul style="list-style-type: none"> As per above. 	<ul style="list-style-type: none"> IG: Variety of snacks (yoghurt, pudding, fruit, juices) offered twice daily 	<ul style="list-style-type: none"> IG: Compared with baseline, the mean difference of energy 	Mean difference of direct costs of intervention from baseline to 6-weeks were

				<ul style="list-style-type: none"> at 10am and 2pm. CG: As per above. 	<ul style="list-style-type: none"> intake was 163kcal (P<0.001) for the snack group. No change in body weight. CG: as per above. Between groups: not compared. 	<ul style="list-style-type: none"> USD\$2.06 per resident per day for the snack group, and USD\$-0.03 for the control group per resident per day.
<ul style="list-style-type: none"> Simmons et al.(2015) [25] 	<ul style="list-style-type: none"> As per above. 	<ul style="list-style-type: none"> As per above. 	<ul style="list-style-type: none"> As per above. 	<ul style="list-style-type: none"> IG: Variety of snacks (yoghurt, pudding, juices, liquid supplements) offered twice daily in the morning and afternoon. CG: As per above. 	<ul style="list-style-type: none"> IG: Compared with the control group, the mean difference of total energy intake was 288kcal (95%CI: 144-432). No change in body weight. CG: as per above. Between groups: not compared. 	<ul style="list-style-type: none"> Mean difference of direct costs of intervention at 6-months compared with the control group was USD\$3.85 per resident per day. Incremental cost-effectiveness ratios 79kcal/USD\$ for the snack group. CEA acceptability curves show supplement intervention consistently exceeds snack intervention for net benefit (e.g. USD\$0.01 value of one unit caloric gain, probability of net benefit is 18%).
<ul style="list-style-type: none"> Elia et al 2017. [29] Data also reported in Parsons et al. [32] 	<ul style="list-style-type: none"> As per above. 	<ul style="list-style-type: none"> As per above. 	<ul style="list-style-type: none"> As per above. 	<ul style="list-style-type: none"> IG: Dietary advice for HPHE snacks and drinks with aid of a diet sheet. Saw dietitian at baseline and 6 weeks to receive advice about HPHE diet. Dietitian discussed plan with care home. CG: none. Compared to 12-week baseline observation period. 	<ul style="list-style-type: none"> IG: Quality of life (EQ-5D-TTO) decreased (μ change: -0.159) (not tested statistically). No change in body weight. kCal decreased (μ change: -93kcal) (not tested statistically). QALY gained μ 0.1128\pm0.0086. CG: N/A. 	<ul style="list-style-type: none"> Direct cost of intervention: not reported. Direct unit cost of health care utilisation: £186\pm38. Significantly lower than ONS group (see above).
<ul style="list-style-type: none"> Lorefalt et. al. (2011) [28] 	<ul style="list-style-type: none"> Sweden Mean age 83-86\pm8-9yrs 	<ul style="list-style-type: none"> Non-randomised controlled trial Intervention: 3 months 	<ul style="list-style-type: none"> Studying the effect of individualised meals on nutritional 	<ul style="list-style-type: none"> IG: aged care home staff provided with 3x2hr education program by 	<ul style="list-style-type: none"> IG: MN prevalence 26% at baseline and 12% at follow-up; body weight 	<ul style="list-style-type: none"> Direct health care cost in IG: median £924, CG: £847 per year. Not

	<ul style="list-style-type: none"> • N=109 • Females=50%. • N= 6 aged care homes 	<ul style="list-style-type: none"> • Follow-up: 3 months for clinical data, 1 year for cost data. • Economic method: Health care unit cost comparison on direct healthcare costs. • Economic cost vs. benefit/effect measured: cost = Cost for each health care contact; Benefit/Effect = none included in economic analysis. 	<p>status among older people living in aged care home vs. control group and to estimate direct health care costs for both groups.</p>	<p>project leader - a nurse with nutrition background. MNA >24 (well nourished) offered snack (e.g. fruit, yoghurt) at midmeals. MNA <24 (risk of malnutrition/ malnourished) had modified food choices within existing food availability and costs: offered appetizer at lunch (e.g. soup, egg, herring), additional snacks (e.g. smoothies, bread and butter, milk and yoghurt) distributed throughout the day according to needs and preference.</p> <ul style="list-style-type: none"> • CG: No change to routine meals. 	<p>change at 3/12 follow-up 2.7±3.9kg; BMI at 3/12 follow-up 25.6±4.9kg/m².</p> <ul style="list-style-type: none"> • CG: Malnutrition prevalence 18% at baseline and 28% at follow-up; Body weight change at 3/12 follow-up -0.6±4.9kg; BMI at 3/12 follow-up 23.7±4.9 kg/m². • Between groups: • Body weight $p=0.0001$; • BMI $p=0.05$. 	<p>compared statistically.</p>
<p>Odlund Olin et. al. (2003) [27]</p>	<ul style="list-style-type: none"> • Sweden • Median age 80-83yrs (IQR 71-89) • N=40. • Female: 52%. • N=1 aged care home recruited (N=2 clusters [wards]). 	<ul style="list-style-type: none"> • Non-randomised clustered controlled intervention trial. • Intervention: 15 weeks • Follow-up: 29 weeks post-baseline/27 weeks post intervention commencement. • Economic method: Cost of Intervention. No analysis. • Economic cost vs. benefit/effect measured: cost = Cost of additional butter and cream; Benefit/Effect = none included in economic 	<p>To determine if a food-first approach will improve energy intake, activities of daily living and infections.</p>	<ul style="list-style-type: none"> • IG: Served regular hospital diet fortified with butter and cream (2100kCal/day). • CG: Served regular hospital diet (1600kCal/day). 	<ul style="list-style-type: none"> • IG: Compared with baseline, IG increased protein intake (median 48.3 [IQR: 41.8-54.3g] vs median 57.9 [IQR: 46.2-61.2g], $P<0.001$). ADL remained unchanged. • CG: worsened in ADL during the intervention (median score 15.5 [IQR: 10.0-17.0] increased to 16.0 [IQR: 15.0-18.0], $P<0.001$). • Between groups: No difference for number of infections. IG increased 	<p>£0.10 per resident per day</p>

		analysis.			energy intake (median 1437 [IQR: 1252-1617kcal] vs median 1840 [IQR: 1497-2012kcal], P<0.01).	
--	--	-----------	--	--	-----------------------------------------------------------------------------------------------	--

AT: Afternoon Tea, BMI: Body Mass Index, CC: Calf Circumference, CEA: Cost Effectiveness Analysis, CUA: Cost Utility Analysis, CG: control group, IG: Intervention group, kCal: kilocalorie, kg: kilogram, MAC: Mid Arm Circumference, ONS: Oral Nutrition Supplements, QALY: Quality Adjusted Life Year, RCT: Randomised Control Trial, USD: United States Dollar, vs.: versus, yrs: years.

Supplementary Table 2 – Cochrane Risk of Bias Table

Cochrane Risk of Bias: Quality assessment for interventional studies which report cost data related to nutrition in residential aged care							
Study	Random sequence generation (selection bias)	Allocation concealment (selection bias)	Blinding of participants and personnel (performance bias)	Blinding of outcome assessment (detection bias)	Incomplete outcome data (attrition bias)	Selective reporting (reporting bias)	Other bias
Rating	High risk of bias	High risk of bias	Unclear	Unclear	Unclear	Low risk of bias	Unclear
Evidence	<p>Edwards & Beck, 2002</p> <p>No randomisation was used (page 701-702). Due to the cross-over design, the CG received the intervention; however, the IG did not cross-over to the control group.</p> <p>Allocations were not concealed from participants and not described for investigators (page 702-703).</p> <p>Participants were not blinded from the intervention allocation; however, due to the study population (Alzheimer's Disease), the study participants could be considered to have been blinded. There was no description of blinding for the investigators (page 702-703).</p> <p>Blinding of investigators not described, but as reported using objective clinical measures, risk of bias is low (page 702-703). However, risk of bias for economic outcome reported is high as there is no method described.</p> <p>No attrition is reported; there was likely some attrition present due to the high risk sample (705-706).</p> <p>None detected</p> <p>No bias from funding organisation likely; however conflicts of interest were not declared.</p>						
Rating	Lee et. al. 2013	Low risk of bias	Unclear	Low risk of bias	Low risk of bias	Low risk of bias	Low risk of bias

Evidence		Stratified by gender then allocated to group based on drawing pieces of folded paper from a bag (page 1581).	It is not clear what was written on the pieces of paper and how they were then used by investigators to allocate group. It was not clear who drew out the pieces of paper. Later it describes participants did not know what group they were allocated to, therefore it is assumed participants did not draw out pieces of paper with the group allocation written on it (pages 1581 & 1582).	The study was double blind. Although participants could have observed they had a different afternoon tea, no discussion of the study allocation or intervention was made with the participant (pages 1581 and 1582).	Double blinded study, with anthropometry measured by non-staff research assistants (page 1583).	All participants accounted for, and <20% attrition (page 1582).	None detected	Appears to be no conflicts of interest or likely conflicts due to financial interests of funders/investigators.
Rating	Lorefalt et. al. 2011	High risk of bias	Unclear	Unclear	High risk of bias for clinical outcomes; low risk of bias for health care outcomes	Low risk of bias	Low risk of bias	Unclear
Evidence		No allocation method described in paper. Does not appear to be randomised.	Allocations and concealments of participants and investigators were not described at all. It is assumed the nurses implementing the intervention were aware that they formed an intervention group but it is not clear. It is not clear if participants were told they were in an intervention or control group.	Allocations and concealments of participants and investigators were not described at all. It is assumed the nurses implementing the intervention were aware that they formed an intervention group but it is not clear. It is not clear if participants were told they were in an intervention or control group.	Although it is not clear, it appears that the MNA and anthropometry outcomes were performed by the nurses who attended training and implemented the interventions. No blinding discussed for investigators or participants. Health care data does appear to be objective.	All 109 participants accounted for at 3 month follow-up (page 95). Assumed all accounted for in 1 year economic follow-up although not described.	None detected	No bias from funding organisation likely; however conflicts of interest were not declared.

Rating		Low risk of bias	Low risk of bias	High risk of bias	High risk of bias	Low risk of bias	Low risk of bias	Low risk of bias
Evidence	Neelemaat et. al. 2012	Computerised random number generator used to assign participants (page 184).	Allocation concealed in consecutiely n umbered opaque envelopes (page 184).	Participants, research assistant and researcher were no longer blinded from the intervention after the allocation was made (page 184).	Researchers were not blinded. Although cost data is mostly objective, results are related to subjectively completed tools of quality of life and physical function, which can be influenced by participant and researcher bias due to knowledge of intervention purpose.	High rate of attrition (32%); however was equally distributed between groups and unlikely to be related to intervention (12% due to death). Data presented with complete cases and per protocol.	None detected	Appears to be no conflicts of interest or likely conflicts due to financial interests of funders/investigators.
Rating	Odlund Olin, et.	High risk of bias	Unclear	Unclear	Unclear	Low risk of bias	Low risk of bias	Unclear

Evidence	al. 2003	No description of how wards were allocated to intervention and control. Seems to be conveniently selected (page 125).	It is not clear if participants in each ward were informed that the ward food was different (i.e. an intervention provided in one), or if any allocation concealment attempt was made.	Participants in the intervention group and RAC staff were told that a change to the meal composition will be made, but not further description. It is not clear if they were told this was part of a study. Nursing staff completed measurement of food consumption. It is not therefore clear if they understood the difference between meals being studied, and if this would affect performance and outcome assessment bias.	Participants in the intervention group and RAC staff were told that a change to the meal composition will be made, but not further description. It is not clear if they were told this was part of a study. Nursing staff completed measurement of outcomes. It is not therefore clear if they understood the difference between meals being studied, and if this would affect performance and outcome assessment bias.	All participants accounted for, and <20% attrition (page 126).	None detected	No bias from funding organisation likely; however conflicts of interest were not declared.
Rating		Unclear	Unclear	High risk of bias	Unclear	Low risk of bias	Low risk of bias	Low risk of bias
Evidence	Simmons et. al. 2010	Study describes that participants are randomised, but gives not detail as to how the sequence was generated (page 368)	Study describes that participants are randomised, but gives not detail as to how the participants were allocated (page 368)	There was no blinding of personnel. There was no description of blinding of residents, but as it is clear and obvious what intervention the received it is assumed they were not blinded (page 371)	The percentage food intake was measured by blinded personnel, but other measures including body weight were not. Use of cost data was objective and therefore low risk. There was poor description of blinding of outcome measures and attempts to make measures subjective, therefore risk of bias is unclear (page 368 and 371).	All participants accounted for, and although attrition was 27%, there was no statistical difference between groups. It is unlikely attrition was due to the intervention. However, it should be noted that the attrition rate per group	None detected	Appears to be no conflicts of interest or likely conflicts due to financial interests of funders/investigators.

					allocation was not described (page 369).		
Rating	Low risk of bias	Unclear	Unclear	Unclear	Low risk of bias	Low risk of bias	Low risk of bias
Evidence	Participants were randomised using a computer-generated random numbers table (page 2309)	Method of allocating the participants to groups was not described in the study.	There was no description of blinding of participants or personnel.	There was no description of blinding outcome assessors. As nurses timed themselves in their delivery of the intervention, this outcome does have some bias. Research staff completed other outcome measures, but it is not described if they were blinded (page 2309).	Attrition was substantial (36%); however, it was mostly even across groups and it was mostly for reasons common in RAC populations including death, transfer to hospice. There were low rates of consent withdrawal and nutritional order changes.	None detected	Appears to be no conflicts of interest or likely conflicts due to financial interests of funders/investigators.
	Simmons et. al. 2015						
Rating	Low risk of bias	Low risk of bias	High risk of bias	High risk of bias	Low risk of bias	High risk of bias	High risk of bias
	Elia et. al. 2017						

Evidence	& Parsons et. al. 2017	<p>"Randomisation, stratified according to malnutrition risk (medium or high risk of malnutrition) and type of care (residential or nursing care) was undertaken independently of the researchers using random number tables produced by Microsoft Excel for Windows 2003. The randomisation codes were generated by the chief investigator prior to commencement of the trial." (Parsons et al. page 135)</p>	<p>"The research dietitian enrolled and assigned participants to the interventions using opaque, sealed envelopes labeled with the random numbers containing the designated interventions. At the point of randomisation both the residents and the researchers were blinded to the designated intervention." (Parsons et al. page 135)</p>	<p>"...a prospective, randomised, parallel, open-label trial, which took place between August 2007 and March 2010" (Parsons et al. page 135). Acknowledged that both groups biased equally; however, likely to impact upon change from baseline variables.</p>	<p>"Due to the nature of the two different interventions researchers and participants were not blinded to the interventions after randomisation" (Parsons et al. page 141). Although cost data is mostly objective, results are related to subjectively completed tools of quality of life and physical function, which can be influenced by participant and researcher bias due to knowledge of intervention purpose.</p>	<p>Attrition was substantial (33%); however, it was mostly even across groups and it was mostly for reasons common in RAC populations including death and decline in health. A well-described intention-to-treat analysis was conducted and reported.</p>	<p>Data was reported in relation to ONS versus dietary advice; however, change from baseline was not tested for statistical significance for most outcomes. Therefore, it is not clear if dietary advice may have also been beneficial. Results are reported in such a way as to favour one intervention over the other. Cost/QALY was not reported for each intervention, but only in regards to ONS minus the dietary advice.</p>	<p>The study was funded by an unrestricted grant from a company which made the supplement; which results and conclusion favoured (see comment on selective reporting). In addition, there is significant concern about the possibility of poorly implementing the second comparator group (dietary advice group). Such an intervention is highly dependent on the skills of the particular dietitian. In addition, one of dietary advice followed by a 6-week follow-up for a malnourished patient would be seen as inadequate intervention, monitoring and follow-up by a treating clinical dietitian. A poor outcome in the dietary advice group would only serve to make the ONS (a standardised intervention which does not necessarily need follow-up or regular modification)</p>
----------	------------------------	----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------	---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------	----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------	----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------	-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------	-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------	-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------

