

**Depressive symptoms and adverse outcomes from hospitalization in older adults: Secondary outcomes of a trial of falls prevention education**

Haines, Terry P.; Williams, Cylie M.; Hill, Anne Marie; McPhail, Steven M.; Hill, D.; Brauer, Sandy G.; Hoffmann, Tammy C.; Etherton-Ber, Chris

*Published in:*  
Archives of Gerontology and Geriatrics

*DOI:*  
[10.1016/j.archger.2014.09.009](https://doi.org/10.1016/j.archger.2014.09.009)

Published: 01/01/2015

*Document Version:*  
Peer reviewed version

*Licence:*  
CC BY-NC-ND

[Link to publication in Bond University research repository.](#)

*Recommended citation(APA):*

Haines, T. P., Williams, C. M., Hill, A. M., McPhail, S. M., Hill, D., Brauer, S. G., Hoffmann, T. C., & Etherton-Ber, C. (2015). Depressive symptoms and adverse outcomes from hospitalization in older adults: Secondary outcomes of a trial of falls prevention education. *Archives of Gerontology and Geriatrics*, *60*(1), 96-102. <https://doi.org/10.1016/j.archger.2014.09.009>

**General rights**


Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

For more information, or if you believe that this document breaches copyright, please contact the Bond University research repository coordinator.

## Accepted Manuscript

Title: Depressive Symptoms and Adverse Outcomes from Hospitalization in Older Adults: Secondary Outcomes of a Trial of Falls Prevention Education

Author: Terry P. Haines Cylie. M. Williams Anne-Marie Hill Steven. M. McPhail D. Hill Sandy. G. Brauer Tammy. C. Hoffmann Chris Etherton-Beer



PII: S0167-4943(14)00160-5  
DOI: <http://dx.doi.org/doi:10.1016/j.archger.2014.09.009>  
Reference: AGG 3062

To appear in: *Archives of Gerontology and Geriatrics*

Received date: 13-11-2013  
Revised date: 25-6-2014  
Accepted date: 16-9-2014

Please cite this article as: Haines, T.P., Williams, Ce.M., Hill, A.-M., McPhail, Sn.M., Hill, D., Brauer, Sy.G., Hoffmann, Ty.C., Etherton-Beer, C., Depressive Symptoms and Adverse Outcomes from Hospitalization in Older Adults: Secondary Outcomes of a Trial of Falls Prevention Education., *Archives of Gerontology and Geriatrics* (2014), <http://dx.doi.org/10.1016/j.archger.2014.09.009>

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

1 Depressive Symptoms and Adverse Outcomes from Hospitalization in Older Adults:  
2 Secondary Outcomes of a Trial of Falls Prevention Education.  
3  
4  
5  
6

7 Associate Professor Terry P Haines

8  
9 Monash University /Monash Health

10 Allied Health Research Unit, Monash Health

11  
12 Kingston Hospital

13  
14 Cnr Warrigal and Kingston Rds,

15  
16 Cheltenham, VIC 3192, Australia

17  
18 Email: [terrence.haines@monash.edu](mailto:terrence.haines@monash.edu)

19  
20 Phone: +61 (0) 3 92651822  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31

32 Dr Cylie. M. Williams

33  
34 Allied Health Research Unit, Monash Health

35  
36 Kingston Hospital

37  
38 Cnr Warrigal and Kingston Rds,

39  
40 Cheltenham, VIC 3192, Australia

41  
42 Email: [Cylie.williams@southernhealth.org.au](mailto:Cylie.williams@southernhealth.org.au)

43  
44 Phone: +61 (0) 3 92651816  
45  
46  
47  
48  
49  
50  
51

52 Dr Anne-Marie Hill

53  
54 The University of Notre Dame Australia

55  
56 19 Mouat Street (PO Box 1225)

57  
58 Fremantle, Western Australia  
59  
60  
61  
62  
63  
64  
65

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

Australia 6959

Telephone +61 8 9433 0239

Facsimile: +61 8 9433 0210

Email: [anne-marie.hill@nd.edu.au](mailto:anne-marie.hill@nd.edu.au)

Dr Steven. M McPhail

Queensland University of Technology/ Queensland Health

[steven\\_mcphail@health.qld.gov.au](mailto:steven_mcphail@health.qld.gov.au)

Ph: +617 3406 2266

Professor Keith. D. Hill

Curtin University, School of Physiotherapy

Ph: +618 9266 3618

[Keith.Hill@curtin.edu.au](mailto:Keith.Hill@curtin.edu.au)

Professor Sandy. G. Brauer

University of Queensland

Head of Physiotherapy | School of Health & Rehab Sciences

The University of Queensland | Brisbane Qld 4072

Ph +61 7 3365 2317 | Fax +61 7 3365 2775

Email [s.brauer@uq.edu.au](mailto:s.brauer@uq.edu.au)

1 Dr Tammy. C. Hoffmann

2 Centre for Research in Evidence-Based Practice, Bond University

3  
4 Faculty of Health Sciences and Medicine

5  
6 Bond University QLD 4229

7  
8 +61 7 559 55522

9  
10 Email: thoffmann@bond.edu.au

11  
12  
13  
14  
15  
16 Professor Chris Etherton-Beer

17  
18 University of Western Australia

19  
20 School of Medicine and Pharmacology Royal Perth Hospital Unit

21  
22 35 Stirling Highway, CRAWLEY WA 6009

23  
24  
25  
26 Australia

27  
28 Ph: 61 8 9224 2750

29  
30 Email: christopher.etherton-beer@uwa.edu.au

31  
32  
33  
34  
35  
36 Corresponding author:

37  
38 Terry Haines,

39 Allied Health Research Unit, Southern Health

40 Kingston Hospital

41 Cnr Warrigal and Kingston Rds,

42 Cheltenham, VIC 3192, Australia

43 Email: [terrence.haines@monash.edu](mailto:terrence.haines@monash.edu)

44  
45  
46  
47  
48 Phone: +61 (0) 3 92651822

49  
50  
51  
52  
53 Keywords: depression, elderly, hospital, falls

54  
55  
56  
57 Key points:

- 58 1. Depressive symptoms are common amongst older hospital inpatients and

generally persist through to discharge.

2. Depressive symptoms are associated with longer length of stay and falls while in hospital
3. Depressive symptoms worsen if exposed to low intensity (video, workbook) falls prevention education material.

Word count: 3236

Accepted Manuscript

**Depressive Symptoms and Adverse Outcomes from Hospitalization in Older  
Adults: Secondary Outcomes of a Trial of Falls Prevention Education.**

Abstract

Depression is common in older people and symptoms of depression are known to substantially increase during hospitalisation. There is little known about predictors of depressive symptoms in older adults or impact of common interventions during hospitalisation. This study aimed to describe the magnitude of depressive symptoms, shift of depressive symptoms and the impact of the symptoms of depression amongst older hospital patients during hospital admission and identify whether exposure to falls prevention education affected symptoms of depression.

Participants (n=1206) were older adults admitted within two Australian hospitals, the majority of participants completed the Geriatric Depression Scale (GDS) short form at admission (n=1168). Participants' mean age was 74.7 ( $\pm$ SD 11) years and 47%(n=551) were male.

At admission 53% (619 out of 1168) of participants had symptoms of clinical depression and symptoms remained at the same level at discharge for 55% (543 out of 987). Those exposed to the low intensity education program had higher GDS scores at discharge than those in the control group (low intensity vs control n=652, adjusted regression coefficient (95% CI) = 0.24 (0.02, 0.45), p=0.03). The only factor other than admission level of depression that affected depressive symptoms change was if the participant was worried about falling.

1 Older patients frequently present with symptoms of clinical depression on admission to  
2 hospital. Future research should consider these factors, whether these are modifiable  
3  
4  
5 and whether treatment may influence outcomes.  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

Accepted Manuscript



**Introduction:**

1  
2  
3  
4  
5 Depression is common amongst older people,(Bryant, Jackson, & Ames, 2009; Djernes,  
6  
7 2006; Solhaug, Romuld, Romild, & Stordal, 2012) and it is well understood that the risk  
8  
9 of experiencing symptoms of depression is increased by cognitive impairment, illness  
10  
11 and limited access to friends and family support networks.(Djernes, 2006; Sheikh &  
12  
13 Yesavage, 1986) It is not surprising then that symptoms of depression have been  
14  
15 shown to increase in frequency and severity during times of hospitalisation.(Givens,  
16  
17 Jones, & Inouye, 2009) Recent research has found that over two thirds of older people  
18  
19 undertaking inpatient rehabilitation had some form of clinically significant psychiatric  
20  
21 comorbidity upon admission and over one third displayed significant symptoms of  
22  
23 depression at discharge.(Gluyas, Lum, Chong, Borg, & Haines, 2011)  
24  
25  
26  
27  
28  
29  
30

31  
32 Clinicians currently have little guidance to identify patients at high risk of developing  
33  
34 depressive symptoms or having worsening symptoms during their hospital admission.  
35  
36 This is important as early identification may permit delivery of services to prevent  
37  
38 development or worsening of depressive symptoms. Another issue of importance to  
39  
40 clinicians is whether health interventions designed to address other geriatric issues  
41  
42 (such as falls prevention) may have the unintended consequence of worsening  
43  
44 depressive symptoms. Previous research has demonstrated that older adults rarely  
45  
46 discuss falls with their health professionals (Lee, McDermott, Hoffmann, O'Connell, &  
47  
48 Haines, In Press) and that some find this discussion very confronting, as they commonly  
49  
50 associate falls with being old and frail. (Yardley et al., 2008) It was these challenges and  
51  
52 observations within the clinical environment that lead to the formulation of the study  
53  
54 questions.  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

1  
2 This study is a secondary analysis of data collected as a part of a randomised trial  
3  
4 investigating the efficacy of two patient education strategies for the prevention of falls  
5  
6 amongst older hospital inpatients. The analyses presented in this paper aim to i)  
7  
8 describe the magnitude symptoms of depression amongst older hospital patients, ii)  
9  
10 describe how symptoms of depression change during hospital admission, iii) identify  
11  
12 factors associated with worsening symptoms of clinical depression, iv) describe the  
13  
14 impact that symptoms of depression had on length of stay in hospital, falls, and the  
15  
16 probability of being discharged to a residential aged care facility, and v) determine  
17  
18 whether provision of either of two patient education programs for the prevention of  
19  
20 falls had an impact on change in depressive symptoms during hospital admission.  
21  
22  
23  
24  
25  
26  
27  
28

## 29 **Method:**

### 30 *Study Design*

31  
32 This study was a 3-group randomized trial. There were two active intervention groups:  
33  
34 a “*low intensity*” education program that involved providing a video and written  
35  
36 materials discussing the prevention of falls, and a “*high intensity*” education program  
37  
38 where the same video and written materials were provided along with face-to-face  
39  
40 education with a health professional. The control group received usual care  
41  
42 alone.(Haines et al., 2011; A.M. Hill, Hill, et al., 2009; A.M. Hill, McPhail, et al., 2009) The  
43  
44 secondary analysis, utilized observational data collected from participants within the  
45  
46 RCT to address the research questions.  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56

### 57 *Participants and setting*

58  
59 Participants were older adults admitted to acute (orthopedic and acute-respiratory  
60  
61  
62  
63  
64  
65

1 medicine) and subacute (geriatric assessment and rehabilitation) wards of the Princess  
2 Alexandra Hospital, Brisbane, Australia, and the acute (medical-surgical) and subacute  
3 (restorative–stroke rehabilitation) wards of Swan Districts Hospital, Perth, Australia.  
4  
5 Patients were enrolled until they were discharged, transferred to a non-study ward, or  
6  
7 died. Patients were excluded if they had previously participated in the patient  
8  
9 education program or were too ill to provide informed consent as determined by  
10  
11 hospital staff. A sample size of 1206 was recruited out of a possible 5162 admissions  
12  
13 across all study wards.  
14  
15  
16  
17  
18  
19  
20  
21

#### 22 *Measurements:*

23  
24 Symptoms of depression were measured using the Geriatric Depression Scale – Short  
25  
26 Form (GDS).(Sheikh & Yesavage, 1986) This was administered at admission and  
27  
28 discharge by a research assistant blinded to group allocation. The GDS scale was  
29  
30 specifically developed for detecting depression in older adults and consists of 15 items.  
31  
32 A cut-off score of six or above out of the maximum 15 indicates the presence of  
33  
34 clinically significant depressive symptoms.(Burns, Lawlor, & Craig, 2002)  
35  
36  
37  
38  
39  
40  
41

42 The number of falls each participant sustained while in hospital was collected using  
43  
44 three data sources as previous research has demonstrated single sources  
45  
46 underestimate the true number of falls (A. M. Hill et al., 2010) Falls meeting the World  
47  
48 Health Organization definition (“A fall is an event which results in a person coming to  
49  
50 rest inadvertently on the ground or floor or other lower level”(World Health  
51  
52 Organisation, 2010)) were included. The sources of falls data were i) computerized  
53  
54 incident reports, ii) hand searching of individual patient medical notes, and iii) weekly  
55  
56 patient interviews (or at patient discharge if earlier than one week). Falls captured  
57  
58  
59  
60  
61  
62  
63  
64  
65

through any of these three approaches were included.

Health-related quality of life (HRQoL) was measured using the EQ-5D-3L.(Rabin & de Charro, 2001) This scale measures health-related quality of life over five domains: mobility, self care, usual activities, pain and/or discomfort, and anxiety/depression. The participant rates each domain in one of three levels of responses: no problem, some problems / moderate, or severe problems / unable. The Dolan scoring approach(Dolan, 1997) was applied creating a multi-attribute utility score with a possible range from -0.59 to 1.0; where 0 and 1 represent death and perfect health respectively.

Cognitive impairment was measured using the Short-Portable Mental Status Questionnaire. (Pfeiffer, 1975) Higher scores on this the Short-Portable Mental Status Questionnaire indicate better cognitive function.

Self-perceived risk of falling and anxiety\_about falling were measured with a patient survey as part of baseline and discharge measures. This survey used single items “I think I will fall while in hospital” and “I am worried about falling while in hospital” respectively. A five point Likert scale (strongly agree through to strongly disagree) was used for each item.

Other demographic and outcome variables were collected from patient medical records. These included: Age, gender, highest level of education, history of number of falls in the past 6 months, whether English was spoken as a first language, length of stay in hospital (measured in days), admission diagnosis, admission and discharge destination.

1  
2  
3  
4  
5  
6  
7 *Procedure:*  
8

9  
10 Pre-study training was provided to hospital staff on study wards regarding  
11 classification of falls and procedures for recording falls on incident reports using  
12 previously developed video materials.(Haines, Cornwell, Fleming, Varghese, & Gray,  
13 2008; A.M. Hill, McPhail, et al., 2009) Patients consenting to participate in the study  
14 undertook a baseline assessment of all study measures. Participants were then  
15 allocated to one of the two intervention groups or the control group within the falls  
16 trial. Discharge measurements for health-related quality of life, depression and  
17 cognitive function were undertaken (with assistance from research assistants to  
18 administer the questionnaires) within 48 hours of discharge.  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31

32  
33  
34 *Analysis:*  
35

36  
37 The baseline demographics between groups were explored with means, standard  
38 deviations and frequencies. Differences between the groups were analysed using t-tests,  
39 chi-squared and logistic regression. Associations between factors predictive of  
40 admission depressive symptoms and change in depressive symptoms were examined  
41 using path analysis(Byrne, 2010) undertaken with Amos 19(Arbuckle, 2006) software  
42 package. Both unstandardized and standardized regression coefficients of associations  
43 were generated. Unlike multiple regression analysis, path analysis was used to  
44 explicitly depict both direct and indirect (mediation) effects between exogenous  
45 (predictor) variables and endogenous (outcome) variables. The path analysis included  
46 both admission GDS and change in depressive symptoms as endogenous variables.  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

1 Variables collected at baseline were used as exogenous variables (including use of  
2 admission GDS as an exogenous variable to predict change in GDS as an endogenous  
3 variable).  
4  
5  
6  
7  
8  
9

10 Univariate analyses were also used to build the model initially, with indirect /  
11 mediation effects then added to the model. Modification indices were used to assist  
12 with the path model development and refinement, as these identify potential  
13 associations not already specified in the model. The model was refined by removing  
14 any exogenous variables that had no significant direct effect on either of the  
15 endogenous variables. The model fit was examined using the Chi Square, Root Mean-  
16 Square Error of Approximation (values <0.05 indicate good model fit), and Comparative  
17 Fit Index statistics (values >0.95 indicate good fit). Moderation analysis was then  
18 undertaken to examine whether group allocation in the randomized controlled trial  
19 influenced the associations identified. For this, pairwise parameter comparisons were  
20 undertaken across the three groups in the trial so that critical ratios for comparisons  
21 between parameters could be calculated. These critical ratios are a z-test for the  
22 difference between coefficients where a value >1.96 represents a p-value (two-tailed)  
23 <0.05 indicative of a statistically significant moderation. (Fairchild & MacKinnon, 2009)  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45

## 46 **Results**

47 There were 1206 participants enrolled in the larger RCT. Of these participants, 1168  
48 completed a baseline GDS and 987 participants completed both baseline and discharge  
49 GDS. Unplanned discharges or transfers prevented discharge assessment from  
50 occurring for some participants which resulted in the lower number of participants  
51 completing both baseline and discharge (n=181). Table 1 displays the demographic  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

1 data for participants within each group. All groups were similar in all aspects except for  
2 the Low vs High intensity group's admission to a rehabilitation setting ( $p=0.039$ ).  
3  
4

5  
6  
7 Clinically significant symptoms of depression (GDS cut-off of 6 or above) were present  
8 in 53% ( $n=619$  of 1168) of this population at admission. When the magnitude and  
9 distribution of depression symptoms was explored for the whole group at admission  
10 and discharge, it was determined that the symptoms of depression worsened from  
11 admission baseline to discharge for 18% ( $n=177$  of 987) of this sample, but remained  
12 unchanged for 55% ( $n=543$  of 987). The change in depressive symptoms was  
13 significantly different from zero ( $p<0.001$ ), indicating that depressive symptoms on  
14 average measured using the GDS decreased during the admission. A scatterplot of the  
15 admission GDS vs discharge GDS shows change in all groups during admission (Figure 1)  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30

31  
32 The  $n$ , mean (sd) discharge GDS score for the control group was  $n=314$ , 6.4 (1.8), for  
33 the low intensity group was  $n=342$ , 6.8 (1.9), and for the high intensity group was  
34  $n=338$ , 6.5 (1.8). Pairwise comparisons of discharge GDS scores adjusted for baseline  
35 GDS scores demonstrated that those exposed to the low intensity education program  
36 had higher GDS scores at discharge than those in the control group (low intensity vs  
37 control  $n=652$ , adjusted regression coefficient (95% CI) = 0.24 (0.02, 0.45),  $p=0.03$ ).  
38  
39 Other adjusted pairwise contrasts were not significant (high intensity vs control  $n=648$ ,  
40 adjusted regression coefficient (95% CI) = -0.18 (-0.39,0.03),  $p=0.10$ , high intensity vs  
41 low intensity  $n= 674$ , adjusted regression coefficient (95% CI) = 0.04 (-0.17, 0.25),  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

Table 2 and Figure 2 present the path analysis that identifies factors predictive of a

1 higher baseline GDS score, and/or a greater change in GDS score during the admission.

2 The factors that predicted higher levels of depressive symptoms at baseline ( $r^2=0.09$ )  
3 were, having a history of falls in the past six months, male gender, not having English as  
4 a first language, having lower levels of education, being admitted from a residential  
5 aged care facility, being recruited for this study from a rehabilitation ward, having a  
6 higher self-perceived risk of falling and being worried about falling while in hospital.  
7 Lower baseline depressive symptoms and higher levels of worry about the risk of  
8 falling were also associated with worsening depressive symptoms over the admission  
9 in hospital ( $r^2=0.26$ ).  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24

25 The direct effect of worry about the risk of falling in hospital on the change in GDS (the  
26 more worried that older adult was about falling predicted worsening depressive  
27 symptoms), and the indirect effect of worry about the risk of falling mediated through  
28 admission depressive symptoms were opposite in direction. However these effects  
29 appeared to cancel each other out when considering the total standardised effect of  
30 being worried about falls at admission and change in GDS score (Table 2). The total  
31 standardized effects indicated that baseline GDS had the greatest influence on change in  
32 depressive symptoms during admission. Two model fit indices (RMSEA =0.019, CFI  
33 =0.984) indicated good fit while the third indicated moderate fit ( $\chi^2=0.128$ ).  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49

50 Lastly, the moderator variable analysis examining the effect of group allocation (within  
51 the randomized controlled trial) in these results identified that there was a significant  
52 difference in path coefficients generated between groups in two out of 45 pairwise  
53 comparisons. These were the direct effect of worry about the risk of falling on change  
54 in depressive symptoms (high intensity intervention vs. control group pairwise  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65



1 comparison), and the direct effect of worry about the risk of falling on baseline  
2 depressive symptoms (high intensity intervention vs. low intensity intervention group  
3 pairwise comparison). The direct effect of worry about the risk of falling illustrated that  
4 the high intensity intervention provided in the randomized trial may have changed the  
5 nature of the association between the participant's worry about falling and increased  
6 depressive symptoms over the admission relative to the control group. In the control  
7 group, this standardized direct effect (regression coefficient -0.02) indicated a weak  
8 relationship between worry about falls and increasing depression. In the intensive  
9 intervention group, this standardized direct effect (regression coefficient -0.18)  
10 indicated that higher levels of worry about falls were associated with increasing  
11 depressive symptoms. The latter moderation effect appears to represent random  
12 inherent variability in the trial data as the assessment of worry about falls and baseline  
13 GDS took place prior to randomisation. These two moderation effects were the only  
14 significant effects identified from 45 pairwise comparisons examined.  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36

## 37 Discussion

38 This study has identified that a high proportion (53%) of older hospital patients  
39 experienced clinically significant depressive symptoms at admission and that these  
40 remained at the same level at discharge for the majority, while they worsened for 18%  
41 of participants. This finding is important as it points to a considerable magnitude of  
42 burden of depressive symptoms amongst hospitalised older adults.  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53

54 Our study for the first time provides experimental evidence of an association between  
55 involvement in a low-intensity falls prevention education program and increasing  
56 depressive symptoms during hospitalisation. The low intensity intervention in this trial  
57  
58  
59  
60  
61  
62  
63  
64  
65

1 consisted of showing a video designed to raise awareness about falls and falls  
2 prevention strategies and providing a booklet with similar information without  
3 additional follow-up. This low-cost education approach may therefore be harmful, as it  
4 has already been demonstrated that it provides no benefit in terms of preventing  
5 falls(Haines 2011). It is possible that viewing this material may have made participants  
6 more concerned about falls, or thought more about the related processes of ageing and  
7 frailty without providing health-professional follow-up to discuss and counsel patients  
8 on these issues. However, it must again be stated that this was an opportunistic  
9 secondary analysis of data arising from this trial which was not pre-planned, and  
10 should therefore be viewed as a hypothesis generating finding rather than a  
11 confirmatory one.  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28

29 A previous systematic review and meta-analysis on risk factors for depression in older  
30 adults within the community found that women were at a higher risk of depression  
31 than males.(Cole & Dendukuri, 2003) Our study, in contrast, found that men were at  
32 higher risk and a much higher prevalence of depressive symptoms at admission. The  
33 finding of gender differences is similar to previous research that has highlighted  
34 women as being at higher or equivalent risk of falling while in the community, but  
35 when in hospital, it is men who are at higher risk.(Haines & Hill, 2011) These findings  
36 may indicate that men have greater difficulty with adjusting to being unwell or  
37 adapting to the hospital environment.  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53

54 Those who had a history of falls within the past six months and those who worried  
55 about falling were also observed to be at greater risk of experiencing depressive  
56 symptoms. It was expected that a recent fall would be a predictor for depressive  
57  
58  
59  
60  
61  
62  
63  
64  
65

1 symptoms as past research has identified a relationship between depressive symptoms  
2 and falls.(Biderman, Cwikel, Fried, & Galinsky, 2002) It was interesting that this  
3 relationship was partially mediated by worry about falling in the future indicating that  
4 addressing fear of falling may be a potential consideration for managing depressive  
5 symptoms in older hospital patients. This therapeutic recommendation may be further  
6 justified by previous findings that depressive symptoms in the older population were  
7 not associated with an individual's physical and functional ability(Gluyas et al., 2011)  
8 despite physical and functional ability being closely associated with falls.(Shumway-  
9 Cook et al., 2009)

10 Lower levels of education have been previously identified within the literature as a risk  
11 for depression in older people(Chang-Quan, Zheng-Rong, Yong-Hong, Yi-Zhou, & Qing-  
12 Xiu, 2010) corroborating the findings of this research. Little is known about why  
13 education levels have been identified as a predictive factor. Cognitive impairment  
14 amongst older adults has previously been associated with lower levels of education,  
15 however, cognitive function scores taken at baseline using the abbreviated mental test  
16 score were not a significant factor in the path model. This result may possibly be due to  
17 multi collinearity with the factors admission from a residential aged care facility,  
18 previous falls and level of education. Similarly, little is known about why English as a  
19 second language is a predictor for depressive symptoms in older adults. Often these  
20 patients are excluded from medical trials(Frayne, Burns, Hardt, Rosen, & Moskowitz,  
21 1996) yet appear to be an important sub-group of the population for intervention. One  
22 previous study amongst older Korean-American immigrants found that English  
23 language proficiency was associated with depressive symptoms,(Park & Bernstein,  
24 2008) while a study of older Iranian-Australian immigrants found anxiety and

1 depressive symptoms were associated with proficiency in English.(Alizadeh-Khoei,  
2 Mathews, & Hossain, 2011) These findings indicate that health services should be  
3 particularly cognisant of the risk of depression amongst their patients with limited  
4 English language skills.  
5  
6  
7  
8  
9

10 An important limitation of this research was that data were not collected on whether  
11 patients were treated for their depressive symptoms while in hospital. It is possible  
12 that some of the associations examined in this study may have been biased by  
13 treatments that were provided. This may explain why we were unable to identify any  
14 factors predictive of change in depressive symptoms other than admission depressive  
15 symptoms (which itself may have been caused by regression to the mean). It is  
16 unknown what, if any, impact depressive symptoms had on physical function and other  
17 behaviours which may moderate the relationship between depressive symptoms and  
18 increased length of stay or falls.  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36

37 This study has highlighted the potential importance of addressing depressive  
38 symptoms amongst older hospital patients, and highlighted patient groups for whom  
39 this may be particularly relevant. Further research is required to determine if the  
40 depressive symptoms observed during the inpatient period persist following discharge  
41 and whether any further burden may be associated with this. Further research is also  
42 required to investigate the effectiveness of therapeutic approaches initiated in the  
43 inpatient setting, and the potential unintended consequences of providing low intensity  
44 falls prevention education interventions.  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58

## 59 **Conclusion**

60  
61  
62  
63  
64  
65

1 Clinically important levels of depressive symptoms are common amongst older hospital  
2 inpatients and generally persist through to discharge. These symptoms may be  
3  
4 exacerbated by participating in a low intensity falls prevention education program.  
5  
6 Some factors predictive of depressive symptoms may indicate a relationship to frailty  
7  
8 (e.g. history of falls, admission from a residential care facility), though other factors  
9  
10 indicate a more complex picture than this.  
11  
12  
13  
14  
15  
16

### 17 **Conflict of Interest:**

18  
19 TPH is director of Hospital Falls Prevention Solutions Pty Ltd. This company provides  
20  
21 consultancy services on the subject of falls prevention in hospitals and provides  
22  
23 training in use of the Safe Recovery patient education program which was used in the  
24  
25 trial described in this manuscript.  
26  
27  
28  
29  
30  
31

### 32 **Description of Author roles and funding:**

33  
34 TPH, AMH, SMM, KDH, SGB, TCH, and CEB designed the study. AMH collected the data  
35  
36 and TPH was responsible for data analysis. All authors critically interpreted the results;  
37  
38 CW and TPH drafted the manuscript and circulated repeatedly among all authors for  
39  
40 critical revision. All authors approved the final version.  
41  
42  
43

44 This research was also funded by a project grant from the National Health and Medical  
45  
46 Research Council of Australia. TPH and TCH were supported by a Career Development  
47  
48 Fellowship, SMM and AMH were supported by Early Career Fellowships from the  
49  
50 National Health and Medical Research Council of Australia.  
51  
52  
53  
54  
55  
56  
57  
58

### 59 **References**

- 1 Alizadeh-Khoei, M., Mathews, R. M., & Hossain, S. Z. (2011). The Role of Acculturation in  
 2 Health Status and Utilization of Health Services among the Iranian Elderly in  
 3 Metropolitan Sydney. *Journal of Cross-Cultural Gerontology*, 26(4), 397-405.
- 4 Arbuckle, J. L. (2006). Amos (Version 19). Chicago: SPSS.
- 5 Biderman, A., Cwikel, J., Fried, A. V., & Galinsky, D. (2002). Depression and falls among  
 6 community dwelling elderly people: a search for common risk factors. *Journal of*  
 7 *Epidemiology and Community Health*, 56(8), 631-636. doi: 10.1136/jech.56.8.631
- 8  
 9 Bryant, C., Jackson, H., & Ames, D. (2009). Depression and anxiety in medically unwell  
 10 older adults: prevalence and short-term course. *International psychogeriatrics / IPA*,  
 11 21(4), 754-763. doi: 10.1017/S1041610209009399
- 12  
 13 Burns, A., Lawlor, B., & Craig, S. (2002). Rating scales in old age psychiatry. *The British*  
 14 *journal of psychiatry : the journal of mental science*, 180, 161-167.
- 15  
 16 Byrne, B. M. (2010). *Structural equation modeling with AMOS* (2nd ed.). New York:  
 17 Routledge.
- 18  
 19 Chang-Quan, H., Zheng-Rong, W., Yong-Hong, L., Yi-Zhou, X., & Qing-Xiu, L. (2010).  
 20 Education and risk for late life depression: a meta-analysis of published literature.  
 21 *International journal of psychiatry in medicine*, 40(1), 109-124.
- 22  
 23 Cole, M. G., & Dendukuri, N. (2003). Risk factors for depression among elderly community  
 24 subjects: a systematic review and meta-analysis. *The American journal of psychiatry*,  
 25 160(6), 1147-1156.
- 26  
 27 Djernes, J. K. (2006). Prevalence and predictors of depression in populations of elderly: a  
 28 review. *Acta psychiatrica Scandinavica*, 113(5), 372-387. doi: 10.1111/j.1600-  
 29 0447.2006.00770.x
- 30  
 31 Dolan, P. (1997). Modeling valuations for EuroQol health states. *Medical care*, 35(11),  
 32 1095-1108.
- 33  
 34 Fairchild, A. J., & MacKinnon, D. P. (2009). A general model for testing mediation and  
 35 moderation effects. *Prevention Science*, 10(2), 87-99.
- 36  
 37 Frayne, S., Burns, R., Hardt, E., Rosen, A., & Moskowitz, M. (1996). The exclusion of non-  
 38 english-speaking persons from research. *Journal of General Internal Medicine*, 11(1),  
 39 39-43. doi: 10.1007/BF02603484
- 40  
 41 Givens, J. L., Jones, R. N., & Inouye, S. K. (2009). The overlap syndrome of depression and  
 42 delirium in older hospitalized patients. *Journal of the American Geriatrics Society*,  
 43 57(8), 1347-1353. doi: 10.1111/j.1532-5415.2009.02342.x
- 44  
 45 Gluyas, C., Lum, C., Chong, S. Y., Borg, C., & Haines, T. P. (2011). Prevalence and  
 46 influence of psychiatric comorbidity on rehabilitation outcome for older hospital  
 47 inpatients. *International psychogeriatrics / IPA*, 1-13. doi:  
 48 10.1017/S1041610211000731
- 49  
 50 Haines, T. P., Cornwell, P., Fleming, J., Varghese, P., & Gray, L. (2008). Documentation of  
 51 in-hospital falls on incident reports: Qualitative investigation of an imperfect process.  
 52 *BMC Health Services Research*, 8(1), 254.
- 53  
 54 Haines, T. P., & Hill, A. M. (2011). Inconsistent results in meta-analyses for the prevention  
 55 of falls are found between study-level data and patient-level data. *Journal of Clinical*  
 56 *Epidemiology*, 64(2), 154-162.
- 57  
 58 Haines, T. P., Hill, A. M., Hill, K. D., McPhail, S., Oliver, D., Brauer, S., . . . Beer, C. (2011).  
 59 Patient education to prevent falls among older hospital inpatients: A randomized  
 60 controlled trial. *Archives of Internal Medicine*, 171(6), 516-524.
- 61  
 62 Hill, A. M., Hill, K., Brauer, S., Oliver, D., Hoffmann, T., Beer, C., . . . Haines, T. P. (2009).  
 63 Evaluation of the effect of patient education on rates of falls in older hospital patients:  
 64 description of a randomised controlled trial. *BMC Geriatr*, 9, 14. doi: 10.1186/1471-  
 65

2318-9-14

- 1 Hill, A. M., Hoffmann, T., Hill, K., Oliver, D., Beer, C., McPhail, S., . . . Haines, T. P.  
2 (2010). Measuring falls events in acute hospitals-a comparison of three reporting  
3 methods to identify missing data in the hospital reporting system. *Journal of the*  
4 *American Geriatrics Society*, 58(7), 1347-1352. doi: 10.1111/j.1532-  
5 5415.2010.02856.x
- 6 Hill, A. M., McPhail, S., Hoffmann, T., Hill, K., Oliver, D., Beer, C., . . . Haines, T. P.  
7 (2009). A randomized trial comparing digital video disc with written delivery of falls  
8 prevention education for older patients in hospital. *Journal of the American*  
9 *Geriatrics Society*, 57(8), 1458-1463.
- 10 Lee, D. C., McDermott, F., Hoffmann, T., O'Connell, B., & Haines, T. (In Press). "They will  
11 tell me if there is a problem": Limited discussion between health professionals, older  
12 adults and their caregivers on falls prevention during and after hospitalization. *Health*  
13 *Education Research*.
- 14 Park, S. Y., & Bernstein, K. S. (2008). Depression and Korean American immigrants.  
15 *Archives of psychiatric nursing*, 22(1), 12-19. doi: 10.1016/j.apnu.2007.06.011
- 16 Pfeiffer, E. (1975). A short portable mental status questionnaire for the assessment of organic  
17 brain deficit in elderly patients. *Journal of the American Geriatrics Society*, 23(10),  
18 433-441.
- 19 Rabin, R., & de Charro, F. (2001). EQ-5D: a measure of health status from the EuroQol  
20 Group. *Annals of medicine*, 33(5), 337-343.
- 21 Sheikh, J. I., & Yesavage, J. A. (1986). Geriatric Depressive Scale (GDS); recent evidence  
22 and development of a shorter version. *Clinical Gerontologist*, 5, 165-171.
- 23 Shumway-Cook, A., Ciol, M. A., Hoffman, J., Dudgeon, B. J., Yorkston, K., & Chan, L.  
24 (2009). Falls in the Medicare population: incidence, associated factors, and impact on  
25 health care. *Physical Therapy*, 89(4), 324-332.
- 26 Solhaug, H. I., Romuld, E. B., Romild, U., & Stordal, E. (2012). Increased prevalence of  
27 depression in cohorts of the elderly: an 11-year follow-up in the general population -  
28 the HUNT study. *International psychogeriatrics / IPA*, 24(1), 151-158. doi:  
29 10.1017/S1041610211001141
- 30 Weisberg, H. F. (1992). *Central tendency and variability*. Newbury Park, CA: Sage  
31 Publications.
- 32 World Health Organisation. (2010). Definition of a fall: :  
33 [http://www.who.int/violence\\_injury\\_prevention/other\\_injury/falls/en/index.html](http://www.who.int/violence_injury_prevention/other_injury/falls/en/index.html) /  
34 (last accessed Jan 2010).
- 35 Yardley, L., Kirby, S., Ben-Shlomo, Y., Gilbert, R., Whitehead, S., & Todd, C. (2008). How  
36 likely are older people to take up different falls prevention activities? *Prev Med*,  
37 47(5), 554-558. doi: 10.1016/j.ypmed.2008.09.001
- 38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

Table 1. Demographics of participants within each group

	Control Group	Materials Only Group	Complete Intervention	Control Vs Low Intensity	Control Vs High Intensity	Low Intensity Vs High Intensity
	Mean(SD) or n (%)	Mean(SD) or n (%)	Mean(SD) or n (%)	p	p	p
Total Number of Participants	381 (32%)	424 (36%)	401 ((33%)			
Age	75.25(10.14)	74.73(11.75)	75.25 (10.99)	0.501	1.000	0.510
Gender (male)	203(53%)	223(53%)	216(54%)	0.856	0.870	0.715
Education level				0.995	0.092	0.737
Primary	113 (30%)	120 (28%)	111 (29%)			
Year 10	163 (43%)	189 (45%)	171 (44%)			
Year 12	38 (10%)	46 (11%)	47 (12%)			
TAFE/Diploma	47 (12%)	46 (11%)	53 (13%)			
Bachelor degree	14 (4%)	17 (4%)	5 (1%)			
Postgraduate	3 (1%)	6 (1%)	1 (1%)			
English as first language	348 (91%)	376 (89%)	376 (94%)	0.212	0.390	0.697
Faller in previous 6 months	209 (55%)	245 (58%)	212 (53%)	0.383	0.550	0.134
Discharged to residential care	51 (13%)	63 (15%)	45 (11%)	0.600	0.920	0.123
Admitted to a rehabilitation setting	139 (37%)	181 (43%)	143 (36%)	0.073	0.811	0.039
Statement response: <i>I am worried about falling while in hospital</i>				0.412	0.619	0.761
Strongly agree	18 (5%)	19 (4%)	26 (6%)			
Agree	99 (26%)	124 (29%)	103 (26%)			
Undecided	13 (3%)	16 (4%)	13 (3%)			
Disagree	169 (44%)	177 (42%)	174 (44%)			
Strongly disagree	72 (19%)	82 (19%)	82 (20%)			
(Missing data)	(9 (2%))	(5 (1%))	(3 (1%))			
Baseline GDS	6.90 (2.02)	7.13 (2.04)	6.69 (2.01)	0.115	0.155	0.002
GDS Change	-0.46 (1.69)	-0.32 (1.59)	-0.19 (1.56)	0.261	0.032	0.282



Table 2. Standardized total effects from path analysis model

	Gender	Faller in previous 6 months	Rehabilitation Admission	English not first language	Worried about falling	Admission from residential aged care facility	Education level	GDS admission score
Faller in previous 6 months	-0.089							
Rehabilitation Admission	-0.006	0.067						
Worried about falling	0.016	-0.182	-0.119					
Admission from residential aged care facility	-0.12							
Education level	0.068							
GDS admission score	0.095	0.12	0.127	-0.075	-0.185	0.07	-0.101	
GDS change score	-0.051	-0.048	-0.057	0.039	0.016	-0.037	0.053	-0.523

\*GDS = Geriatric Depression Scale – Short Form

1  
2  
3  
4  
5  
6  
7 Figure 1. Scatterplots of overall admission GDS and discharge GDS and for each group.  
8  
9

10 Figure 2: Path analysis showing standardized regression coefficients, r2 values and model fit statistics  
11

12 GDS = Geriatric Depression Scale – Short Form

13 Df = Degrees of freedom

14 CFI = Comparative fit index

15 RMSEA = Root mean square residual and standardized root mean square residual  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49

Figure 1

