

PG-SGA SF is a suitable screening tool for identifying the risk of malnutrition in outpatients with cancer

Lee, Gahee; de Groot, Lynette; Ackerie, Antoinette; Van der Meij, Barbara S

Licence:
CC BY-NC-ND

[Link to output in Bond University research repository.](#)

Recommended citation(APA):

Lee, G., de Groot, L., Ackerie, A., & Van der Meij, B. S. (2019). *PG-SGA SF is a suitable screening tool for identifying the risk of malnutrition in outpatients with cancer*. Poster session presented at Dietitians Association of Australia 36th National Conference, Gold Coast, Queensland, Australia.

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

For more information, or if you believe that this document breaches copyright, please contact the Bond University research repository coordinator.

PG-SGA SF IS A SUITABLE SCREENING TOOL FOR IDENTIFYING THE RISK OF MALNUTRITION IN OUTPATIENTS WITH CANCER

Gahee Lee^{1,2}, Lynette De Groot², Antoinette Ackerie² and Barbara van der Meij^{1,2,3}

¹ Nutrition and Dietetics, Faculty of Health Sciences and Medicine, Bond University, Gold Coast, QLD, Australia

² Dietetics and Foodservices, Mater Health, Brisbane, QLD, Australia

³ Mater Research Institute, University of Queensland, Brisbane, QLD, Australia

In cancer centres, a valid malnutrition screening tool is necessary to identify patients at risk of malnutrition and ensure early nutrition intervention. The aim of this study was to explore the prevalence of malnutrition in outpatients with cancer and to investigate the validity of the Patient-Generated Subjective Global Assessment Short Form (PG-SGA SF) as compared to 2 reference standards: the Patient-Generated Subjective Global Assessment (PG-SGA) and the newly developed criteria for malnutrition by the Global Leadership Initiative on Malnutrition (GLIM). A sensitivity of 80% and specificity of 60% were deemed acceptable for the PG-SGA SF. In 141 adult outpatients with cancer receiving IV day treatment, we assessed the PG-SGA SF and compared to PG-SGA and GLIM. According to the PG-SGA, 117 (83%) of 141 patients were identified as well-nourished (PG-SGA A), and 23 (16.3%) and 1 (0.7%) as moderately (PG-SGA B) and severely (PG-SGA C) malnourished. The PG-SGA SF had a sensitivity of 86% and specificity of 67% as compared to PG-SGA, and the agreement between these tools was 'Fair' ($k=0.3$). PG-SGA SF had a relatively low sensitivity and specificity as compared to the GLIM, 42% and 53% respectively, and a 'Poor' agreement ($k<0$). In conclusion, approximately 1 out of 5 outpatients with cancer are malnourished. PG-SGA SF meets the professional standard and could be a suitable nutrition screening tool in outpatients with cancer. The GLIM criteria are different to PG-SGA ratings and further research is required to determine their value in oncology outpatients.