

Bond University  
Research Repository



## Cutaneous markers of photo-damage and risk of Basal cell carcinoma of the skin: a meta-analysis

Khalesi, Mohammad; Whiteman, David C.; Doi, Suhail A R; Clark, Justin; Kimlin, Michael G; Neale, Rachel E

*Published in:*  
Cancer Epidemiology, Biomarkers and Prevention

*DOI:*  
[10.1158/1055-9965.EPI-13-0424](https://doi.org/10.1158/1055-9965.EPI-13-0424)

*Licence:*  
Other

[Link to output in Bond University research repository.](#)

*Recommended citation(APA):*  
Khalesi, M., Whiteman, D. C., Doi, S. A. R., Clark, J., Kimlin, M. G., & Neale, R. E. (2013). Cutaneous markers of photo-damage and risk of Basal cell carcinoma of the skin: a meta-analysis. *Cancer Epidemiology, Biomarkers and Prevention*, 22(9), 1483-9. <https://doi.org/10.1158/1055-9965.EPI-13-0424>

### General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

For more information, or if you believe that this document breaches copyright, please contact the Bond University research repository coordinator.

## **Cutaneous markers of photo-damage and risk of basal cell carcinoma of the skin: A meta-analysis**

Mohammad Khalesi<sup>1,2,3</sup>, David C. Whiteman<sup>1,2</sup>, Suhail A.R. Doi<sup>4</sup>, Justin Clark<sup>4</sup>, Michael G. Kimlin<sup>2,3</sup>, and Rachel E. Neale<sup>1,2</sup>

<sup>1</sup>Department of Population Health, Queensland Institute of Medical Research, Brisbane, Queensland, Australia; <sup>2</sup>NHMRC Centre for Research Excellence in Sun and Health, Brisbane, Queensland, Australia; <sup>3</sup>AusSun Research Lab, Institute of Health and Biomedical Innovation, Queensland University of Technology, Brisbane, Queensland, Australia; <sup>4</sup>School of Population Health, University of Queensland, Brisbane, Australia

### **Markers of photo-damage and risk of basal cell carcinoma**

**Key words:** basal cell carcinoma; meta-analysis; actinic keratoses; telangiectasia; solar elastosis; solar lentigines

### **Financial support:**

Mohammad Khalesi is funded by the NHMRC Centre for Research Excellence in Sun and Health. David Whiteman is funded by Australian Research Council. Michael Kimlin is funded by Cancer Council Queensland.

**Corresponding Author:** Dr Mohammad Khalesi, M.D, PhD Scholar, Cancer Aetiology and Prevention Group, Department of Population Health, Queensland Institute of Medical Research, 300 Herston Road, Herston, Queensland 4006, Australia; E-mails: mohammad.khalesi@qimr.edu.au, khalesi\_mohammad@yahoo.com; Phone: +61 7 3845 3565; Fax: +61 7 3845 3503

### **Conflict of interest**

The authors have no conflict of interest to declare.

**Acknowledgment:** None

**Prior presentation:** None

**Word count:** 2400

**Total number of tables:** 3

**Total number of figures:** 1 (4 parts)

**Total number of supplementary tables:** 1

**Abbreviations:** BCC: basal cell carcinoma; SCC: squamous cell carcinoma; AK: actinic keratosis; OR: odds ratio; RR: relative risk; CI: confidence interval; QE: quality effects; RE: random effects

## **ABSTRACT**

**BACKGROUND:** Epidemiologic research has demonstrated that cutaneous markers of photo-damage are associated with risk of basal cell carcinoma (BCC). However there has been no previous attempt to calculate pooled risk estimates.

**METHODS:** We conducted a systematic review and meta-analysis after extracting relevant studies published up to January 2013 from five electronic databases. Eligible studies were those that permitted quantitative assessment of the association between histologically-confirmed BCC and actinic keratoses, solar elastosis, solar lentigines, or telangiectasia.

**RESULTS:** Seven eligible studies were identified and summary odds ratios (OR) were calculated using both random and quality effects models. Having more than ten actinic keratoses was most strongly associated with BCC, conferring up to a 5-fold increase in risk (OR: 4.97; 95% CI: 3.26, 7.58). Other factors, including solar elastosis, solar lentigines, and telangiectasia had weaker but positive associations with BCC with ORs around 1.5.

**CONCLUSIONS:** Markers of chronic photo-damage are positively associated with BCC. The presence of actinic keratoses was the most strongly associated with BCC of the markers examined.

**IMPACT:** This work highlights the relatively modest association between markers of chronic ultraviolet exposure and BCC.

## INTRODUCTION

Basal cell carcinoma (BCC) is the most commonly diagnosed cancer in Caucasian populations. BCC is not routinely captured in registries, but it is nevertheless clear that the incidence rates of BCC vary significantly around the world, ranging from two in east Asia to 1600/100,000/year in Queensland, Australia (1-7). The incidence rates appear to be continuing to rise in several populations (8-11). Although mortality from BCC is low they can cause considerable morbidity. Arguably the greatest burden, however, is on health system budgets. The total annual cost of managing keratinocyte cancers has been estimated to be about 500 million dollars in Australia (12), and since BCCs account for more than 70% of all keratinocyte cancers (13), they are likely to account for a large proportion of this total cost. The personal and health system costs of BCC emphasise the need for scientific inquiry into better understanding the causes of these common cancers.

Ultraviolet radiation (UVR) is the major aetiological agent in the pathogenesis of BCC (14-16). However, while comparisons of BCC incidence between fair-skinned populations living in regions with different ambient UVR highlight the importance of sun exposure, studies of personal sun exposure suggest lower levels of risk, with typically less than a doubling in risk with high self-reported levels of sun exposure (17). There is a plateau of BCC incidence with age and a reportedly stronger association with recreational than with occupational or total exposure (15), and it has been suggested that childhood might represent a critically important exposure window (14). In addition, while BCC occurs predominantly on the head and neck, approximately 25% arise on less sun exposed body sites such as the trunk (18-21). This is in contrast with squamous cell carcinoma (SCC) where only 8% of lesions occur on the trunk (18) and the strongest risks appear to result from a lifetime of high sun exposure (22).

Although epidemiological studies have tended to downplay the role of chronic cumulative sun exposure in the aetiology of BCC, most studies have relied largely on self-report to capture lifetime sun exposure, which is far from ideal. Sun exposure is ubiquitous and its

recall is prone to non-differential misclassification, shown by poor repeatability (23). Differential reporting according to skin cancer history is of potential concern in case-control studies. Additionally, report of exposure to the sun does not necessarily capture the relevant dose at a cellular level due to differences in skin type. There are several markers of chronic sun exposure, including actinic keratosis (24), elastosis, telangiectasia, and solar lentigines (25-28) and assessment of the association between these and BCC provides a more objective measure of the impact of cumulative UVR exposure. While there are studies that have explored this issue, there has been no previous attempt to synthesize this literature. We therefore conducted a systematic review and meta-analysis to better understand the impact of chronic UVR-related skin changes on the risk of BCC.

## **MATERIALS AND METHODS**

### **Literature search**

Eligible studies published from 1965 up to January 2013 were identified by computerised literature searching using the MEDLINE, Cumulative Index to Nursing and Allied Health Literature (CINAHL), Embase, Web of Knowledge, and Cochrane databases, and hand-searching of the reference lists of the retrieved articles.

For computer searches, we used the following MeSH terms or text words (using both the United Kingdom, the United States spellings and plurals where applicable): "Carcinoma, Basal Cell"[Mesh], "Basal Cell Carcinomas", "Basal Cell Epithelioma", "BCC", "Nonmelanoma skin cancer", "Non-melanoma skin cancer", "Non melanoma skin cancer", "NMSC", "Keratosis, Actinic"[Mesh], "solar keratosis", "Elastosis", "lentigines", "lentigo", "Telangiectasis"[Mesh], and "Telangiectasia".

### **Study selection**

We included observational studies of case-control and cohort designs in the meta-analysis provided that they permitted quantitative assessment of the association between

histologically-confirmed BCC and actinic keratoses (AK), solar elastosis, solar lentigines, and telangiectasia. Studies published in languages other than English were not eligible (29, 30). A single reviewer (MK) read the abstracts of all identified studies and excluded those that were clearly not relevant. He subsequently read the full texts of the remaining articles to determine if they met the study inclusion criteria. Where multiple reports from one study were found, the most recent or most complete publication was used. We did not exclude any studies from the analysis because of study quality, as quality was adjusted for in our quantitative model.

### **Data extraction**

After initial review of the abstracts identified by the primary computerised literature search we identified 70 potentially eligible studies, and hand-searched the reference lists of these. After review we excluded 63 studies because no photo-damage marker was reported ( $n = 37$ ), they were not independent of other included studies ( $n = 18$ ), were not in English ( $n = 2$ ) (29, 30), they reported combined data for BCC, SCC or AK together ( $n = 5$ ) (31-35), or the study was conducted in a high risk group ( $n = 1$ ) (36). We retrieved seven articles for further assessment, all of which met the eligibility criteria: one cohort study and six case-control studies. Of the eligible case-control studies, one was population-based, and five were clinic/hospital-based (Table 1).

Reported data for any relevant variable for which analysis was performed were extracted.

We also extracted a wide range of variables that may influence study validity or contribute to heterogeneity. These included country, year of publication, study design, sample size, case definitions, method of case ascertainment, whether the analysis was adjusted for age, sex, time spent outdoors or skin color/type and results (RR, OR and 95% CIs).

### **Quality assessment**

We used a quality scoring checklist (Supplementary table) which included 16 questions for case-control studies and 13 questions for cohort studies based on a published template for

assessing observational studies (37). We specifically tailored it to our meta-analysis to rate the methodological quality of individual studies (by including such items as method of assessment (observed or self-reported) and whether the variable was adjusted for time spent outdoors or skin colour/type). A single scorer (MK) did the scoring. The quality score was calculated as the total score divided by the number of items (16 for case-control studies and 13 for cohort studies). Normalized scores ranged from 0 (worst) to 1 (best). A summary of the quality score for each paper is shown in Table 1.

### **Statistical analysis**

The primary outcome was the odds of BCC in those with the marker of interest compared to those without the marker. We preferentially used adjusted odds ratios (OR) reported by the authors, only using crude estimates if no adjusted estimates were reported. One cohort study reported relative risks (RRs) rather than ORs – we converted these to ORs using the OR2RR online software ([http://epigear.com/index\\_files/or2rr.html](http://epigear.com/index_files/or2rr.html)). Most studies presented data as the marker of interest present versus absent. For those that presented data in more categories we combined the ORs, weighting for the size of the category, to enable a meta-analysis of the odds of present versus absent.

Heterogeneity was determined to be present when the value of  $\tau^2$  was greater than zero and/or the  $Q$ -statistic was significant at  $P < 0.1$  (38). Although the standard approach for handling heterogeneity between studies is to use the random effects model (39), the present study uses bias adjustment via the quality effects model (40, 41). When a meta-analysis is performed in studies with large heterogeneity, the confidence intervals (CI) are substantially wider in random effects models than in quality effects models (41, 42). We have reported the random effects results for comparative purposes. Due to the small number of studies, we did not conduct analyses stratified according to potential sources of heterogeneity including study design, sources of cases and controls, study location, and adjustment for potential confounders. We performed leave-one-out sensitivity analyses in which we omitted each

study in turn to determine whether the results could have been influenced significantly by any particular study.

Publication bias was examined visually by funnel plots as well as with the aid of the Begg rank correlation method and the Egger regression method (43, 44). We considered the funnel plot to be asymmetric if the intercept on Egger's regression deviated from zero with a P value of less than 0.1. All analyses were carried out using MetaXL ([www.epigear.com](http://www.epigear.com)).

## **RESULTS**

### **Actinic keratosis**

All seven studies presented data on the association between AK and BCC (Table 2). Compared to people with no AK, those with AK were at a markedly higher risk of BCC (Quality effects model OR: 3.28; 95% CI: 2.71, 3.96) (Figure 1a). We also assessed the effect of having a greater number of AKs by meta-analysing the three studies that presented OR for different categories of AK count (45-47). There was evidence of a dose-response, with the OR increasing from 2.22 (95% CI: 1.71, 2.89) in those with fewer than 10 AKs to 4.97 (95% CI: 3.26, 7.58) in people with at least 10 AKs. There was no heterogeneity in risk estimates across studies and the pooled ORs were approximately similar when we used a random effects model (Table 2).

### **Solar elastosis**

Five studies presented data on the association between solar elastosis and BCC (Table 2). Having solar elastosis was associated with an increased risk of BCC (quality effects model OR: 1.58; 95% CI: 1.04, 2.40) (Figure 1b). In one study the presence of solar elastosis was associated with reduced risk of BCC (OR: 0.1, 95% CI: 0.05, 0.42) (48). There was significant heterogeneity across studies but no individual study appeared to account for this. Even taking out the study with the paradoxical result did not remove the heterogeneity. The



pooled OR using the random effects model was considerably lower and no longer significant (random effects model OR: 1.13; 95% CI: 0.52, 2.45).

### **Solar lentigines**

Seven studies presented data on solar lentigines and BCC risk (Table 2). The presence of solar lentigines was associated with an approximately 1.5-fold increased risk of BCC using the quality effects model (OR: 1.44; 95% CI: 1.10, 1.89) (Figure 1c), and a 1.7-fold increased risk using the random effects model (OR: 1.71; 95% CI: 1.17, 2.48) (Table 2). There was evidence of significant heterogeneity but no individual study appeared to account for this.

### **Telangiectasia**

Three studies presented data on the association between telangiectasia and BCC (Table 2). Compared to people with no telangiectasia, those with telangiectasia were at a higher risk of BCC (quality effects model OR: 1.59; 95% CI: 1.31, 1.93) (Figure 1d). The pooled ORs using quality effects and random effects models were the same, with no heterogeneity in risk estimates across studies.

### **PUBLICATION BIAS**

There was no evidence of publication bias using the Egger weighted regression method and the Begg rank correlation method for the analyses of all markers of interest with the exception of solar lentigines which showed significant publication bias with both methods (Table 3).

### **DISCUSSION**

This is the first study to systematically evaluate available epidemiologic evidence about the magnitude of the relationship between cutaneous markers of photo-damage and risk of BCC using estimates of OR derived through meta-analysis. The presence of AK was most strongly associated with BCC, with more than 10 AK conferring up to a 5-fold increase in risk. Other

photo-damage markers including solar elastosis, solar lentigines, and telangiectasia, had weaker but positive and similar associations with BCC, with ORs around 1.5.

This analysis is based on observational studies which were predominantly case-control in design. Although the markers of interest used in this study were not prone to recall bias, in most studies it was unclear if observers were blinded to skin cancer status prior to examination, potentially resulting in biased assessments (47-51). The most likely result of this would have been to overestimate the association between chronic skin damage and BCC. Only one cohort study (45) has included measures of actinic skin damage, so we were unable to estimate the potential for observer bias. The studies included in this analysis ranged in size from 127 cases to 528 cases and varied in quality. We took account of both size and quality in this analysis to derive the best estimates of association between actinic skin damage and BCC.

Our results demonstrate that markers of chronic sun exposure do increase the risk of BCC, suggesting that cumulative exposure is aetiologically relevant. However with the exception of AK the associations were modest and the strength of association was similar to that found for self-reported cumulative sun exposure. This is in contrast to SCC where the presence of AK increases risk by 30-40 times and telangiectasia and solar elastosis confer 3- and 6-fold increases in risk respectively (46, 52). These relatively lower associations with BCC might suggest that different patterns of sun exposure, such as intense exposure are important. It is also possible that the patterns of sun exposure that give rise to BCCs on different anatomic sites are different, and that chronic UVR is more important for lesions arising on sun exposed sites, with intermittent exposure underpinning the development of BCCs on the trunk (53).

To our knowledge this is the first study to systematically evaluate available epidemiologic evidence about the magnitude of the relationship between photo-damage markers of the skin and risk of BCC using estimates of OR derived through meta-analysis. We used a very comprehensive search strategy and, although the number of studies was too small to enable

stratified analyses, we took account of variable study quality of studies. It is interesting to note that, despite the enormous burden of BCC, there are few studies that have used objective measures of sun exposure in risk assessment, and only one of these was prospective. In addition, while the proportion of BCCs arising on the trunk in the included studies ranged from 7% to 37% for the four studies (47-49, 51) that reported site distribution, there was no attempt to stratify by site or restrict the analyses to lesions arising on exposed sites, and this may have led to an underestimation of the role of chronic UVR in the aetiology of BCC arising on sites that are routinely sun exposed.

This meta-analysis has highlighted the relatively modest role of chronic UVR in the aetiology of BCC, suggesting that other factors must also be contributing to the marked differences observed at an ecological level. A better understanding of the patterns of sun exposure that are most important for BCC development, and the ways these interact with other factors such as genetic susceptibility, will enable the development of more strategic and informed strategies to reduce the impact of BCC on individuals and health systems.

## REFERENCES

1. Sng J, Koh D, Siong WC, Choo TB. Skin cancer trends among Asians living in Singapore from 1968 to 2006. *J Am Acad Dermatol* 2009;61(3): 426-32.
2. Birch-Johansen F, Jensen A, Mortensen L, Olesen AB, Kjaer SK. Trends in the incidence of nonmelanoma skin cancer in Denmark 1978-2007: Rapid incidence increase among young Danish women. *Int J Cancer* 2010;127(9): 2190-8.
3. Bath-Hextall F, Leonardi-Bee J, Smith C, Meal A, Hubbard R. Trends in incidence of skin basal cell carcinoma. Additional evidence from a UK primary care database study. *Int J Cancer* 2007;121(9): 2105-8.
4. Demers AA, Nugent Z, Mihalcioiu C, Wiseman MC, Kliewer EV. Trends of nonmelanoma skin cancer from 1960 through 2000 in a Canadian population. *J Am Acad Dermatol* 2005;53(2): 320-8.
5. Jung GW, Metelitsa AI, Dover DC, Salopek TG. Trends in incidence of nonmelanoma skin cancers in Alberta, Canada, 1988-2007. *Br J Dermatol* 2010;163(1): 146-54.
6. Reizner GT, Chuang TY, Elpern DJ, Stone JL, Farmer ER. Basal cell carcinoma in Kauai, Hawaii: the highest documented incidence in the United States. *J Am Acad Dermatol* 1993;29(2 Pt 1): 184-9.
7. Staples MP, Elwood M, Burton RC, Williams JL, Marks R, Giles GG. Non-melanoma skin cancer in Australia: the 2002 national survey and trends since 1985. *Med J Aust* 2006;184(1): 6-10.
8. Ko CB, Walton S, Keczkcs K, Bury HP, Nicholson C. The emerging epidemic of skin cancer. *Br J Dermatol* 1994;130(3): 269-72.
9. Holme SA, Malinowszky K, Roberts DL. Changing trends in non-melanoma skin cancer in South Wales, 1988-98. *Br J Dermatol* 2000;143(6): 1224-9.
10. Doherty VR, Brewster DH, Jensen S, Gorman D. Trends in skin cancer incidence by socioeconomic position in Scotland, 1978-2004. *Br J Cancer* 2010;102(11): 1661-4.

11. Skellett AM, Hafiji J, Greenberg DC, Wright KA, Levell NJ. The incidence of basal cell carcinoma in the under-30s in the UK. *Clin Exp Dermatol* 2012;37(3): 227-229.
12. Fransen M, Karahalios A, Sharma N, English DR, Giles GG, Sinclair RD. Non-melanoma skin cancer in Australia. *Med J Aust* 2012 197(10): 565-8.
13. Diepgen TL, Mahler V. The epidemiology of skin cancer. *Br J Dermatol* 2002;146: 1-6.
14. Gallagher RP, Hill GB, Bajdik CD, Fincham S, Coldman AJ, McLean DI, et al. Sunlight exposure, pigmentary factors, and risk of nonmelanocytic skin cancer. I. Basal cell carcinoma. *Arch Dermatol* 1995;131(2): 157-63.
15. Kricger A, Armstrong BK, English DR, Heenan PJ. Does intermittent sun exposure cause basal cell carcinoma? A case-control study in Western Australia. *Int J Cancer* 1995;60(4): 489-94.
16. Rosso S, Zanetti R, Martinez C, Tormo MJ, Schraub S, Sancho-Garnier H, et al. The multicentre south European study 'Helios'. II: Different sun exposure patterns in the aetiology of basal cell and squamous cell carcinomas of the skin. *Br J Cancer* 1996;73(11): 1447-54.
17. Armstrong BK, Kricger A. The epidemiology of UV induced skin cancer. *J Photochem Photobiol B* 2001;63(1-3): 8-18.
18. NCCI Non-melanoma Skin Cancer Working Group. The 2002 national non-melanoma skin cancer survey. 2003; Melbourne: Carlton, National Cancer Control Initiative.
19. Scrivener Y, Grosshans E, Cribier B. Variations of basal cell carcinomas according to gender, age, location and histopathological subtype. *Br J Dermatol* 2002;147(1): 41-7.
20. Bastiaens MT, Hoefnagel JJ, Bruijn JA, Westendorp RG, Vermeer BJ, Bouwes Bavinck JN. Differences in age, site distribution, and sex between nodular and superficial basal cell carcinoma indicate different types of tumors. *J Invest Dermatol* 1998;110(6): 880-4.

21. Pelucchi C, Di Landro A, Naldi L, La Vecchia C. Risk factors for histological types and anatomic sites of cutaneous basal-cell carcinoma: an Italian case-control study. *J Invest Dermatol* 2007;127(4): 935-44.
22. Kennedy C, Bajdik CD, Willemze R, de Gruijl FR, Bouwes Bavinck JN. The Influence of Painful Sunburns and Lifetime Sun Exposure on the Risk of Actinic Keratoses, Seborrheic Warts, Melanocytic Nevi, Atypical Nevi, and Skin Cancer. *J Invest Dermatol* 2003;120(6): 1087-1093.
23. English DR, Armstrong BK, Kricger A. Reproducibility of reported measurements of sun exposure in a case-control study. *Cancer Epidemiol Biomarkers Prev* 1998;7(10): 857-863.
24. Vitasa BC, Taylor HR, Strickland PT, Rosenthal FS, West S, Abbey H, et al. Association of nonmelanoma skin cancer and actinic keratosis with cumulative solar ultraviolet exposure in Maryland watermen. *Cancer* 1990;65(12): 2811-2817.
25. Kennedy C, Bastiaens MT, Bajdik CD, Willemze R, Westendorp RGJ, Bouwes Bavinck JN. Effect of Smoking and Sun on the Aging Skin. *J Invest Dermatol* 2003;120(4): 548-554.
26. Lucas RM, Ponsonby A-L, Dear K, Taylor BV, Dwyer T, McMichael AJ, et al. Associations between Silicone Skin Cast Score, Cumulative Sun Exposure, and Other Factors in the Ausimmune Study: A Multicenter Australian Study. *Cancer Epidemiol Biomarkers Prev* 2009;18(11): 2887-2894.
27. Green AC, Hughes MCB, McBride P, Fournanier A. Factors Associated with Premature Skin Aging (Photoaging) before the Age of 55: A Population-Based Study. *Dermatology* 2011;222(1): 74-80.
28. Thomas NE, Kricger A, From L, Busam KJ, Millikan RC, Ritchey ME, et al. Associations of cumulative sun exposure and phenotypic characteristics with histologic solar elastosis. *Cancer Epidemiol Biomarkers Prev* 2010;19(11): 2932-41.

29. Ruiz Lascano A, Kuznitzky R, Garay I, Ducasse C, Albertini R. [Risk factors for basal cell carcinoma. Case-control study in Cordoba]. *Medicina (B Aires)* 2005;65(6): 495-500.
30. Sanchez G, Nova J, de la Hoz F. Risk factors for basal cell carcinoma: A study from the National Dermatology Center of Colombia. *Actas Dermosifiliogr* 2012;103 (4): 294-300.
31. Rocha FP, Menezes A, Almeida Junior HL, Tomasi E. Risk markers and risk factors for actinic keratosis and basal cell carcinoma: a case-control study. *An Bras Dermatol* 2004;79(4): 441-454.
32. Healy E, Collins P, Barnes L. Nonmelanoma skin cancer in an Irish population: an appraisal of risk factors. *Ir Med J* 1995;88(2): 58-9.
33. Ferreira FR, Nascimento LFC, Rotta O. Fatores de risco para câncer da pele não melanoma em Taubaté, SP: um estudo caso-controle. *Rev Assoc Med Bras* 2011;57: 431-437.
34. Lichte V, Dennenmoser B, Dietz K, Häfner HM, Schlagenhauff B, Garbe C, et al. Professional risk for skin cancer development in male mountain guides – a cross-sectional study. *J Eur Acad Dermatol Venereol* 2010;24(7): 797-804.
35. Herity B, O'Loughlin G, Moriarty MJ, Conroy R. Risk factors for non-melanoma skin cancer. *Ir Med J* 1989;82(4): 151-2.
36. Liboutet M, Portela M, Delestaing G, Vilmer C, Dupin N, Gorin I, et al. MC1R and PTCH gene polymorphism in French patients with basal cell carcinomas. *J Invest Dermatol* 2006;126(7): 1510-7.
37. Shamliyan TA, Kane RL, Ansari MT, Raman G, Berkman ND, Grant M, et al. Development quality criteria to evaluate nontherapeutic studies of incidence, prevalence, or risk factors of chronic diseases: pilot study of new checklists. *J Clin Epidemiol* 2011;64(6): 637-657.

38. Takkouche B, Cadarso-Suárez C, Spiegelman D. Evaluation of Old and New Tests of Heterogeneity in Epidemiologic Meta-Analysis. *Am J Epidemiol* 1999;150(2): 206-215.
39. DerSimonian R, Laird N. Meta-analysis in clinical trials. *Control Clin Trials* 1986;7(3): 177-88.
40. Doi SA, Barendregt JJ, Mozurkewich EL. Meta-analysis of heterogeneous clinical trials: An empirical example. *Contemp Clin Trials* 2011;32(2): 288-298.
41. Doi SA, Thalib L. A quality-effects model for meta-analysis. *Epidemiology* 2008;19(1): 94-100.
42. Doi SA, Thalib L. An alternative quality adjustor for the quality effects model for meta-analysis. *Epidemiology* 2009;20(2): 314.
43. Begg CB, Mazumdar M. Operating Characteristics of a Rank Correlation Test for Publication Bias. *Biometrics* 1994;50(4): 1088-1101.
44. Egger M, Smith GD, Schneider M, Minder C. Bias in meta-analysis detected by a simple, graphical test. *BMJ* 1997;315(7109): 629-634.
45. Green A, Battistutta D, Hart V, Leslie D, Weedon D. Skin cancer in a subtropical Australian population: incidence and lack of association with occupation. The Nambour Study Group. *Am J Epidemiol* 1996;144(11): 1034-40.
46. Kricger A, Armstrong BK, English DR, Heenan PJ. Pigmentary and cutaneous risk factors for non-melanocytic skin cancer--a case-control study. *Int J Cancer* 1991;48(5): 650-62.
47. Dessinioti C, Tzannis K, Sypsa V, Nikolaou V, Kypreou K, Antoniou C, et al. Epidemiologic risk factors of basal cell carcinoma development and age at onset in a Southern European population from Greece. *Exp Dermatol* 2011;20(8): 622-6.
48. Walther U, Kron M, Sander S, Sebastian G, Sander R, Peter RU, et al. Risk and protective factors for sporadic basal cell carcinoma: results of a two-centre case-control



- study in southern Germany. Clinical actinic elastosis may be a protective factor. *Br J Dermatol* 2004;151(1): 170-8.
49. Naldi L, DiLandro A, D'Avanzo B, Parazzini F. Host-related and environmental risk factors for cutaneous basal cell carcinoma: evidence from an Italian case-control study. *J Am Acad Dermatol* 2000;42(3): 446-52.
50. Gon A, Minelli L. Risk factors for basal cell carcinoma in a southern Brazilian population: a case-control study. *Int J Dermatol* 2011;50(10): 1286-90.
51. Corona R, Dogliotti E, D'Errico M, Sera F, Iavarone I, Baliva G, et al. Risk factors for basal cell carcinoma in a Mediterranean population: role of recreational sun exposure early in life. *Arch Dermatol* 2001;137(9): 1162-8.
52. English DR, Armstrong BK, Kricger A, Winter MG, Heenan PJ, Randell PL. Demographic characteristics, pigmentary and cutaneous risk factors for squamous cell carcinoma of the skin: A case-control study. *Int J Cancer* 1998;76(5): 628-634.
53. Neale RE, Davis M, Pandeya N, Whiteman DC, Green AC. Basal cell carcinoma on the trunk is associated with excessive sun exposure. *J Am Acad Dermatol* 2007;56(3): 380-386.

**Table 1. Characteristics of the 7 studies included in the meta-analysis of cutaneous markers of photo-damage and risk of basal cell carcinoma of the skin.**

First Author	Year of publication	Study Location	Source of cases	Source of controls	Sample size	Actinic keratosis	Solar elastosis	Solar Lentigines	Telangiectasia	Quality score
<i>Cohort</i>					<i>Cases/Cohort</i>					
<b>Green A (45)</b>	1996	Australia	population	population	250/2095	✓	✓	✓	✓	0.84
<i>Population-based case-control</i>					<i>Cases/Controls</i>					
<b>Kricker S (46)</b>	1991	Australia	population	population	226/1021	✓	✓	✓	✓	0.81
<i>Clinic/Hospital-based case-control</i>					<i>Cases/Controls</i>					
<b>Dessinioti C (47)</b>	2011	Greece	Clinic/Hospital	Clinic/Hospital	199/200	✓		✓		0.75
<b>Gon A (50)</b>	2011	Brazil	Clinic/Hospital	Clinic/Hospital	127/280	✓	✓	✓	✓	0.69
<b>Walther U (48)</b>	2004	Germany	Clinic/Hospital	Clinic/Hospital	213/411	✓	✓	✓		0.56
<b>Corona R (51)</b>	2001	Italy	Clinic/Hospital	Clinic/Hospital	166/158	✓	✓	✓		0.69
<b>Naldi L (49)</b>	2000	Italy	Clinic/Hospital	Clinic/Hospital	528/512	✓		✓		0.81

**Table 2. Meta-analysis results using the quality effects and random effects models: Risk of basal cell carcinoma associated with cutaneous markers of photo-damage.**

	Studies	Quality effects model OR (95% CI)	Random effects model OR (95% CI)	<i>I</i> <sup>2</sup> (%)	<i>P</i> het
<b>Actinic keratosis</b>					
Present vs. Absent	7	3.28 (2.71, 3.96)	3.19 (2.57, 3.95)	33.0	0.17
Less than 10	3	2.22 (1.71, 2.89)	2.26 (1.71, 2.99)	20.0	0.28
More than 10	3	4.97 (3.26, 7.58)	4.96 (2.83, 8.69)	67.0	0.05
<b>Solar elastosis</b>					
Present vs. Absent	5	1.58 (1.04, 2.40)	1.13 (0.52, 2.45)	90.0	0.00
<b>Solar lentigines</b>					
Present vs. Absent	7	1.44 (1.10, 1.89)	1.71 (1.17, 2.48)	85.0	0.00
<b>Telangiectasia</b>					
Present vs. Absent	3	1.59 (1.31, 1.93)	1.59 (1.31, 1.93)	00.0	0.90

**Table 3. Results of publication bias tests using the Egger weighted regression method and the Begg rank correlation method.**

	<i>P</i> for bias	
	Begg method	Egger method
<b>Actinic keratosis</b>		
Present vs. Absent	0.764	0.696
Less than 10	0.296	0.020
More than 10	0.52	0.841
<b>Solar elastosis</b>		
Present vs. Absent	0.221	0.344
<b>Solar lentigines</b>		
Present vs. Absent	0.035	0.029
<b>Telangiectasia</b>		
Present vs. Absent	1.000	0.605

## Figure legends:

**Figure 1 (a-d):** Forest plots of the association between actinic keratosis (Figure 1a), solar elastosis (Figure 1b), solar lentigines (Figure 1c), telangiectasia (Figure 1d), and BCC using quality effects models. Each line represents an individual study result with the width of the horizontal line indicating 95% CI, the position of the box representing the odds ratio, and the size of the box being proportional to the weight of the study.

