

Bond University
Research Repository



What are the optimal dietary strategies for the management of cancer-related nausea and vomiting in adults: A PEN guideline update

Crichton, Megan; Marx, Wolfgang; Marshall, Skye; Isenring, Elisabeth

Published in:
Nutrition and Dietetics

DOI:
[10.1111/1747-0080.12567](https://doi.org/10.1111/1747-0080.12567)

Licence:
Free to read

[Link to output in Bond University research repository.](#)

Recommended citation(APA):
Crichton, M., Marx, W., Marshall, S., & Isenring, E. (2019). What are the optimal dietary strategies for the management of cancer-related nausea and vomiting in adults: A PEN guideline update. *Nutrition and Dietetics*, 76(S1), 77. Article 306. <https://doi.org/10.1111/1747-0080.12567>

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

For more information, or if you believe that this document breaches copyright, please contact the Bond University research repository coordinator.

WHAT ARE THE OPTIMAL DIETARY STRATEGIES FOR THE MANAGEMENT OF CANCER-RELATED NAUSEA AND VOMITING IN ADULTS: A PEN GUIDELINE UPDATE

Megan Crichton¹, Wolfgang Marx², Skye Marshall¹ and Elizabeth Isenring¹

¹ Bond University Nutrition and Dietetics Research Group

² Deakin University Food and Mood Centre

The symptoms of nausea and vomiting are primarily managed with anti-emetic medications in cancer patients; however, dietary interventions and nutraceuticals are often recommended by health professionals. Therefore, a literature review was undertaken to update the PEN guideline and toolkit. To identify existing dietary recommendations and determine their rigor for controlling cancer-related nausea and vomiting, literature was searched using three electronic databases. The quality of evidence was graded using PEN Evidence Grading Checklist. Anecdotal reports and expert opinion form the basis for many dietary modifications such as eating bland foods and consuming small meals and snacks regularly (Grade of Evidence:C-Limited). Dietary counselling during and up to three months after radiation treatment may decrease severity of nausea and vomiting for patients with colorectal and head and neck cancers (Grade of Evidence: C-Limited). The use of ginger supplements appears to be safe for chemotherapy patients, and may provide some benefit in combination with standard anti-emetics for chemotherapy-induced nausea and vomiting as well as fatigue and quality of life (Grade of Evidence:B-Fair). Managing taste changes may be beneficial in relieving nausea and vomiting (Grade of Evidence:C-Limited). Habitual alcohol intake is related to lower incidence of nausea and vomiting; however, adherence to national alcohol guidelines and/or individualised advice from treating specialist doctors is recommended (Grade of Evidence:C-Limited). Updated patient education materials were produced. Although diet is generally not a first line of therapy, dietary and nutraceutical interventions may provide additional benefits when used in conjunction with anti-emetics; however, existing literature is limited, warranting further research.