

Bond University
Research Repository



Clever searching for evidence: New search filters can help to find the needle in the haystack

Sanders, Sharon; Del Mar, Chris

Published in:
British Medical Journal

DOI:
[10.1136/bmj.330.7501.1162](https://doi.org/10.1136/bmj.330.7501.1162)

Licence:
Other

[Link to output in Bond University research repository.](#)

Recommended citation(APA):
Sanders, S., & Del Mar, C. (2005). Clever searching for evidence: New search filters can help to find the needle in the haystack. *British Medical Journal*, 330(7501), 1162-1163. <https://doi.org/10.1136/bmj.330.7501.1162>

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

For more information, or if you believe that this document breaches copyright, please contact the Bond University research repository coordinator.

5-21-2005

Clever searching for evidence

Sharon Sanders
University of Queensland

Chris Del Mar
Bond University, chris_del_mar@bond.edu.au

Follow this and additional works at: http://epublications.bond.edu.au/hsm_pubs

Recommended Citation

Sanders, Sharon and Del Mar, Chris, "Clever searching for evidence" (2005). *Faculty of Health Sciences & Medicine Publications*. Paper 5. http://epublications.bond.edu.au/hsm_pubs/5

This Editorial is brought to you by the Faculty of Health Sciences & Medicine at [ePublications@bond](mailto:epublications@bond). It has been accepted for inclusion in Faculty of Health Sciences & Medicine Publications by an authorized administrator of [ePublications@bond](mailto:epublications@bond). For more information, please contact [Bond University's Repository Coordinator](#).



Clever searching for evidence

Sharon Sanders and Chris Del Mar

BMJ 2005;330;1162-1163
doi:10.1136/bmj.330.7501.1162

Updated information and services can be found at:
<http://bmj.com/cgi/content/full/330/7501/1162>

- These include:*
- Data supplement** *"Extra references"*
<http://bmj.com/cgi/content/full/330/7501/1162/DC1>
- References** This article cites 12 articles, 4 of which can be accessed free at:
<http://bmj.com/cgi/content/full/330/7501/1162#BIBL>
1 online articles that cite this article can be accessed at:
<http://bmj.com/cgi/content/full/330/7501/1162#otherarticles>
- Rapid responses** 2 rapid responses have been posted to this article, which you can access for free at:
<http://bmj.com/cgi/content/full/330/7501/1162#responses>
You can respond to this article at:
<http://bmj.com/cgi/eletter-submit/330/7501/1162>
- Email alerting service** Receive free email alerts when new articles cite this article - sign up in the box at the top right corner of the article
-
- Topic collections** Articles on similar topics can be found in the following collections
[Other Journalology](#) (363 articles)
[Other evidence based practice](#) (448 articles)
-

Notes

To order reprints of this article go to:
<http://www.bmjournals.com/cgi/reprintform>

To subscribe to *BMJ* go to:
<http://bmj.bmjournals.com/subscriptions/subscribe.shtml>

period, covering acute presentation, management, rehabilitation, and longer term care. They emphasise the need for early imaging and input from a paediatric neurology tertiary centre. They also appropriately emphasise that these children should be investigated for an underlying prothrombotic tendency and should undergo echocardiography.

With respect to acute care, the working group found no studies specifically examining the efficacy of acute treatments and made its recommendations based on consensus opinion. In the main, whether or not to anticoagulate a child with cardiac embolism is therefore a decision that should be based on the individual patient. Similarly no current evidence supports the use of thrombolytic agents. By contrast good evidence exists for the use of aspirin in a dose of 5 mg per kg body weight per day unless imaging suggests intracranial haemorrhage or the child has sickle cell disease.

Alongside the acute medical (and more rarely surgical) management, the guidelines recommend that at an early stage the child's disability be assessed and that acute clinicians liaise with community child health services.

It is disappointing but unsurprising that specific rehabilitative techniques and approaches that are recommended do not have an evidence base: the information that is available is largely derived from practice and experience in children with cerebral palsy.

The guidelines tackle longer term rehabilitation needs, emphasising the requirement for psychological, cognitive, and family assessments together with the likelihood that rehabilitation for many children will extend into their adult lives.

Against the background of a paucity of hard data, the working group has made various audit recommendations that relate to acute care and longer term care of children who have had a stroke.

A working group report such as this raises issues around paediatric disability that go beyond its remit. These are both general and specific. The general point

to be made is that models for assessment and rehabilitation in stroke have much in common with those for most neurologically disabling conditions in childhood. These include prompt and appropriate diagnostic formulation, comprehensive assessment to include an understanding of family functioning and dynamics, appropriate health interventions at an early stage, and realistic multidisciplinary rehabilitation based on evidence based practice wherever that is possible. Hence one would wish to see paediatric stroke services integrated into and as a component of wider paediatric neurodisability services.

Specifically, what is disappointing about the deliberations of the working group is that they have not addressed the issues that relate to perinatal stroke and its usual sequel of congenital hemiplegia. Lynch et al have summarised recent knowledge of this subject.³ The diagnostic issues in this group of children are at least as complex as when stroke has its origin later in childhood, and specifically the links with intrapartum adversity are a current subject of interest and potential litigation.⁴ Expertise in obstetric, fetal, and neonatal medicine will need to be added to any future working group if these wider aspects of stroke in childhood are to be addressed satisfactorily.

Lewis Rosenbloom *consultant paediatric neurologist*

Alder Hey Children's Hospital, Liverpool L3 0BQ
(83waterloo@supanet.com)

Competing interests: None declared.

- 1 Royal College of Physicians, London. Intercollegiate Stroke Working Party. June 2004. *National clinical guidelines for stroke*. 2nd ed. www.rcplondon.ac.uk/pubs/books/stroke/stroke_guidelines_2ed.pdf (accessed 15 Apr 2005).
- 2 Royal College of Physicians, London. Paediatric Stroke Working Group. Stroke in childhood. *Clinical guidelines for diagnosis, management and rehabilitation*. November 2004. www.rcplondon.ac.uk/pubs/books/childstroke/childstroke_guidelines.pdf (accessed 15 Apr 2005).
- 3 Lynch, JK, Hirtz DG, DeVeber G, Nelson KB. Report of the National Institute of Neurological Disorders and Stroke workshop on perinatal and childhood stroke. *Pediatrics* 2002;109:116-23.
- 4 Cowan F, Rutherford M, Groenendaal F, Eken P, Mercuri E, Bydder GM. Origin and timing of brain lesions in term infants with neonatal encephalopathy. *Lancet* 2003;361: 736-42.

Clever searching for evidence

New search filters can help to find the needle in the haystack

Do people who write about and propagate evidence based medicine use its principles properly? A paper in this week's *BMJ* and two other recent *BMJ* papers report on such authors' abilities to find the best evidence.¹⁻³ On the way, they have provided excellent search strategies, using filters they call "hedges" (as in hedging one's bets) that help to separate the wheat (scientifically strong studies of diagnosis, treatment, and systematic reviews) from the chaff (less rigorous ones) in one of the most frequently accessed medical literature databases, Medline.

Why is this important? We still often need to search large databases such as Medline to find original research data because reviews may not cover our questions, may be out of date, and may not be relevant

enough to real clinical problems. Databases of primary research are staggeringly large (there are more than 12 million citations in Medline, and 7 million in Embase). Most research papers are written as communications from scientist to scientist and relatively few have immediate clinical relevance. Most of the remainder are not rigorous enough to warrant applying clinically.⁴ Subsequently, the proportion of useful information is very small.^{w1}

This lack of high quality, clinically relevant research studies leads to insecurity about the results of literature searches (table). Finding evidence can often seem easy,

 Additional references w1-w5 are on bmj.com

Papers p 1179

Lack of security about the results of a search

Search	Evidence	
	Positive	Negative
Positive	Success	Suboptimal evidence (potentially misleading)
Negative	Uncertainty—is there really no evidence?	Uncertainty—or did we just not find it?

but those searching may either be blissfully unaware that they have not found the best evidence, or might suspect that there is still better to be found.⁵ But when a search retrieves nothing worth while, was it because there is none, or is it that the person looking just did not find it?

Skills limitations—both perceived and real—are important barriers to effective searching.⁵ Efforts have been made to simplify use of the large databases by allowing searchers to search with simple statements and minimal knowledge of the technicalities of the way the database works.^{w2} But, until this approach has been found effective in pragmatic trials, we have to deal with some associated consequences. Simple searching using free text can give the impression that searching in the medical literature databases is easy and can be performed in the same manner as a general internet search. But this is wrong.^{w3} We still have to know how to formulate a search in each particular database. Even the best filters¹⁻³ cannot compensate for a poorly structured content search strategy.

For example, in Pubmed a searcher may enter the term “Myocardial infarct*”, thinking that the truncation will improve their search by retrieving articles using the terms myocardial infarction, myocardial infarctions, and myocardial infarcts. But the truncation (*) removes the automatic term mapping feature, which searches for articles using both the text word and MeSH term, reducing retrieval and potentially missing articles.

Furthermore, many clinicians do not search at all^{6,7,w4} or look for answers in easily located but less evidence based sources.^{7,8} To delay clinical activity to look for the best evidence is not even considered by most doctors. There seems to be a “90 second rule”—that is, clinicians don’t even try to find information unless they think they can do it in a minute and a half.^{9,10} Yet, with practice and access to certain sources, this is possible during routine ward rounds.¹¹

Searching is only one necessary skill. Whenever we run evidence based practice workshops, we are struck by the overemphasis that clinicians place on searching for and accessing evidence and their dismay at finding they also need to acquire other skills—asking an initial useful question, appraising, and applying evidence—as well.

Is using evidence in these ways too hard, and a waste of time for people who are only ever going to be amateurs, and who could refer their questions to experts? Is this analogous to pathology and medical imaging where clinicians need to know the principles of how investigations are done, but do not have to do the tests themselves?

Ideally, a literature searching service would be woven into the medical record, or made available at

some point even closer to the clinical decision. Although this is still a dream for most clinicians, literature search services are available that provide evaluated and summarised evidence, and answer specific questions.¹² Efforts to nationally coordinate and develop these services are currently under way in the United Kingdom.^{w5}

Another alternative to clinicians searching is to use compendiums of the evidence. An increasing number of excellent resources is available that either synthesise primary research (such as the *Cochrane Library*), or summarise it (such as *Clinical Evidence*, *Bandolier*, *TRIP Database Plus*, and evidence based clinical guidelines). Valuable sources of abstracted and appraised clinically relevant primary studies (such as *Best Evidence*) are also available. But these abstracted services are not always easy to access, tend to focus on interventions rather than diagnosis or prognosis, are more prone to being out of date—and are now so prolific that they, in turn, have to be searched for and appraised.

Clinicians risk missing something important if they do not dip their fingers into the mess of searching for, finding, and using the best evidence, at least for some of their questions. Being able to weigh the evidence in one’s hand, to first feel the heft of it, could lead to a greater ability to use it well. Using the evidence in processed form potentially loses so much in understanding issues of quality, the quantity of an effect, and generalisability. Finding the needle will then be only a part of the battle with the haystack.

Sharon Sanders *senior research officer*

Discipline of General Practice, University of Queensland, Herston, QLD 4006, Australia
(s.sanders@uq.edu.au)

Chris Del Mar *dean, Health Sciences and Medicine*

Bond University, Gold Coast, QLD 4229 Australia
(cdelmar@bond.edu.au)

Competing interests: None declared.

- Haynes RB, McKibbin KA, Wilczynski NL, Walter SD, Werre SR. Optimal search strategies for retrieving scientifically strong studies of diagnosis from Medline: analytical survey. *BMJ* 2005;330:1179–82.
- Haynes RB, Wilczynski NL. Optimal search strategies for retrieving scientifically strong studies of diagnosis from Medline: analytical survey. *BMJ* 2004;328:1040–4.
- Montori VM, Wilczynski NL, Morgan D, Haynes RB. Optimal search strategies for retrieving systematic reviews from Medline: analytical survey. *BMJ* 2005;330:68–73. (8 January).
- Haynes B, Glasziou P, Straus S. Advances in evidence-based information resources for clinical practice. *ACP J Club* 2000;132:A11–4.
- Green ML, Ruff TR. Why do residents fail to answer their clinical questions? A qualitative study of barriers to practicing evidence-based medicine. *Acad Med* 2005;80:176–82.
- Gorman PN, Helfand M. Information seeking in primary care: how physicians choose which clinical questions to pursue and which to leave unanswered. *Med Decis Making* 1995;15:113–9.
- Green ML, Ciampi MA, Ellis PJ. Residents’ medical information needs in clinic: are they being met? *Am J Med* 2000;109:218–23.
- Montori VM, Tabini CC, Ebbert JO. A qualitative assessment of 1st-year internal medicine residents’ perceptions of evidence-based clinical decision making. *Teach Learn Med* 2002;14:114–8.
- Ramos K, Linscheid R, Schafer S. Real-time information-seeking behavior of residency physicians. *Fam Med* 2003;35:257–60.
- Ely JW, Osheroff JA, Ebell MH, Bergus GR, Levy BT, Chambliss ML, et al. Analysis of questions asked by family doctors regarding patient care. *BMJ* 1999;319:358–61.
- Sackett DL, Straus SE. Finding and applying evidence during clinical rounds: the “evidence cart.” *JAMA* 1998;280:1336–8.
- Del Mar CB, Silagy CA, Glasziou P, Weller D, Spinks AB, Bernath V, et al. Feasibility of an evidence-based literature search service for general practitioners. *Med J Aust* 2001;175:134–7.