

Bond University
Research Repository



Effects of combined strength and sprint training on lean mass, strength, power and sprint performance in masters road cyclists

Del Vecchio, Luke; Stanton, Robert; Reaburn, Peter; Macgregor, Campbell; Meerkin, Jarrod; Villegas, Jerome; Korhonen, Marko T

Published in:
Journal of Strength and Conditioning Research

DOI:
[10.1519/JSC.0000000000001960](https://doi.org/10.1519/JSC.0000000000001960)

Licence:
Other

[Link to output in Bond University research repository.](#)

Recommended citation(APA):
Del Vecchio, L., Stanton, R., Reaburn, P., Macgregor, C., Meerkin, J., Villegas, J., & Korhonen, M. T. (2019). Effects of combined strength and sprint training on lean mass, strength, power and sprint performance in masters road cyclists. *Journal of Strength and Conditioning Research*, 33(1), 66-79.
<https://doi.org/10.1519/JSC.0000000000001960>

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

For more information, or if you believe that this document breaches copyright, please contact the Bond University research repository coordinator.

Effects of Combined Strength and Sprint Training on Lean Mass, Strength, Power and Sprint Performance in Masters Road Cyclists

Luke Del Vecchio¹, Robert Stanton¹, Peter Reaburn¹, Campbell Macgregor¹, Jarrod Meerkin², Jerome Villegas³ and Marko T. Korhonen⁴

¹School of Medical and Applied Sciences, Central Queensland University, Rockhampton, Australia

²Measureup, Sydney, Australia

³Sports Performance and Athletic Development Consultant, Canberra, Australia

⁴Gerontology Research Centre, Department of Health Sciences, University of Jyväskylä, Finland

All Correspondence to:

Luke Del Vecchio

School of Medical and Applied Sciences

Central Queensland University

Building 81/1.16, Bruce Highway

Rockhampton QLD 4700

Australia

Email: l.delvecchio@cqu.edu.au

Telephone Number: +61 0749232427

1 **Abstract**

2 Strength and sprint training exercises are integral part of training in many younger endurance
3 cyclists to improve cycling efficiency and sprinting ability. This study was undertaken to
4 examine whether muscle and performance characteristics could be improved in endurance-
5 trained masters cyclist by adding strength and sprint training stimuli into their training
6 regimen. Twenty five masters road cyclists were assigned to a combined strength and sprint
7 training group (CT; n=9, 53.5 ± 9.3 years), a sprint training group (ST, n=7, 49.4 ± 4.8 years)
8 or a control group (CG, n=9, 56.9 ± 8.6 years). Before and after the 12 week intervention,
9 whole body lean mass (WBLM), total lower limb lean mass (LLLM), countermovement jump
10 height (CMJ), peak isometric torque of quadriceps (QPT) and hamstring (HPT) muscles were
11 examined. For evaluation of sport-specific performance, 10 second sprint cycling peak power
12 (PP10), total 30 second work (TW), peak power output (PPO) and flying 200 meter time trial
13 performance (TT) were assessed. No pre-training differences were observed between CT, ST
14 and CG groups for any of the dependant variables. After training, a significant ($p<0.05$)
15 between group difference was observed in TW between CT and CG groups. A significant
16 effect of time ($p<0.05$) was observed for LLLM in CT and ST groups, and for TT in the CT
17 group. These results suggest including strength and sprint exercises in training can increase
18 lower limb lean mass and sprint performance in endurance trained masters road cyclists.
19 Further research is warranted to find out an ideal pattern of training to maintain aerobic
20 capabilities along with sprint performance in aging road cyclists.

21 **Key Words:** Masters Cyclists, Muscle Mass, Combined, Strength Training, Sprint Training

22

23

24

1 **Introduction**

2 Masters athletes are typically older than 35 years of age and systematically train for, and
3 compete in, organized forms of sport (32). Over recent years there has been a significant
4 increase in the number of masters athletes continuing to train and compete at high
5 performance levels within individual and multi-sport (duathlon, triathlon) endurance events
6 designed for masters athletes (38,22). Of the individual events, particularly road cycling is
7 becoming increasingly popular among masters athletes. For example, the number of
8 competitive masters road cyclists in Australia has grown from about 4,000 in 2013 to 10,000
9 in 2015 (4).

10 In younger cyclists, maximal strength and hypertrophy exercises (4-10 RM) has been shown
11 to increase cycling efficiency and power output at VO_{2max} (34, 42). There is also evidence
12 that various explosive strength training exercises are used in high-level road cyclists in order
13 to improve sprinting ability that is decisive factor in the finish and breaks in road cycling (30).
14 Although training-induced muscular hypertrophy and strength gains may slightly decrease
15 with age, due to factors such hormonal changes, the adaptive capacity could be maintained up
16 to very old age (20). Only few studies have addressed the effects of strength training on
17 cycling performance in masters endurance cyclists and older individuals (11, 23, 35). For
18 instance, Louis (23) reported an improvement in cycling efficiency, following 3 weeks of
19 hypertrophy training (70% of 1RM) in a group of masters road cyclists. In older non-athletes,
20 strength training (~80% of 1RM) has been shown to improve cycling peak power output (11).

21 Previous research has shown that an age-related decline in lean mass contributes to the age-
22 related declines in aerobic and anaerobic performance in both untrained older adults (12) and
23 masters athletes (32). Importantly, high-volume endurance training has been shown to lead
24 reduced muscle fiber size, muscle mass and reduced absolute power and force production in

1 both single fiber and whole-muscle level in masters long-distance runners (7, 20, 41). In
2 contrast, strength training has proved to be an effective countermeasure to maintain or
3 increase muscle mass and functional characteristics of masters endurance runners (31) and
4 masters sprint runners (10,33). However, the effectiveness of strength training to increase
5 lean mass in endurance-trained masters cyclists is currently unknown.

6 Recently, there has been growing interest in the effects of sprint training and it's specific
7 form, high intensity interval training (HIIT) as an alternative modality for increasing physical
8 performance and muscle mass in older adults (3,28). HIIT regimes are characterised by brief
9 repeated intense bursts of activity (e.g. 4-6 x 30s), at maximal intensities. In healthy older
10 men HIIT has been shown to increase lean muscle mass (28). In younger cyclists, HIIT
11 improves cycling performance including sprint performance (9). However, to the best of our
12 knowledge, limited studies to date have investigated the effect of HIIT on cycling
13 performance and lean mass in masters endurance cyclists.

14 Based on the available studies, it might be suggested that replacing a portion of endurance
15 training with a combination of strength and sprint training, may be beneficial to limit the age-
16 related decline in lean mass, strength, power and sprint performance. In terms of overall
17 cycling performance, sprint and / or strength training is important for a number of reasons.
18 First, increase in muscle strength can improve cycling efficiency. Second, leg power is
19 needed to accelerate rapidly during a breakaway attack and the sprint to the finish typical in
20 road racing. Third, leg strength and power are needed during hill climbing. The purpose of
21 this study was to examine the effect of a 12 week concurrent strength and sprint training
22 program on muscle and performance characteristics in male masters road cyclists. We
23 hypothesised that 12 weeks of concurrent strength and sprint cycling training, would
24 significantly increase lean mass, strength, power and sprint performance in already
25 endurance-trained cyclists

1 **Methods**

2 **Experimental approach to the problem**

3

4 It was hypothesised that concurrent strength and sprint cycling training added to regular
5 endurance cycling training would lead to a significant increase in lean body mass, muscular
6 strength and power, and sprint performance in master road cyclists. A parallel, three-group,
7 intervention (pre-post-test) experimental design was used. To investigate the possible effects
8 of CT on strength, power and sprint performance in master endurance cyclists, Dual Energy
9 X-Ray Absorptiometry (DXA) measures of WBLM, LLLM, CMJ, QPT and HPT were
10 examined. For evaluation of sport-specific performance, PP10, TW, PPO and TT were
11 measured before and after a 12 week intervention period. All subjects performed
12 familiarization trials before the testing days. We used as the independent variable, the group,
13 whereas the dependent variables were WBLM, LLLM, CMJ, QPT, HPT, PP10, TW, PPO
14 and TT.

15 **Participants**

16 The study was approved by the Central Queensland University Human Research Ethics
17 Committee. Twenty-five healthy male masters cyclists aged between 41 and 76 years with no
18 background of strength training were recruited and provided written informed consent. The
19 subjects were required to be involved in regular cycling training and/or road cycling
20 competition for a minimum of two years and to be achieving a minimum of eight hours of
21 endurance cycling training per week. All subjects underwent pre-exercise screening to ensure
22 they had no established cardiovascular, metabolic or respiratory disease nor signs or
23 symptoms of disease (29).

24 Random allocation of participants into training groups was not possible as the majority of
25 participants had both work and family commitments that limited their availability to
26 participate in the ST or CT programs. As a result, subjects were allocated to either a control

1 group (CG, n=10), sprint cycling group (ST, n=7) or concurrent strength and sprint cycling
2 training group (CT, n=10). For personal reasons, one participant from the CT group and one
3 subject from the CG group withdrew from the study, subsequently reducing the CT group to
4 nine participants (CT, n=9) and the control group to nine participants (CG, n=9). Subjects
5 were instructed not to change their diet or lifestyle over the experimental period. The physical
6 characteristics of each group are shown in Table 1.

7
8 Table 1 about here.

9 10 **Procedures**

11 Subjects attended the laboratory (22°C, 60% RH) following an overnight fast and did not
12 consume caffeine the morning of the test. All tests were carried out between 0700 and 0900
13 hours. Pre- and post-intervention testing included measures of anthropometry, DXA, jumping
14 performance on force-plate, peak isometric torque of quadriceps and hamstring muscle
15 groups, ten second sprint cycling peak power, total 30 second work and maximal aerobic
16 power on a cycle ergometer. The flying 200 meter time trial performance test was performed
17 at a local, outdoor cycling velodrome. Twenty-five masters road cyclists, engaged in the same
18 endurance training program were assigned to one of the following three groups: concurrent
19 strength and sprint cycling training group (CT), sprint cycling training group (ST) and a
20 control group (CG). The CT group replaced four (50%) of their usual endurance cycling
21 sessions (table 3) with two strength training sessions and two sprint training sessions, the ST
22 group replaced two of their usual endurance cycling sessions with two sprint training
23 sessions; and the CG group maintained their normal endurance training.

24 **1. Body composition**

25 Stature (m) and body mass (kg) were measured with a stadiometer and medical scales (Seca,
26 Birmingham, UK) with participant's unshod and wearing cycling apparel. Dual Energy X-

1 Ray Absorptiometry (DXA) (Hologic Discovery-W, Bedford, MA.) was used to measure
2 WBLM and LLLM. A Certified Clinical Densitometrist (CM) performed all DXA data
3 collection and analysis procedures. Prior to each measurement session an automatic
4 calibration procedure was performed to assess and maintain the measurement precision and
5 accuracy of the DXA. During the procedure, subjects lay motionless in a supine position on a
6 table for eight minutes, while an X-ray fan array passed above the table. WBLM and LLLM
7 were determined using manufacturer-supplied software (APEX version 4.0, Hologic
8 Discovery).

9 **2. Warm up**

10
11 Following the DXA scan, and prior to all performance measures, a 15-minute warm up
12 consisting of 5 minutes of cycling at 50 watts on a cycle ergometer (Velotron Dynafit Pro,
13 RaceMate, Seattle, WA, USA). Followed by 10 body weight squats, 10 heel raises, 10
14 countermovement jumps (CMJ). All were undertaken at moderate intensity. Participants then
15 completed each of the following performance measures.

16 **3. Muscular power**

17 Muscular power was assessed using a CMJ test. CMJ trials were performed three times on an
18 AMTI force plate (Advanced Medical Technology Inc., Watertown, USA). The analogue
19 signal sampled at 1000Hz was converted to a digital signal using a Powerlab 30 series data
20 acquisition system (AD Instruments, Sydney, Australia), and data were collected using
21 custom-written LabView software Version 2011 (National Instruments, Texas, USA). The
22 vertical force-time data were filtered using a fourth-order Butterworth low-pass filter with a
23 cut-off frequency of 17 Hz. Participants were instructed to perform a fast downward
24 movement (to 90° knee flexion) immediately followed by a fast upward movement, and to
25 jump as high as possible. Hands were kept on the hips to minimize any influence of the arm

1 swing. Each trial was followed by 2 minutes of passive rest, and the mean of three jumps
2 (cm) was used for further analysis.

3 **4. Muscular strength**

4 Quadriceps and hamstring peak isometric torque (QPT and HPT) of the dominant leg was
5 measured using a Biodex System 3 isokinetic dynamometer (Biodex Medical Systems,
6 Shirley, NY, USA). Subjects performed three x 10-second maximal isometric knee
7 extensions (QPT) and three 10-second maximal isometric knee flexions with strong verbal
8 encouragement. The effort with the highest peak torque ($\text{Nm}\cdot\text{kg}^{-1}$) was used for subsequent
9 data analysis (24).

10 **5. Anaerobic performance**

11 Sprint cycling performance was measured using 10 and 30 second sprint tests on a Velotron
12 ergocycle (Racermate, Seattle, USA) with a 5-minute passive rest period between tests.
13 Following familiarisation of the protocol and a warm-up consisting of pedalling at a self-
14 selected cadence at a set resistance of 50 W for five minutes interspersed with three practice
15 maximal accelerations over 2-3 seconds, the resistance of the ergocycle was adjusted at 75
16 $\text{g}\cdot\text{kg}^{-1}$ of body mass (39). Peak power ($\text{W}\cdot\text{kg}^{-1}$) in the 10-second test and total 30 second work
17 ($\text{kJ}\cdot\text{kg}^{-1}$) was used for subsequent data analysis.

18 **7. Peak power output**

19 A graded maximal exercise test to measure peak power output (PPO) was completed on an
20 electrically-braked, computer controlled cycle ergometer (Velotron Dynafit Pro, RaceMate,
21 Seattle, USA). Gas analysis was undertaken using a Fitmate Pro (Cosmed, Rome, Italy)
22 following a 5-minute warm-up at 30 W cycling and a pedalling cadence of 90 rpm throughout
23 the test. The work increments for each 1-minute stage were 15 W. The test ceased when two
24 or more criteria for attainment of $\text{VO}_{2\text{peak}}$ were achieved. These criteria included no
25 significant increase in O_2 uptake with an increase in work rate, attainment of the age-

1 predicted maximum heart rate, and/or volitional exhaustion (36). Peak power output (PPO)
2 was calculated from the last completed work rate, plus the fraction of time spent in the final
3 non-completed work rate multiplied by 25 watts (16).

4 **8. Flying 200 meter time sprint time**

5 Forty-eight hours after the laboratory tests, flying 200 meter sprint time was assessed at a
6 local concrete and banked (31 degrees), 333 meter cycling velodrome with participants using
7 their own road bikes to perform a total of three flying 200 meter attempts. Following a ten lap
8 warm up, participants then performed two familiarisation attempts of the flying 200 meter
9 time trial before ten minutes of passive seated rest. The flying 200 meter time trial
10 commenced by each participant cycling around the velodrome two times in attempt to build
11 up speed, and on the third lap, participants were instructed to come down the bank of the
12 velodrome at maximal speed when crossing the starting line. Flying 200 meter sprint time
13 was recorded by three, experienced observers using hand-held stopwatches (Hart sports timer
14 898, Hart Sport, Aspley, Australia). Observers were instructed to start the stopwatches when
15 the participant crossed the start line with the front end of the front wheel and stop the
16 stopwatches when the participant crossed the finish line with the front wheel. The mean of
17 three trials was recorded for subsequent analysis.

18 **Sprint cycling training Program**

19 The sprint cycling training program was designed in consultation with an accredited track
20 cycling coach and supervised by the same coach for each of the twice weekly sessions. Both
21 CT and ST groups performed two 60-90 minute sprint cycling training sessions per week,
22 separated by 48 hours. Sprint cycling sessions consisted of a five to ten minute warm-up (10-
23 15 x 333 meter laps at a self-selected pace) after which subjects performed 1-3 sets x 1-3
24 repetitions of maximal effort sprints ranging in distance from 65 meters to 333 meters with 2-
25 3 mins of active then passive recovery between repetitions and 10 minutes passive rest

1 between sets. At the completion of the track training session, subjects performed a five to ten
2 minute cool down (10-15 laps of the velodrome at a self-selected pace). Using an undulating
3 periodization program; participants commenced the track program using a 92 inch gear and
4 throughout the 12-week period, progressed to a 104 inch gear (table 2). As a result, the ST
5 group reduced their usual weekly endurance cycling training by three hours per week. The
6 overall training adherence rate calculated as a percentage of the total sprint cycling training
7 sessions successfully completed was $82 \pm 5.1\%$ for ST group across the 12-week study period.

8 Table 2 about here

9 Table 3 about here

10

11 **Strength training program**

12 The CT group replaced four of their usual weekly endurance cycling training sessions with
13 two evening group track sprint-cycling training sessions as described above, and two morning
14 group gym-based strength training sessions per week. As a result the CT group reduced their
15 usual weekly endurance cycling training by six hours per week. Participants were advised to
16 perform two 60 minute recovery rides (50-70% MHR, 90-110rpm) and not undertake other
17 cycling training sessions throughout the training week to avoid overtraining and excessive
18 fatigue. All four training sessions were supervised by an accredited strength and conditioning
19 coach. Strength training sessions were conducted on alternate days to the track sprint training
20 days. The strength training program and relative volumes of the different modes of strength
21 during the course of the study are summarized in Table 4. During each training session,
22 subjects performed the following exercises in order [1] Plyometric and explosive strength
23 exercises: double leg vertical and horizontal hops or jumps, single leg alternating box jumps,
24 leg press throws. [2] Strength training exercises: single-leg leg presses, seated hip flexions.

1 [3] Hypertrophy exercises: leg curls, leg extensions, seated calf-raises, supine hip extensions,
2 chest press, bench rows, abdominal curl ups and lower back extensions. Recovery time of
3 two minutes between sets and exercises was strictly controlled, and each strength training
4 session lasted approximately 90 minutes. The strength training program incorporated an
5 undulating periodization approach, to reduce the potential for overtraining and to optimise
6 adaptation. Subjects completed electronic training logs (Accelaware, Sports Performance
7 Systems, Brisbane, Australia) describing all their training parameters (number of repetitions,
8 sets, loads, distances, track sprint cycling times) to monitor progress and to provide
9 motivation for maximal effort during the training program. The overall strength training
10 adherence rate, calculated as a percentage of training sessions successfully completed, was 85
11 \pm 3.8% for CT group across the 12-week study period.

12 Table 4 about here

13 **Control group**

14 The CON group were asked to maintain eight-hours per week, of their current endurance
15 cycling training program (table 3). In comparison to the CON group, the CT undertook two
16 hours per week of endurance training for 12-weeks, whilst the ST group undertook five hours
17 per week of endurance training for 12-weeks (table 5).

18 Table 5 about here

20 **Data analysis**

21 The training related effects were measured using a three (group) x two (time) repeated
22 measures analysis of variance (ANOVA). If a main effect was observed, a Tukey post-hoc
23 test was undertaken to identify the source of the differences. A p value of <0.05 was
24 considered statistically significant. Twenty-three of the twenty-four dependant variables were

1 normally distributed, as assessed by Shapiro-Wilk's test ($p > .05$), although one variable did
2 not meet the assumption of normality (Post_PP10; $p = 0.033$). For this variable, data were log
3 transformed and the equivalent non-parametric statistic used. This did not change the
4 outcome for this variable, and thus for ease of interpretation, we report findings from
5 parametric statistics only. Cohen's conventions for effect size (ES) were used for
6 interpretation for no effect ($ES < 0.2$), small effect ($0.2-0.49$), moderate effect ($0.5-0.79$), and
7 large effect (> 0.8) (5). SPSS Version 20 (IBM, Corp, New York) software was used for all
8 statistical analyses.

9 **Results**

10 Pre and post-test values for each dependant variable for each of the intervention groups are
11 shown in Table 6. No pre-training differences were observed between CT group, ST group
12 and the CG group for any of the dependant variables.

13 **Lean mass**

14 No changes in WBLM occurred during the intervention in all groups ($F(2, 22) = 2.4$, $p =$
15 0.11) (Table 6). There were no significant between group effects for LLLM ($F(2,22) = 2.7$, p
16 $= 0.89$). However there was a significant effect of time ($F(1, 22) = 10.61$, $p = 0.04$). LLLM
17 increased in CT group ($p = 0.01$, 4.5%, $ES = 0.35$), and in the ST group ($p = 0.03$, 3.5%, ES
18 $= 0.45$).

20 **Muscular power**

21 No changes in CMJ occurred during the intervention in all groups ($F(1, 24) = 0.48$, $p = 0.69$).
22 (table 6).

23 **Muscular strength**

24 No changes in either QPT or HPT occurred during the intervention in all groups. ($F(2, 22) =$
25 2.61 , $p = 0.96$); ($F(2, 22) = 2.32$, $p = 0.14$) (table 6).

1 **Sprint cycling performance**

2 A significant group x time interaction was observed for PP10 ($F(2, 22) = 3.50, p = 0.48$),
3 however subsequent post hoc analysis revealed no differences between groups (table 6). A
4 significant group x time interaction was also observed for TW ($F(2, 22) = 5.59, p = 0.01$,
5 6.9%, $ES = -0.59$), subsequent a Tukey post hoc analysis revealed a difference in TW
6 between ST and CG groups ($p = 0.02$).

7 **Peak power output**

8 No changes in PPO occurred during the intervention in all groups ($F(2, 22) = 1.61, p = 0.22$)
9 (table 6).

10 **Flying 200 meter sprint time trial**

11 A significant group x time interaction was observed for TT ($F, (2, 22) = 11.70, p = 0.00$)
12 however subsequent post hoc analysis revealed no differences between groups. There was
13 also a significant effect of time ($F(1, 22) = 7.21, p = 0.01$). TT decreased in the CT group (p
14 $< 0.01, -7.7\%$, $ES = 0.85$). In the CON group, TT increased ($p = 0.07, -8.8\%$, $ES = 0.85$).

15
16 Table 6 about here

18 **Discussion**

19 The success in many endurance events such as road cycling and running could be dependent
20 not only good aerobic capabilities but also muscle characteristics and related sprint
21 performance. The purpose of this study was to examine whether lean mass, strength, power
22 and sprint performance could be improved by short term concurrent training in a group of
23 masters road cyclists who had no previous experience in strength and sprint training. The
24 major finding was that 12 weeks of concurrent strength and sprint training increased LLLM
25 and improved TT performance in masters road cyclists.

1 There are very few training studies on aging athletes and we are not aware of any previous
2 interventions on road cyclists. Our findings of training induced change in LLLM are in
3 agreement with previous research, which has reported increases in muscle mass or fibre area
4 in response to concurrent strength training in masters sprint and endurance runners (10,31,33).
5 For example, Piacentini et al. (31) reported a non-significant 2% increase in lean mass in a
6 group (n=6, 44.2 ± 3.9 years) of male and female masters endurance runners following six
7 weeks of concurrent endurance running and strength training. However the duration of the
8 latter study (six weeks), may not have been long enough to observe significant changes in
9 lean mass, as it is generally understood that muscle hypertrophy requires greater than eight
10 weeks of strength training (37). The ST group in the current study demonstrated a 3.5%
11 increase in LLLM which is surprisingly higher than the increases in lean mass reported in
12 younger cohorts who have undergone sprint interval training programs lasting between eight
13 weeks to eight months (18,27). These differences may be explained by the use of heavy
14 gearing in the present study with the ST gearing progressively increased over the 12 week
15 training program, thus providing a form of progressive overload that may have stimulated an
16 increase LLLM. Taken together, the results of the current study suggest ST positively affects
17 lean mass in masters cyclists. These findings support the use of ST as an alternative exercise
18 intervention to increase lower limb lean mass in masters road cyclists.

19 In the present study, CMJ did not significantly increase following 12 weeks of CT. These
20 results are in contrast to the findings of Cristea et al. (10) who reported a significant
21 improvement in squat jump height in a group of male masters sprint runners (n=7, 71.0 ± 5.0
22 years) who completed a 20 week progressive strength training program. However, the
23 previous researchers used a squat jump test which does not utilize the stretch-shortening
24 cycle, making a true comparison of the present results difficult. In contrast, the lack of a
25 significant increase in CMJ following 12 weeks of concurrent resistance and sprint training

1 observed in the current study are in agreement with the findings of Piacentini et al. (29) who
2 reported six weeks of concurrent endurance running and strength training did not
3 significantly improve CMJ in a group ($n=6$, 44.2 ± 3.9 years) of male and female masters
4 endurance runners. Despite not reaching significance, the participants in the Piacentini et al.
5 (29) study improved countermovement jump height by 3.2% which is similar to the 2.7%
6 increase in CMJ observed in the CT group. A lack of a significant improvement in CMJ in
7 the current study, may also be attributed to a possible interference effect known to affect
8 explosive strength when strength training is combined with endurance training (14). Despite
9 reducing their endurance training volume, the CT group still performed more than two
10 scheduled endurance sessions a week throughout the whole study period. Taken together,
11 these results suggest 12 weeks of CT or ST may not significantly improve muscular power in
12 masters road cyclists.

13 In the present study 12 weeks of CT did not significantly improve QPT or HPT in the CT
14 group. Age-related declines in muscular strength is commonly associated with the age-related
15 loss of lean mass observed in masters runners, swimmers and cyclists (1). These age-related
16 declines in muscular strength and muscle mass may contribute to the observed reduction in
17 cycling performance with age. It has been shown that strength improvements are lower when
18 endurance training is combined with a strength training program (17) as a result of conflicting
19 cellular stimuli (26). In the current study, participants in the CT group performed more than
20 the prescribed limit of endurance cycling training sessions throughout the 12 week CT
21 program, which could explain why no changes were observed in QPT and HPT observed in
22 the CT group. Similarly, 12 weeks of ST did not significantly improve QPT or HPT. To the
23 best of our knowledge, no studies to date, have investigated the effects of ST on muscle
24 strength in masters cyclists. However, in younger cohorts, repeat sprint training has been
25 shown to increase lower limb strength (8, 15). For example, Harridge et al. (15) reported a

1 significant increase in maximal isometric knee extensor torque (7%) following six weeks of
2 sprint cycling training, performed four times per week, in a group of recreationally active,
3 younger males ($n=7$, 22 ± 2 years). Taken as a whole, the results of the present study showed
4 12 weeks of CT or ST does not significantly increase knee flexion or knee extension strength
5 in masters road cyclists.

6 The ability to generate brief, high powered outputs is an important component of competitive
7 cycling performance (2). In the present study 12 weeks of CT did not significantly increase
8 PP10 or TW in the CT group. No research to date, has investigated the effects of CT on PP10
9 or TW in healthy older adults or masters cyclists. However, in a cross sectional analysis of
10 highly trained masters cyclists ($n= 173$, 35-64 years). Gent and Norton (13) reported PP10
11 and TW declined by 8.1% and 8.0% per decade. In contrast, 12 weeks of ST did not
12 significantly improve PP10 or TW in the ST group. These results are in contrast to similar
13 studies in younger cohorts (9), which have reported significant improvements in TW. For
14 example Creer et al. (9) reported 4 weeks of sprint cycling training, performed two times per
15 week, significantly increased total 30 second work (6.0%) as measured by cycle ergometry,
16 in a group of younger, trained cyclists ($n=10$, 25.1 ± 2.3 years). The lack of improvement in
17 PP10 & TW in the CT group may be a consequence of insufficient recovery between exercise
18 training and testing. In particular, subjects in all groups continued their endurance training at
19 the completion of the 12 week program up until the date of testing. Future research is
20 warranted to better understand the effect of CT and ST on anaerobic performance in masters
21 road cyclists.

22 In the current study, PPO was unaffected by 12 weeks of either CT or ST. To date, the effects
23 of CT or ST on PPO in masters cyclists is unknown. However, the use of strength training to
24 improve endurance cycling performance in healthy, younger and older adults is well
25 supported (6, 19, 25, 34, 42). For example Loveless et al. (25) reported 8 weeks of maximal

1 leg-strength training significantly improved cycling peak aerobic power, in a group of healthy,
2 younger males ($n=7$, 25.0 ± 2.0 years). Additionally, Izquierdo et al. (19) reported 16 weeks
3 of progressive strength-training significantly increased cycling peak aerobic power in a group
4 of healthy, older males ($n=11$, 64-74 years). In the present study the ST group did not
5 significantly improve PPO. These results are in contrast to the findings from studies in
6 younger cyclists, which have reported significant increases in PPO following sprint cycling
7 training (21, 39). For example, Laursen et al. (21) reported a significant improvement in peak
8 aerobic power following two weeks of sprint cycling training in a group of trained, younger
9 cyclists ($n=14$, 23.5 ± 3.5 years). Unsurprisingly, the current study observed no significant
10 change in PPO following the 12 week training period. These results suggest, reducing cycling
11 endurance training volume and replacing it with either CT or ST, does not negatively affect a
12 primary marker of endurance performance in masters road cyclists.

13 In the present study 12 weeks of CT significantly improved TT (8.1%) in the CT group.
14 Typical for road cycling competition is that a large group of riders are often together until the
15 end of the race and the ability to sprint to the finish line determines the place in the race. Thus,
16 sprinting speed is of particular importance to cycling performance. To date, no studies have
17 investigated the effects of CT on sprint cycling TT performance in masters cyclists. However,
18 studies investigating the effects of concurrent strength and sprint running training have
19 reported favourable effects on sprint running performance (10,33). For example, Cristea et al.
20 (10) reported a significant improvement in 60 meter sprint running time (2%) following 20
21 weeks of progressive strength training program performed 4 times per week in a group of
22 male masters sprint runners. In addition, Reaburn et al. (33) reported a significant
23 improvement in 100 meter (4%) and 300 meter (2%) sprint running time following eight
24 weeks of concurrent strength and sprint running training performed four times per week.
25 Surprisingly, 12 weeks of ST did not significantly improve TT performance in the ST group.

1 The lack of improvement in TT performance in the ST group, could be attributed to a small
2 sample size or inadequate recovery. Training logs show, several participants in the ST group
3 did not reduce their endurance training volume, on the days leading into the final TT. Finally,
4 in the present study there was no between group differences in TT performance amid the CT
5 group and ST groups, suggesting that the addition of strength training to a ST program may
6 not provide additional benefits to sprint cycling performance. Taken together, these results
7 suggest 12 weeks of CT significantly improves TT performance, which can benefit the
8 masters road cyclists by improving sprint speed to the finish line.

9 We acknowledge several limitations to the current study. Firstly, improvements observed in
10 sprint cycling performance in the CT may have resulted from a placebo effect. For example,
11 the ST group undertook two modified ST sessions per week, in comparison, the CT
12 undertook four modified CT sessions per week, which may have doubled the placebo effect.
13 Secondly, the CT had greater adherence to the sprint training sessions when compared to the
14 ST group, which may further explain the larger improvements in sprint performance observed
15 in the CT group and ST groups. Future studies should match total sprint and strength training
16 volumes. Thirdly, the specialised population of this group, limited the statistical power of this
17 study. Finally, it should also be acknowledged that sprints performed during a competitive
18 road-cycling event often occur in a fatigued state, whereas in the present study, sprinting time
19 trials were performed in non-fatigued state, further limiting the applications of these findings.

21 **Practical Applications**

22 Previous research suggests masters cyclists face an age-related decline in lean mass, muscular
23 strength and power, and sprinting performance. These declines may contribute to the age-
24 related decline in competitive cycling performance, particularly the ability to accelerate
25 rapidly or sprint to the finish line during a race. The results of the present study suggest that

1 12 weeks of CT significantly improves lower body lean mass and sprint cycling time trial
2 performance. In the ST group, 12 weeks of ST significantly improved lower body lean mass
3 only. Based on these findings, improvements in sprint-cycling performance in masters
4 endurance cyclists can be made by undertaking 12-weeks of CT during the general
5 preparation phase of training. Thereafter, the effects of CT could be maintained by
6 performing one strength session and one sprint training session per week throughout the late
7 preparation and competitive periods. Moreover, performing sprint training at a cycling
8 velodrome, including the use of banking, can be used to develop speed, acceleration and
9 maximum velocity. Finally, the use of progressively heavier gearing ratios can enhance
10 cycling specific strength development. However, a more definitive study over several months
11 should be undertaken to clarify the optimal trimming and amount of CT, particularly how the
12 replacement of a portion of endurance training impacts road cycling performance. Finally, it
13 should also be emphasized that this study provides only initial findings about the good
14 adaptive capacity and training specificity in masters cyclists. In future it is essential to obtain
15 knowledge of potential negative effects of combined training with decreased aerobic training
16 on overall competitive cycling performance as a base for planning optimal training for
17 masters cyclists.

20 REFERENCES

- 21 1. Alway SE, Coggan AR, Sproul MS, Abduljalil AM, Robitaille PM. Muscle torque in
22 young and older untrained and endurance-trained men. *J Gerontol A Biol Sci Med Sci*
23 51: B195-B201, 1996.
- 24 2. Atkinson G, Davison R, Jeukendrup A, Passfield L. Science and cycling: current
25 knowledge and future directions for research. *J Sports Sci* 21: 767-787, 2003.

- 1 3. Bell KE, Séguin C, Parise G, Baker SK., & Phillips, S. M. Day-to-day changes in
2 muscle protein synthesis in recovery from resistance, aerobic, and high-intensity
3 interval exercise in older men. *J Gerontol A Biol Sci Med Sci* 70: 1024-1029, 2015.
- 4 4. Cycling Australia 2015 Annual Report. (2015, September/October). Retrieved
5 February 28, 2016 from
6 http://www.cycling.org.au/Portals/10/CA_2015_AnnualReport_FINALemail.pdf
- 7 5. Cohen J. *Statistical power analysis for the behavioral sciences*. Hillsdale, N.J:
8 Earlbaum, 1988.
- 9 6. Bastiaans J, Diemen Av, Veneberg T, Jeukendrup A. The effects of replacing a
10 portion of endurance training by explosive strength training on performance in trained
11 cyclists. *Eur J Appl Physiol* 86: 79-84, 2001.
- 12 7. Brisswalter J & Nosaka K. (2013). Neuromuscular factors associated with decline in
13 long-distance running performance in master athletes. *Sports Med* 43: 51-63, 2013.
- 14 8. Cantrell GS, Schilling BK, Paquette MR, Murlasits Z. Maximal strength, power, and
15 aerobic endurance adaptations to concurrent strength and sprint interval training. *Eur*
16 *J Appl Physiol* 114: 763-771, 2014.
- 17 9. Creer A, Ricard M, Conlee R, Hoyt G, Parcell A. Neural, metabolic, and performance
18 adaptations to four weeks of high intensity sprint-interval training in trained cyclists.
19 *Int J Sports Med* 25: 92-98, 2004.
- 20 10. Cristea A, Korhonen M, Häkkinen K, Mero A, Alén M, Sipilä S, Viitasalo J,
21 Koljonen M, Suominen H, Larsson L. Effects of combined strength and sprint training

- 1 on regulation of muscle contraction at the whole-muscle and single-fibre levels in
- 2 elite master sprinters. *Acta Physiol* 193: 275-289, 2008.
- 3 11. Fatouros I, Kambas A, Katrabasas I, Nikolaidis K, Chatzinikolaou A, Leontsini D,
4 Taxildaris K. Strength training and detraining effects on muscular strength, anaerobic
5 power, and mobility of inactive older men are intensity dependent. *Br J Sports Med*
6 39: 776-780, 2005.
- 7 12. Frontera WR, Meredith CN, O'Reilly KP, Evans WJ. Strength training and
8 determinants of VO₂max in older men. *J Appl Physiol* 68: 329-333, 1990.
- 9 13. Gent DN and Norton K. Aging has greater impact on anaerobic versus aerobic power
10 in trained masters athletes. *J Sports Sci* 31: 97-103, 2013.
- 11 14. Häkkinen K, Alen M, Kraemer W, Gorostiaga E, Izquierdo M, Rusko H, Mikkola J,
12 Häkkinen A, Valkeinen H, Kaarakainen E. Neuromuscular adaptations during
13 concurrent strength and endurance training versus strength training. *Eur J Appl*
14 *Physiol* 89: 42-52, 2003.
- 15 15. Harridge S, Bottinelli R, Canepari M, Pellegrino M, Reggiani C, Esbjörnsson M,
16 Balsom P, Saltin B. Sprint training, in vitro and in vivo muscle function, and myosin
17 heavy chain expression. *J App Phys* 84: 442-449, 1998.
- 18 16. Hawley JA, Noakes TD. Peak power output predicts maximal oxygen uptake and
19 performance time in trained cyclists. *Eur J Appl Physiol* 65: 79-83, 1992.
- 20 17. Hennessy LC, Watson AWS. The interference effects of training for strength and
21 endurance simultaneously. *J Strength Cond Res* 8:12-19, 1992.

- 1 18. Heydari M, Freund J, and Boutcher SH. The effect of high-intensity intermittent
2 exercise on body composition of overweight young males. *Journal of obesity* 2012,
3 2012.
- 4 19. Izquierdo M, Häkkinen K, Ibañez J, Antón A, Garrués M, Ruesta M, Gorostiaga EM.
5 Effects of strength training on submaximal and maximal endurance performance
6 capacity in middle-aged and older men. *J Strength Cond Res* 17: 129-139, 2003.
- 7 20. Klitgaard H, Manton M, Schiaffino S, Ausoni S, Gorza L, Laurent-Winter, C, &
8 Saltin B. Function, morphology and protein expression of ageing skeletal muscle: a
9 cross-sectional study of elderly men with different training backgrounds. *Acta Physiol*
10 *Scand*, 140:41-54, 1990.
- 11 21. Laursen PB, Blanchard MA, Jenkins DG. Acute high-intensity interval training
12 improves Tvent and peak power output in highly trained males. *Can J Appl Physiol*
13 27: 336-348, 2002.
- 14 22. Lepers, R., Rüst, C. A., Stapley, P. J., & Knechtle, B. Relative improvements in
15 endurance performance with age: evidence from 25 years of Hawaii Ironman racing.
16 *Age*, 35: 953-962, 2013.
- 17 23. Louis J, Hausswirth C, Easthope C, Brisswalter J. Strength training improves cycling
18 efficiency in master endurance athletes. *Eur J Appl Physiol* 112: 631-640, 2012.

- 1 24. Louis J, Hausswirth C, Bieuzen F, & Brisswalter, J. (2009). Muscle strength and
2 metabolism in master athletes. *Int J Sports Med* 30: 754-759, 2009.
- 3 25. Loveless DJ, Weber CL, Haseler LJ, and Schneider DA. Maximal leg-strength
4 training improves cycling economy in previously untrained men. *Med Sci Sports*
5 *Exerc* 37: 1231, 2005.
- 6 26. Lundberg TR, Fernandez-Gonzalo R, Tesch PA. Exercise-induced AMPK activation
7 does not interfere with muscle hypertrophy in response to resistance training in men.
8 *J Appl Physiol* 116: 611-620, 2014.
- 9 27. Macpherson R, Hazell TJ, Olver TD, Paterson DH, and Lemon P. Run sprint interval
10 training improves aerobic performance but not maximal cardiac output. *Med Sci*
11 *Sports Exerc* 43: 115-122, 2011.
- 12 28. Nederveen J, Joanisse S, Séguin C, Bell K, Baker S, Phillips S, Parise G. The effect of
13 exercise mode on the acute response of satellite cells in old men. *Acta Physiol* 215:
14 177-190, 2015.
- 15 29. Norton K. Sports Medicine Australia pre-exercise screening system. *Sports Medicine*
16 *Australia*, 2005.
- 17 30. Paton CD & Hopkins WG. (2005). Combining explosive and high-resistance training
18 improves performance in competitive cyclists. *J Strength Cond Res* 19:826-830, 2005.
- 19 31. Piacentini MF, De Ioannon G, Comotto S, Spedicato A, Vernillo G, and La Torre A.
20 Concurrent strength and endurance training effects on running economy in master
21 endurance runners. *J Strength Cond Res* 27: 2295-2303, 2013.
- 22 32. Reaburn P, Dascombe B. Endurance performance in masters athletes. *Eur Rev Aging*
23 *Phys Act* 5: 31-42, 2008.

- 1 33. Reaburn P, Logan P, Mackinnon L. The Effect of Hypertrophy Resistance Training
2 on Anaerobic Work Capacity in Veteran Sprint Runners. Canberra, Australia:
3 Australian Sports Commission, 1994.
- 4 34. Rønnestad BR, Mujika I. Optimizing strength training for running and cycling
5 endurance performance: A review. *Scand J Med Sci Sports* 24: 603-612, 2014.
- 6 35. Ryngøy A.F. The effect of two different types of resistance training on the cycling
7 performance of highly trained veteran cyclists. (unpublished masters thesis).
8 University of Norland, Bodø, Norway, 2012.
- 9 36. Schell J, Leelarthapin B. *Physical fitness assessment in exercise and sport science*.
10 Leelar Biomediscience Services, 1990.
- 11 37. Staron R, Karapondo D, Kraemer W, Fry A, Gordon S, Falkel J, Hagerman F, Hikida
12 R. Skeletal muscle adaptations during early phase of heavy-resistance training in men
13 and women. *J App Phys* 76: 1247-1255, 1994.
- 14 38. Tanaka H and Seals D. Endurance exercise performance in Masters athletes:
15 age-associated changes and underlying physiological mechanisms. *J Physiol* 586: 55-
16 63, 2008.
- 17 39. Westgarth-Taylor C, Hawley JA, Rickard S, Myburgh KH, Noakes TD, Dennis SC.
18 Metabolic and performance adaptations to interval training in endurance-trained
19 cyclists. *Eur J Appl Physiol Occup Physiol* 75: 298-304, 1997.

- 1 40. Widrick Trappe SW, Costill DL, & Fitts RH. Force-velocity and force-power
2 properties of single muscle fibers from elite master runners and sedentary men. *Am J*
3 *Physiol Cell Physiol* 271: C676-C683, 1996.
- 4 41. Yamamoto LM, Klau JF, Casa DJ, Kraemer WJ, Armstrong LE, and Maresh CM. The
5 effects of resistance training on road cycling performance among highly trained
6 cyclists: a systematic review. *J Strength Cond Res* 24: 560-566, 2010.
- 7 42. Zajac A, Jarzabek R, and Waskiewicz Z. The Diagnostic Value of the 10-and 30-
8 Second Wingate Test for Competitive Athletes. *J Strength Cond Res* 13: 16-19, 1999.

ACCEPTED

Table 1: Physical and training characteristics of participants.

	CT	ST	CG
	(n=9)	(n=7)	(n=9)
Age (years)	53.5 ± 9.3	49.4 ± 4.8	56.9 ± 8.6
Stature (m)	1.80 ± 0.08	1.80 ± 0.10	1.75 ± 0.10
Body mass (kg)	81.9 ± 6.1	78.5 ± 6.1	83.5 ± 10.0
Training hours (hr/week)	8.2 ± 1.0	8.1 ± 1.3	8.0 ± 1.2

CT = combined strength and sprint group; ST = sprint training group; CG = control group; data are Mean ± SD.

Table 2 Flying 200-m Track Cycling Program

	Session 1	Session 2
Weeks 1-4	<p>Warm up:</p> <ul style="list-style-type: none"> • 15 minute easy rolling laps (low gear) with gradual windup from 30km/h up to 40km/h <p>Conditioning phase (@80% of max speed):</p> <ul style="list-style-type: none"> • 3-5 minutes active recovery between reps • 15 minutes passive recovery between sets <p>Set 1:</p> <ul style="list-style-type: none"> • 3 x 65m @ G92 standing start. 1 x 100m seated from 20kph <p>Set 2:</p> <ul style="list-style-type: none"> • 3 x 65m @ G94 standing start. 3 minutes active recovery between repetitions 	<p>Warm up:</p> <ul style="list-style-type: none"> • 15 minute easy rolling laps (low gear) with gradual windup from 30km/h up to 40km/h <p>Conditioning phase (@80% of max speed):</p> <ul style="list-style-type: none"> • 3-5 minutes active recovery between reps • 15 minutes passive recovery between sets <p>Set 1:</p> <ul style="list-style-type: none"> • 1 x Flying 100m @ G94 • 1 x Flying 100m @ G96 • 1 x Flying 100m @ G 98 <p>Set 2:</p>

- 1 x 200m seated from 20kph

Set 3: 3 x 65m @ G96 standing start.

- 1 x 333m seated from 30kph

Cool-down:

- 10-15 laps at a very low intensity or 5-10 minutes on rollers

Weeks 5-8

Warm up:

- 15 minute easy rolling laps (low gear) with gradual windup from 30km/h up to 40km/h

Conditioning phase (@ 90% max speed):

- 3-5 minutes active recovery between reps
- 15 minutes passive recovery between sets

Set 1:

- 3 x 65m @ G96 standing start.
- 1 x 100m seated from 20kph

Set 2:

- 3 x 65m @ G98 standing start.

- 1x flying 33m @G98

- 1x flying 33m @G100

Cool-down:

- 10-15 laps at a very low intensity or 5-10 minutes on rollers

Warm up:

- 15 minute easy rolling laps (low gear) with gradual windup from 30km/h up to 40km/h

Conditioning phase (@ 90% max speed):

- 3-5 minutes active recovery between reps
- 15 minutes passive recovery between sets

Set 1:

- 1 x Flying 100m @ G98
- 1 x Flying 100m @ G100
- 1 x Flying 100m @ G102

Set 2:

- 1 x 200m seated from 20kph

Set 3: 3 x 65m @ G100 standing start.

- 1 x 333m seated from 30kph

Cool-down:

- 10-15 laps at a very low intensity or 5-10 minutes on rollers

Weeks 9-12

Warm up:

- Rollers 10 minutes, including several short sprints

Conditioning phase (@90-98% max speed):

- 3-5 minutes active recovery between reps
- 15 minutes passive recovery between sets

Set 1:

- 3 x 65m @ G98 standing start.
- 1 x 100m seated from 20kph

Set 2:

- 3 x 65m @ G96 standing start.
- 1 x 200m seated from 20kph

Set 3: 3 x 65m @ G194 standing start.

- 1x flying 33m @ G102

- 1x flying 33m @ G104

Cool-down:

- 10-15 laps at a very low intensity or 5-10 minutes on rollers

Warm up:

- Rollers 10 minutes, including several short sprints

Conditioning phase (@90-98% max speed):

Set 1:

- 1 x Flying 100m @ G100
- 1 x Flying 100m @ G98
- 1 x Flying 100m @ G96

Cool-down:

- 10-15 laps at a very low intensity or 5-10 minutes on rollers

- 1 x 333m seated from 30kph

Cool-down:

- 10-15 laps at a very low intensity or 5-10 minutes on rollers

G= gear ratio; K1= started gate sprints; fly-session = all sprints completed from a flying-start

Table 3: Endurance Training Program

Day	Training Session Description
Monday	Recovery ride– 60 minutes @ 60% of HRmax
Tuesday	Endurance ride 90-180 minutes @ 60-70% of HRmax
Wednesday	Tempo ride – 60-90 minutes (3 x 15 minute efforts @ 85% of HRmax incorporated into the ride)
Thursday	Recovery ride 60 minutes @ 60% of HRmax
Friday	Cross training (swim, spin bike, cross-trainer) 45-60 minutes @ 60-70% of HRmax
Saturday	Group ride 60-90 minutes @ 75-85% HRmax; practise skills such as drafting
Sunday	Rest day

HRmax = age predicted maximal heart rate.

Table 4 strength training program

Day 1	Hypertrophy Phase				Day 2	Hypertrophy Phase			
Weeks	1	2	3	4	Weeks	1	2	3	4
Warm up	5 minutes stationary bike				Warm up	5 minutes stationary bike			
Plyometric Exercises					Plyometric Exercises				
ankle hops	2 x 8	2 x 10	2 x 12	2 x 10	ankle hops	2 x 8	2 x 10	2 x 12	2 x 10
side to side ankle hops	2 x 8	2 x 10	2 x 12	2 x 10	side to side ankle hops	2 x 8	2 x 10	2 x 12	2 x 10
standing jump & reach	2 x 8	2 x 10	2 x 12	2 x 10	standing jump & reach	2 x 8	2 x 10	2 x 12	2 x 10
Strength Exercises					Strength Exercises				
Leg Press	50% 2 x 12	60% 2 x 10	65% 2 x 10	60% 2 x 12	Leg Press	50% 2 x 12	60% 2 x 10	65% 2 x 10	60% 2 x 12
Seated Hip Flexion	50% 2 x 12	60% 2 x 10	65% 2 x 10	60% 2 x 12	Seated Hip Flexion	50% 2 x 12	60% 2 x 10	65% 2 x 10	60% 2 x 12
Hypertrophy Exercises					Hypertrophy Exercises				
Leg Curls	40% 12 x 4	50% 12 x 4	55% 12 x 4	45% 12 x 4	Leg Curls	40% 12 x 4	50% 12 x 4	55% 12 x 4	45% 12 x 4
Leg Ext	40% 12 x 4	50% 12 x 4	55% 12 x 4	45% 12 x 4	Leg Ext	40% 12 x 4	50% 12 x 4	55% 12 x 4	45% 12 x 4
seated Calve Raise	40% 12 x 4	50% 12 x 6	55% 12 x 4	45% 12 x 4	Calve Raise standing	40% 12 x 4	50% 12 x 4	55% 12 x 4	45% 12 x 4
Chest Press *	40% 12 x 4	50% 12 x 7	55% 12 x 4	45% 12 x 4	Shoulder Press	40% 12 x 4	50% 12 x 4	55% 12 x 4	45% 12 x 4

Trunk Stability					Trunk Stability				
plank	2 x 20secs	2 x3 0secs	1 x 60	3 x 30secs	plank	2 x 20secs	2 x3 0secs	1 x 60	3 x 30secs
prone back extensions	2 x 10	2 x 12	2 x 15	3 x 12	prone back extensions	2 x 10	2 x 12	2 x 15	3 x 12
Recovery and Cool Down					Recovery and Cool Down				
Static Stretching	2-3 x 30 second holds all muscle groups				Static Stretching	2-3 x 30 second holds all muscle groups			
Foam Roller	3 - 5 mins foam-rolling over all major muscle groups				Foam Roller	3 - 5 mins foam-rolling over all major muscle groups			

ACCEPTED

Day 1	Strength Phase				Day 2	Strength Phase			
Weeks	5	6	7	8	Weeks	5	6	7	8
Warm up	5 minutes stationary bike				Warm up	5 minutes stationary bike			
Plyometric Exercises					Plyometric Exercises				
Front Box Jump	2 x 10	2 x 12	2 x 15	2 x 12	Front Box Jump	2 x 10	2 x 12	2 x 15	2 x 12
jump from box	2 x 10	2 x 12	2 x 15	2 x 12	jump from box	2 x 10	2 x 12	2 x 15	2 x 12
lateral box jump	1 x 5	2 x 6	2 x 8	2 x 6	lateral box jump	1 x 5	2 x 6	2 x 8	2 x 6
Explosive Strength Exercises					Strength Exercises				
single leg press throw	20% 2x5	30% 2x5	40% 2x5	30% 3x5	single leg press throw	20% 2x5	30% 2x5	40% 2x5	30% 3x5
Strength Exercises					Strength Exercises				
Single Leg, Leg Press	70% 3 x 8	75% 3 x 8	80% 3 x 6	75% 3 x 8	Single Leg, Leg Press	70% 3 x 8	75% 3 x 8	80% 3x6	75% 3 x 8
Seated Hip Flexion	70% 3 x 8	75% 3 x 8	80% 3 x 6	75% 3 x 8	Seated Hip Flexion	70% 3 x 8	75% 3 x 8	80% 3x6	75% 3 x 8
Hypertrophy Exercises					Hypertrophy Exercises				
Leg Extensions	50% 12 x 3	60% 10 x 3	65% 10 x 3	55% 12 x 3	Leg Extension	50% 12 x 3	60% 10 x 3	60% 10 x 3	55% 12 x 3
Leg Ext	50% 12 x 3	60% 10 x 3	65% 10 x 3	55% 12 x 3	Leg Curls	50% 12 x 3	60% 10 x 3	60% 10 x 3	55% 12 x 3
Calve Raise	50% 12 x 3	60% 10 x 3	65% 10 x 3	55% 12 x 3	Calve Raise standing	50% 12 x 3	60% 10 x 3	60% 10 x 3	55% 12 x 3

Chest Press	50% 12 x 3	60% 10 x 3	65% 10 x 3	55% 12 x 3	Chest Press	50% 12 x 3	60% 10 x 3	60% 10 x 3	55% 12 x 3
Prone row	50% 12 x 3	60% 10 x 3	65% 10 x 3	55% 12 x 3	Prone Row	50% 12 x 3	60% 10 x 3	60% 10 x 3	55% 12 x 3
Trunk Stability					Trunk Stability				
Abdominal Curl up	2 x 10	2 x 12	2 x 15	3 x 12	Abdominal Curl up	2 x 10	2 x 12	2 x 15	3 x 12
bird dog	2 x 10	2 x 12	2 x 15	3 x 12	bird dog	2 x 10	2 x 12	2 x 15	3 x 12
Recovery and Cool Down					Recovery and Cool Down				
Static Stretching	2-3 x 30 second holds all muscle groups				Static Stretching	2-3 x 30 second holds all muscle groups			
Foam Roller	3 - 5 mins foam-rolling over all major muscle groups				Foam Roller	3 - 5 mins foam-rolling over all major muscle groups			

Day 1	Power Phase				Day 2	Power Phase			
Weeks	9	10	11	12	Weeks	9	10	11	12
Warm up	5 minutes stationary bike				Warm up	5 minutes stationary bike			
Plyometric Exercises					Plyometric Exercises				
alternating step push offs	1 x 10	1 x 15	1 x 20	2 x 15	alternating step push offs	1x10	1x15	1x20	2x15
single leg box push offs	1 x 10	1 x 15	1 x 20	2 x 15	single leg box push offs	1x10	1x15	1x20	2x15
squat depth jumps	2 x 10	2 x 15	2 x 20	2 x 15	squat depth jumps	2x10	2x15	2x20	2x15
Explosive Strength Exercises					Strength Exercises				
single leg press throw	45% 3x5	50% 3x5	60% 3x5	55% 3x5	single leg press throw	45% 3x5	50% 3x5	60% 3x5	55% 3x5
Strength Exercises					Strength Exercises				
SL Leg Press	85%, 3 x 5	90%, 3x3	95% 3x2	90% 4x3	SL Leg Press	85%, 3 x 5	90%, 3x3	95% 3x2	90% 4x3
Seated Hip Flexion	85%, 3 x 5	90%, 3x3	95% 3x2	90% 4x3	Seated Hip Flexion	85%, 3 x 5	90%, 3x3	95% 3x2	90% 4x3
Hypertrophy Exercises					Hypertrophy Exercises				
Leg Curls	60% 12 x 2	70% 10 x 2	75% 10 x 2	70% 12 x 2	Leg Curl	60% 12 x 2	70% 10 x 2	75% 10 x 2	70% 12 x 2
Leg Ext	60% 12 x 2	70% 10 x 2	75% 10 x 2	70% 12 x 2	Leg Ext	60% 12 x 2	70% 10 x 2	75% 10 x 2	70% 12 x 2
seated Calve Raise	60% 12 x 2	70% 10 x 2	75% 10 x 2	70% 12 x 2	Calve Raise standing	60% 12 x 2	70% 10 x 2	75% 10 x 2	70% 12 x 2
Chest Press	60% 12 x 2	70% 10 x 2	75% 10 x 2	70% 12 x 2	Chest Press	60% 12 x 2	70% 10 x 2	75% 10 x 2	70% 12 x 2

Prone row	60% 12 x 2	70% 10 x 2	75% 10 x 2	70% 12 x 2	Prone row	60% 12 x 2	70% 10 x 2	75% 10 x 2	70% 12 x 2
Trunk Stability					Trunk Stability				
Advanced Curl up	2 x 10	2 x 12	2 x 15	3 x 12	Advanced Curl up	2 x 10	2 x 12	2 x 15	3 x 12
back extension bench	2 x 10	2 x 12	2 x 15	3 x 12	back extension bench	2 x 10	2 x 12	2 x 15	3 x 12
Recovery and Cool Down					Recovery and Cool Down				
Static Stretching	2-3 x 30 second holds all muscle groups				Static Stretching	2-3 x 30 second holds all muscle groups			
Foam Roller	3 - 5 mins foam-rolling over all major muscle groups				Foam Roller	3 - 5 mins foam-rolling over all major muscle groups			

Table 5. Training volume over the 12 week intervention

Training Modality	CT Group	ST Group	CON Group
Sprint training (hrs/wk)	3.0 ± 0.5	3.0 ± 0.5	0.0
Strength training (hrs/wk)	3.0 ± 0.5	0.0	0.0
Endurance training (hrs/wk)	2.0 ± 0.3	5.0 ± 0.3	8.2 ± 1.0
Total weekly training(hrs/wk)	8.0 ± 1.3	8.0 ± 0.8	8.2 ± 1.0

ACCEPTED

Table 6: Changes in lean mass, laboratory measures and 200m sprint cycle performance following 12 weeks of CT or ST

	CT		Effect Size	ST		Effect Size	CG		Effect Size	P Values	
	Pre	Post		Pre	Post		Pre	Post		Between-Group	Time
WBLM (kg)	61.8 ± 5.2	63.1 ± 5.4	0.26 Small effect	61.4 ± 4.7	61.6 ± 5.1	0.16 No effect	61.5 ± 5.5	60.6 ± 6.2	-0.15 No effect	0.113	0.441
LLM (kg)	17.6 ± 1.9	18.4 ± 2.3 [†]	0.35 Small effect	17.0 ± 1.5	17.6 ± 1.4 [†]	0.45 Small effect	16.0 ± 2.0	16.0 ± 1.9	0.00 No effect	0.089	0.004
CMJ (cm)	24.4 ± 3.8	24.9 ± 4.4	0.12 No effect	25.1 ± 12.0	22.7 ± 24.3	-0.12 No effect	23.9 ± 7.0	21.7 ± 6.3	-0.33 Small effect	0.698	0.495

QPT (Nm·kg ⁻¹)	2.8 ± 0.5	3.0 ± 0.4	0.47 Small effect	3.1 ± 0.4	2.9 ± 0.4	-0.50 Small effect	2.6 ± 0.6	2.4 ± 0.8	-0.28 Small effect	0.096	0.813
HPT (Nm·kg ⁻¹)	1.0 ± 0.2	1.1 ± 0.2	0.32 Small effect	1.0 ± 0.2	1.1 ± 0.1	0.53 Moderate effect	1.0 ± 0.1	0.9 ± 0.3	-0.44 Small effect	0.149	0.122
PP10 (W·kg)	11.3 ± 1.8	11.5 ± 1.9	0.10 No effect	11.6 ± 1.2	12.0 ± 1.1	0.38 Small effect	10.5 ± 1.2	9.8 ± 1.9	-0.44 Small effect	0.048	0.780
TW (J·kg)	247.4 ± 35.0	255.1 ± 35.8	0.22 Small effect	256.0 ± 28.5	262.4 ± 19.2*	0.26 Small effect	227.5 ± 20.8	211.8 ± 30.9†	-0.59 Moderate effect	0.011	0.896
PPO (watts)	341.6 ± 62.6	338.8 ± 60.0	0.04 No effect	362.5 ± 37.7	378.1 ± 48.9	0.35 Small effect	316.6 ± 54.4	308.3 ± 59.9	-0.14 No effect	0.222	0.881
TT (sec)	16.0 ± 1.9	14.7 ± 1.3†	0.85 Large effect	14.7 ± 1.1	14.2 ± 0.6	0.61 Moderate effect	15.4 ± 1.0	15.9 ± 1.2	-0.45 Small effect	0.000	0.014

CT = combined strength and sprint group; ST = sprint training group; CG = control group; effect size = between group effect size. * = Between-group difference estimated by ANOVA: Tukey post-hoc test ($P < 0.05$); † = Significant effect of time ($p < 0.05$); WBLM = whole body Lean mass; LLLM = total lower limb lean mass; CMJ = counter movement jump height; QPT = quadriceps peak isometric torque; HPT = hamstring peak isometric torque; PP10 = ten second sprint peak power; TW = total 30 second work; TT = flying 200 meter sprint time; PPO = peak power output in incremental cycle ergometer test.