

Bond University
Research Repository



Determining the isometric hip abduction and adduction profile in a competitive surfing cohort

Furness, James; Pope, Rodney R

Licence:
CC BY-NC-ND

[Link to output in Bond University research repository.](#)

Recommended citation(APA):
Furness, J., & Pope, R. R. (2019). *Determining the isometric hip abduction and adduction profile in a competitive surfing cohort*. PO-E-21-SUN1. Poster session presented at World Confederation for Physical Therapy Congress 2019, Geneva, Switzerland.
<https://www.abstractstosubmit.com/wcpt2019/archive/#/viewer/abstract/2117>

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

For more information, or if you believe that this document breaches copyright, please contact the Bond University research repository coordinator.

Background

- Previous injury epidemiology research has shown the knee to be an injury prone region within competitive surfers



Frequency of acute knee injury ranges from 19-35%^{1,2,3}

- Research supports the influence that hip strength has on knee alignment and subsequent injury⁴
- It is theorized that the long hip abductors and adductors may provide mechanical support to the medial and lateral knee



- Minimal scientific research has been conducted specific to the hip region in a competitive surfing cohort

Purpose

- The primary aim of this study was to profile the isometric hip abductor and adductor muscle strength of competitive surfers using a clinical assessment method
- A secondary aim was to compare the surfing cohort to a group of age matched recreational athletes

Methods

- A cross sectional study design was used
- A total 14 elite surfers were recruited from the Surfing Australia High Performance Centre (4 females)
- 21 recreational athletes were recruited from Bond University (4 females)

- Hip abduction and adduction isometric strength was measured using a hand-held dynamometer (Commander PowerTrack II™ (Nihon Medix, Matsudo, Japan))



- Abduction (Image A) and Adduction (Image B) strength was measured in supine
- A belt was used to stabilize the participants pelvis
- The examiner maintained a stride stance with the testing elbow positioned against their own iliac spine to limit any movement of the hand-held dynamometer
- The hand-held dynamometer was positioned 5cm proximal to the medial / lateral malleolus



- Two maximal isometric contractions (separated by 10 seconds) were sustained for 3 seconds for each movement and repeated on each side
- The testing order was randomised to test side
- To calculate Torque (Newton/ meters) the lever arm was measure from the Greater Trochanter to 5cm proximal to the malleoli
- Normalised force (newtons per kilogram, N/kg) and torque (newton meters per kilogram, Nm/kg) were derived and compared between groups

Results cont.

Table 1: Demographics


	Competitive Surfers	Recreational Athletes
Age (years)	24.5 ± 6.5	25.3 ± 2.7
Weight (kg)	71.3 ± 8.3	80.0 ± 12.4*
Land Based Training (hours)	4.7 ± 2.6	5.9 ± 2.5
Weekly Surfing Frequency (hours)	11.6 ± 4.5	-

*The recreational athlete group were significantly (p=0.03) heavier (71.3±8.3 vs. 80.0±12.4 kg) and therefore strength values were normalised based on body weight.

Table 2: Isometric hip abduction and adduction strength profile of competitive surfers (n = 14)

	Hip Abduction	Hip Adduction
Force (N)	146.7 ± 32.9	131.7 ± 29.9
Normalised Force (N/kg)	2.0 ± 0.3	1.8 ± 0.3
Torque (Nm)	113.9 ± 26.6	102.4 ± 24.9
Normalised Torque (Nm/kg)	1.6 ± 0.2	1.4 ± 0.2

Normalised isometric hip adduction force and torque values for competitive surfers and recreational athletes



Results cont.

- No significant differences were identified for Nm/kg and N/kg for abduction between the competitive surfers and the recreational athletes (1.6 vs. 1.6 Nm/kg; 2.0 vs. 2.1 N/kg, respectively)
- As seen in the bar graph, the competitive surfer group displayed significantly higher scores for normalised adduction when compared to the recreational athletes (1.4 vs. 1.2 Nm/kg; p<0.01 and 1.8 vs. 1.6 N/kg; p=0.03, respectively)
- The surfer group exhibited an adduction to abduction strength ratio of 0.9 which was significantly (p<0.01) higher than the recreational athlete group (0.8).

Discussion

This study presents an isometric strength profile for the hip abductors and adductors specific to a competitive surfing cohort.

Results indicate that the competitive surfing cohort had significantly stronger adductors and consequently a higher adduction/ abduction ratio when compared with the recreational athlete group.

Implications

This research provides preliminary information specific to hip abduction and adduction strength in a competitive surfing cohort. The data presented provides a baseline specific to competitive surfers who are involved in both land based training and surfing.

The findings may guide rehabilitation or be used as a bench mark to highlight deficiency. Further research is needed to determine how strength at the hip plays a role in injury prevention and performance enhancement.

References

- Inada, K., Matsumoto, Y., Kihara, T., Tsuji, N., Netsu, M., Kanari, S., ... Arima, S. (2018). Acute injuries and chronic disorders in competitive surfing: From the survey of professional surfers in Japan. *Sports Orthopaedics and Traumatology*, 34(3), 256-260. doi:https://doi.org/10.1097/JSM.0000000000000596
- Nathanson, A., Bird, S., Dao, L., & Tam-Sing, K. (2007). Competitive Surfing Injuries. *The American Journal of Sports Medicine*, 35(1), 113-117. doi:10.1177/0363546506293702
- Santos, T. R. T., Oliveira, B. A., Ocarino, J. M., Holt, K. G., & Fonseca, S. T. (2015). Effectiveness of hip muscle strengthening in patello014.0089