

Bond University
Research Repository



The Role of Open Access in Reducing Waste in Medical Research

Glasziou, Paul

Published in:
PLoS Medicine

DOI:
[10.1371/journal.pmed.1001651](https://doi.org/10.1371/journal.pmed.1001651)

Licence:
CC BY

[Link to output in Bond University research repository.](#)

Recommended citation(APA):
Glasziou, P. (2014). The Role of Open Access in Reducing Waste in Medical Research. *PLoS Medicine*, 11(5), [e1001651]. <https://doi.org/10.1371/journal.pmed.1001651>

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

For more information, or if you believe that this document breaches copyright, please contact the Bond University research repository coordinator.



Editorial

The Role of Open Access in Reducing Waste in Medical Research

Paul Glasziou*

Bond University, Robina, Australia

Twenty years ago an editorial by Doug Altman in the *BMJ* [1], “The Scandal of Poor Medical Research”, decried the poor design and reporting of research, stating that “huge sums of money are spent annually on research that is seriously flawed through the use of inappropriate designs, unrepresentative samples, small samples, incorrect methods of analysis, and faulty interpretation”. Since then, change has been gradual, while the list of problems has lengthened, and documentation of their magnitude has accumulated. Recent years, however, have seen a crescendo of concern. Public awareness has been accelerated with the publication of Ben Goldacre’s *Bad Pharma* [2], which clearly articulated the problems posed by biased non-publication and reporting of pharmaceutical research. Wider awareness of these issues helped spark the AllTrials campaign (<http://www.alltrials.net/>), which asks for “all trials registered; all results reported”. Of course, the problems of poor design and reporting, as well as selective non-publication, extend well beyond drug trials to most areas of research: drug and non-drug, basic and applied, interventional and observational, animal and human. A 2009 paper in *The Lancet* [3] estimated that three problems—flawed design, non-publication, and poor reporting—together meant over 85% of research funds were wasted, implying a global total loss of over US\$100 billion per year. This year, a follow-up series [4] more extensively documented this wastage, confirming the earlier estimate, but adding details and a series of more explicit recommendations for action.

The waste sounds bad, but the reality is worse. The estimate that 85% of research is wasted referred only to activities prior to the point of publication. Much waste clearly occurs after publication: from poor access, poor dissemination, and poor uptake of the findings of research. The development of open access to research [5] is important to reduce this post-publication waste. Poor access—including paywalls, restrictions on re-publication and re-use, etc.—limits both researcher-to-researcher and

researcher-to-clinician communications. As *PLOS Medicine* editorial leaders pointed out in a PubMed Commons response to the *Lancet* series [6], open access is more than free access and includes “free, immediate access online; unrestricted distribution and re-use rights in perpetuity for humans and technological applications; author(s) retains rights to attribution; papers are immediately deposited in a public online archive, such as PubMed Central” [7]. Globally, the most important access problem is arguably due to language barriers, and with the growth of research in non-English-speaking countries, particularly China, this problem is likely to grow. Language barriers make even free-access research unusable, but by eliminating restrictions on re-publication and re-use, open access can at least reduce barriers to translation.

Solving the problems of pre-publication waste and post-publication access could hugely accelerate medical research. Even the complete solution of these problems, however, would be insufficient to close the research–practice gap. Paradoxically, the plethora of research is itself a barrier to its use. A recent analysis of trials and reviews by specialty found an unmanageable scatter of research [8]. For example, in neurology the annual output was 2,770 trials across 896 journals, and 547 systematic reviews across 292 journals. So, in addition to access, clever systems of synthesis, filtering, findability, and usability are needed if the users of research are to cope with this information deluge [9]. The enormous marketing budgets of pharmaceutical companies demonstrate the importance they

place on investing resources in getting the message of their research to decision makers. Unfortunately, little such investment is made in non-commercial research, and this research is consequently neglected. This concern has led to the development of different approaches given names such as “evidence-based medicine”, “knowledge translation”, and “implementation science”.

To get full value from research investment, we need to reduce both the annual US\$100 billion of pre-publication (research production) waste and the unquantified cost of post-publication (research dissemination) barriers (Figure 1). Open access will not in itself fix the problems of poor research question selection, poor study design, selective non-publication, or poor or biased reporting, but these can be ameliorated considerably through appropriate editorial policies and peer review processes. Open-access medical journals must maintain particularly high standards for these processes in order to avoid merely increasing access to a biased selection of (often flawed) research. At the same time, improving research quality but keeping access restricted would mean continued waste in the use and uptake of good science.

“As the system encourages poor research,” wrote Altman in 1994 [1], “it is the system that should be changed. We need less research, better research, and research done for the right reasons.” To that must be added a need for research that is communicated effectively to those who need it. If over a 100

Citation: Glasziou P (2014) The Role of Open Access in Reducing Waste in Medical Research. *PLoS Med* 11(5): e1001651. doi:10.1371/journal.pmed.1001651

Published: May 27, 2014

Copyright: © 2014 Paul Glasziou. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Funding: PG is supported by NHMRC grant 0527500. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Competing Interests: The author declares that no competing interests exist.

* E-mail: Paul_Glasziou@bond.edu.au

Provenance: Commissioned; not externally peer reviewed




Figure 1. Chain of research production and dissemination.
doi:10.1371/journal.pmed.1001651.g001

billion dollars of medical research money were being wasted by corruption, the public and political outcry would be overwhelming. That resources of this magnitude are being wasted through incompetence and inattention should

be seen as a similar scandal. Badly designed and poorly thought through systems of research and dissemination subtract massively from global human health: they demand attention—and action.

Author Contributions

Wrote the first draft of the manuscript: PG. Contributed to the writing of the manuscript: PG. ICMJE criteria for authorship read and met: PG. Agree with manuscript results and conclusions: PG.

References

- Altman DG (1994) The scandal of poor medical research. *BMJ* 308: 283.
- Goldacre B (2012) *Bad pharma: how drug companies mislead doctors and harm patients*. London: Fourth Estate.
- Chalmers I, Glasziou P (2009) Avoidable waste in the production and reporting of research evidence. *Lancet* 374: 86–89. Available: <http://www.thelancet.com/journals/lancet/article/PIIS0140-6736%2809%2960329-9/fulltext>. Accessed 23 April 2014.
- Macleod MR, Michie S, Roberts I, Dimagl U, Chalmers I, et al. (2014) Biomedical research: increasing value, reducing waste. *Lancet* 6736: 1–6.
- Joseph H (2013) The open access movement grows up: taking stock of a revolution. *PLoS Biol* 11: e1001686. Available: <http://www.plosbiology.org/article/info%3Adoi%2F10.1371%2Fjournal.pbio.1001686>. Accessed 23 April 2014.
- Glasziou P, Altman DG, Bossuyt P, Boutron I, Clarke M, et al. (2014) Research: increasing value, reducing waste 5 Reducing waste from incomplete or unusable reports of biomedical research. *Lancet* 383: 267–276.
- Barbour V, Peiperl L (2014) Comment on PMID 24411647: Reducing waste from incomplete or unusable reports of biomedical research. *PubMed Commons*. Available: <http://www.ncbi.nlm.nih.gov/myncbi/ginny.barbour.1/comments/>. Accessed 28 April 2014.
- Hoffmann T, Eructi C, Thorning S, Glasziou P (2012) The scatter of research: cross sectional comparison of randomised trials and systematic reviews across specialties. *BMJ* 3223: 1–9. doi:10.1136/bmj.e3223
- Glasziou P, Haynes RB (2005) The paths from research to improved health outcomes. *Evid Based Med* 10: 4–7.