

Bond University
Research Repository



An assessment of current antenatal care practices and identification of modifiable risk factors for prematurity and low birth weight infants in pregnancy in Solomon Islands

Cafaro, James; Randle, Elloise; Wyche, Penelope; Higgins, Mark; Fink, James; Jones, Peter

Published in:
Rural and Remote Health

DOI:
[10.22605/RRH3230](https://doi.org/10.22605/RRH3230)

Licence:
CC BY

[Link to output in Bond University research repository.](#)

Recommended citation(APA):

Cafaro, J., Randle, E., Wyche, P., Higgins, M., Fink, J., & Jones, P. (2015). An assessment of current antenatal care practices and identification of modifiable risk factors for prematurity and low birth weight infants in pregnancy in Solomon Islands. *Rural and Remote Health*, 15(3), 1-3. Article 3230.
<https://doi.org/10.22605/RRH3230>

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

For more information, or if you believe that this document breaches copyright, please contact the Bond University research repository coordinator.

SHORT COMMUNICATION

An assessment of current antenatal care practices and identification of modifiable risk factors for prematurity and low birth weight infants in pregnancy in Solomon Islands

J Cafaro¹, E Randle¹, P Wyche¹, M Higgins¹, J Fink², PD Jones¹

¹Bond University, Robina, Gold Coast, Queensland, Australia

²Queensland Health, Gold Coast University Hospital, Queensland, Australia

Submitted: 3 June 2014; Revised: 13 January 2015; Accepted: 7 May 2015; Published: 10 September 2015

Cafaro J, Randle E, Wyche P, Higgins M, Fink J, Jones PD

An assessment of current antenatal care practices and identification of modifiable risk factors for prematurity and low birth weight infants in pregnancy in Solomon Islands

Rural and Remote Health 15: 3230. (Online) 2015

Available: <http://www.rrh.org.au>

Solomon Islands is a sovereign country located 2000 km north-east of Brisbane and of the eastern tip of Papua New Guinea. The country consists of an archipelago of hundreds of islands spread over 11 000 km². Makira-Ulawa is a remote province 200 km east of Honiara. It has a rapidly growing population of approximately 40 000 with an estimated fertility rate of 6.2 births per woman. Solomon Islands is a poor country with over 50% of its gross domestic income dependent on foreign aid. There are very few resources available for health care with just over AU\$100 available for each person for health care or less than 2% of what is spent on health care in Australia.

KiraKira is the provincial capital of Makira-Ulawa province. It is a small rural township of approximately 3000 people¹. It

has the only hospital in the province, which is staffed by a single government-employed medical doctor, supported by approximately 40 nurses and midwives. There has been no previous published data on health outcomes in Solomon Islands outside of the National referral hospital in Honiara.

We performed a retrospective audit of hospital records of all births at KiraKira Hospital from 01/01/2011 to 19/09/2013 to calculate the incidence of low birth weight (<2.5 kg), prematurity (<37 weeks gestation) and proportion of women receiving antenatal care. In addition we undertook a cross sectional study of women presenting for routine antenatal care during a 2 week period in September 2013 that included urine analysis using Multistix 10SG (Elkart, IN, USA). Bacteriuria was defined as having >2 white cells identified in



the urine. It was classified as symptomatic if there were associated symptoms including fever, abdominal and flank pain and dysuria. The study was approved by the Bond University Ethics Committee (approval no. BUREC RO 1708).

The results for the audit are shown in Table 1 and those for the cross sectional audit are shown in Table 2. These tables include recent statistics from the Australian Institute of Health and Welfare for comparison².

This study has confirmed that there are high rates of teenage pregnancy, prematurity, low birth weight and perinatal mortality in KiraKira. The perinatal mortality rate of 31 per 1000 is more than double the rate of 14 per 1000 reported for Solomon in 2012 as part of the Millennium Development Goals project³. Aside from the obvious challenge of one doctor trying to provide health care for a community of 40 000 this variation is of concern. A possible explanation for this difference is that data from remote provinces was not available to researchers and the lower reported figure is based on outcomes achieved at the national referral hospital. If that were the case then the overall official figures for Solomon Islands are overly optimistic.

An interesting observation in the cross sectional study was the finding that 59% of pregnant women were found to have bacteriuria; 25% of these with symptoms suggestive of a urinary tract infection and the remaining 33% being classified as having asymptomatic bacteriuria. The expected rate of asymptomatic bacteriuria and urinary tract infection during pregnancy in developed countries such as Australia is between 2 and 10%. Such high rates have previously been described in the developing world with one study from Nigeria finding 87% of women had asymptomatic bacteriuria⁴.

Asymptomatic bacteriuria is an important finding due to the complications it poses to the pregnant mother and the unborn child. Its presence carries a risk of pyelonephritis of up to 20–40% in the mother. It is also associated with increased prematurity and low birth weight. The United States of

America Preventative Task Force recommends screening at 12–16 weeks⁵, as an intervention has been shown to greatly reduce low birth weight and prematurity. The results of this small cross sectional study provide data suggesting that screening the urine using a dipstick should become part of antenatal care in KiraKira.

Seventy-five percent of women who deliver in KiraKira access antenatal care. The main area where there could be improvement would be encouraging women to present earlier in their pregnancy. Our study found that only 6% of women presented for antenatal care within the first trimester. This compares to two-thirds of women visiting a doctor in Australia before the end of the first trimester.

Three-quarters of the pregnant women in this cross sectional study admitted to chewing betel nut. Betel nut is described as the fourth most used psychoactive substance globally (after tobacco, alcohol and caffeine). In addition to proven carcinogenic effects the chewing of betel nut during pregnancy has been shown to increase the incidence of low birth weight infants and also to increase rates of miscarriage, stillbirth, prematurity and fetal malformation⁶. There appeared to be lack of awareness among the women of KiraKira that betel nut poses any risks for the fetus. In KiraKira, betel nut use is nine times more prevalent than alcohol and tobacco use during pregnancy. Mothers should be encouraged to stop chewing betel nut during pregnancy as this habit is more than likely contributing to the very high rates of prematurity and low birth weight.

In summary, the findings of this study suggest that there are three low cost interventions that could be implemented in KiraKira that could reduce the rate of prematurity and low birth weight infants. First, pregnant women should be encouraged to attend for their first antenatal visit in the first trimester of pregnancy. Second, mothers should be screened for bacteriuria using a dipstick for urine analysis. Finally, pregnant mothers should be encouraged to stop chewing betel nut.



Table 1: Deliveries at KiraKira Hospital 2011–2013 compared with data from all Australian deliveries in 2011

Measure	Outcome	Australian comparison ²
Deliveries	1233	295 070
Low birth weight (<2.5 kg)	150 (12.2%)	6.3%
Prematurity (<37 weeks)	293 (23.8%)	8.3%
Perinatal deaths	38 (31 per 1000)	9.1 per 1000
Maternal deaths	1	0.05 per 1000
Caesarian sections	22 (1.8%)	33.0%
Instrumental deliveries	134 (10.9%)	12.0%
Episiotomy at delivery	221 (17.9%)	18.0%
>4 antenatal visits	75%	94.4%
Median date of first antenatal visit	26 weeks	12 weeks
First antenatal visit by 14 weeks	74 (6%)	65.5%
Teenage pregnancies	144 (14%)	3.7%

Table 2: Sixty-three Melanesian women presenting for antenatal care during September 2013 in KiraKira Hospital compared with data for Australian women in 2011

Measure	Outcome	Australian comparison ²
Number of women	63	Not available
Average age (years)	27	30
Teenagers	8 (13%)	3.7%
Primiparous	16 (25%)	43%
Grand multi >5	11 (17.4%)	3.6%
Asymptomatic bacteriuria	21 (33%)	2–10% ³
Symptomatic bacteriuria/urinary tract infection	16 (25%)	2–10%
Tobacco during pregnancy	6 (9.5%)	13.2%
Betel nut during pregnancy	48 (76%)	Not applicable

References

1. UNICEF. Solomon Island, Statistics. (Online) 2013. Available: http://www.unicef.org/infobycountry/solomonislands_statistics.html (Accessed 2 August 2014).
2. Li Z, Zeki R, Hilder L, Sullivan EA. *Australia's mothers and babies 2011*. Preinatal statistics series no. 28 Cat. no. PER59. Canberra: AIHW National Perinatal Epidemiology and Statistics Unit, 2013.
3. Lozano R. Progress towards millennium development goals 4 and 5 on maternal and child mortality: an updated systematic analysis. *The Lancet* 2011; **378**: 1139-1165.
4. Macejko A. Asymptomatic bacteriuria and symptomatic urinary tract infections during pregnancy. *Urologic Clinics of North America* 2007; **34**: 35-42.
5. Patterson TF. Bacteriuria in pregnancy. *Infectious Disease Clinics of North America* 1987; **1**: 807-822.
6. Yang M-S, Lee C-H, Chang S-J, et al. The effect of maternal betel quid exposure during pregnancy on adverse birth outcomes among aborigines in Taiwan. *Drug and Alcohol Dependence* 2008; **95**: 134-139.