

Bond University  
Research Repository



## The effectiveness of nutrition education programmes on improving dietary intake in athletes: A Systematic Review

Boidin, Aimee; Tam, Ryan; Mitchell, Lachlan; Cox, Gregory R.; O'Connor, Helen

*Published in:*  
British Journal of Nutrition

*DOI:*  
[10.1017/S0007114520003694](https://doi.org/10.1017/S0007114520003694)

*Licence:*  
Other

[Link to output in Bond University research repository.](#)

*Recommended citation(APA):*  
Boidin, A., Tam, R., Mitchell, L., Cox, G. R., & O'Connor, H. (2021). The effectiveness of nutrition education programmes on improving dietary intake in athletes: A Systematic Review. *British Journal of Nutrition*, 125(12), 1359-1373. Article 0007114520003694. <https://doi.org/10.1017/S0007114520003694>

### General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

For more information, or if you believe that this document breaches copyright, please contact the Bond University research repository coordinator.

The effectiveness of nutrition education programs on improving dietary intake in athletes: A Systematic Review

Aimee Boidin<sup>1</sup>, Ryan Tam<sup>2</sup>, Lachlan Mitchell<sup>3</sup>, Gregory R Cox<sup>1</sup>, Helen O'Connor<sup>2,4</sup>

<sup>1</sup>Faculty of Health Sciences and Medicine, Bond University, Queensland, Australia. 14 University Drive, Robina, Queensland, Australia

<sup>2</sup>Faculty of Health Sciences, The University of Sydney, NSW, Australia. 75 East Street, Lidcombe, NSW, Australia.

<sup>3</sup>National Nutrition Surveillance Centre, School of Public Health, Physiotherapy and Sport Science, University College Dublin, Ireland. Belfield, Dublin 4, Ireland.

<sup>4</sup>Charles Perkins Centre, The University of Sydney, NSW, Australia. John Hopkins Drive, Camperdown, NSW, Australia.

Corresponding Author: Lachlan Mitchell

[lachlan.mitchell@ucd.ie](mailto:lachlan.mitchell@ucd.ie)

National Nutrition Surveillance Centre, University College Dublin, Ireland

Keywords: athletes, nutrition, education, program, dietary intake

Word Count: 5204

Running head: Nutrition education to improve dietary intake in athletes



This peer-reviewed article has been accepted for publication but not yet copyedited or typeset, and so may be subject to change during the production process. The article is considered published and may be cited using its DOI

10.1017/S0007114520003694

The British Journal of Nutrition is published by Cambridge University Press on behalf of The Nutrition Society

**Abstract**

Nutrition education programs for athletes aim to enhance nutrition knowledge and more importantly support positive dietary change to enhance performance, health and well-being. This systematic review assessed changes in the dietary intakes of athletes in response to nutrition education programs. A search was conducted which included studies providing quantitative dietary intake assessment of athletes of any calibre aged between 12-65y in response to a nutrition education program. Standardized differences (effect sizes) were calculated (when possible) for each dietary parameter. The search yielded 6285 papers with 22 studies (974 participants (71.9% female)) eligible for inclusion. Studies described athletes competing at high school (n=3) through to college level or higher (n=19). Study designs were either single-arm with an intervention only group (12 studies; n=241) or double-arm including an intervention and control group (10 studies; n=689). No control groups received an alternative or 'sham' intervention. Face-to-face lectures (9/22) and individual nutrition counselling (6/22) were the most common education interventions. Non-weighed, three-day diet records (10/22) were the most frequently utilised dietary assessment method. Although 14/22 studies (n=5 single and n=9 double) reported significant change in at least one nutrition parameter, dietary changes were inconsistent. Poor study quality and heterogeneity of methods prohibits firm conclusions regarding overall intervention success or superior types of educational modalities. Of note, carbohydrate intakes 'post-intervention' when assessed often failed to meet recommended guidelines (12/17 studies). Given the substantial investment made in nutrition education interventions with athletes, there is a need for well-designed and rigorous research to inform future best practice.

*Registration:* Prospero [CRD42018083952](https://doi.org/10.1111/1452003694).

*Key Words:* intervention, energy, macronutrient, diet, quality

## Introduction

Nutrition education interventions aim to help athletes align their dietary intake with current sports nutrition guidelines <sup>(1; 2)</sup>. These interventions, incorporated into many elite institute, professional, or collegiate sports programs, vary widely from individual consultations to group education; some incorporate practical skills such as cooking or shopping <sup>(2)</sup>. Despite the time and cost associated with athlete education, there is limited information on how these nutrition interventions influence dietary intake. A number of reviews have evaluated the level of nutrition knowledge in athletes <sup>(2; 3)</sup>, and one published recently also reports on how this improves with nutrition education <sup>(4)</sup>. Generally, the level of athlete nutrition knowledge varies widely amongst athlete groups <sup>(2; 3)</sup>, but improves, at least in the short term, even after brief nutrition education interventions <sup>(4)</sup>. However, due to the wide range of knowledge assessment tools with limited validation the authors were unable to identify the most effective nutrition education modality to improve nutrition knowledge in athletes <sup>(4)</sup>.

Although it is often assumed that greater nutrition knowledge results in better dietary intake, evidence suggests other factors (e.g. taste, cost, convenience) are equally important <sup>(5)</sup>. Athletes may fail to meet their dietary intake due to the challenges in navigating specific barriers such as available time for food selection and preparation due to high daily training commitments, suppressed appetite between training sessions, food culture and traditions unique to their sport, religious or environmental considerations and the body composition and physique requirements required for success <sup>(1; 2)</sup>. Observations made by coaches and sports dietitians indicate that athletes who possess confidence in their nutrition knowledge are more likely to incorporate this knowledge into their lifestyle by choosing appropriate foods to match their sport <sup>(1; 2)</sup>. However, the effectiveness of different types of nutrition education interventions to promote change in the dietary intake of athletes has yet to be evaluated.

Therefore, the primary aim of this systematic review was to investigate the effectiveness of nutrition education interventions on change in dietary intake in athletes. A secondary aim was to compare the effectiveness of different education delivery modalities (e.g. group versus individual or in person to virtual education modalities). Given the substantial professional and institutional investment in nutrition education for athletes, a comprehensive evaluation regarding its effectiveness to modify dietary intake is relevant to informing future best practice.

## **Method**

### *Search Strategy*

The systematic literature search to identify studies was conducted by one researcher (AB) from the earliest record until June 2019. Databases searched included PubMed, Embase, CINAHL and SPORTDiscus (EBSCOHost) using key words and controlled vocabulary, ‘athletes’, ‘sport’, ‘nutrition’, ‘diet’, ‘food’, ‘education\*’, ‘programs\*’, ‘counsel\*’, ‘health education’, ‘intervention’, ‘strategy\*’, ‘curriculum’, ‘lesson\*’, ‘class’, ‘workshop\*’, ‘program evaluation’, ‘dietary intake’, ‘energy intake’, ‘energy balance’, ‘behav\*’, ‘feeding behavior’, ‘intake\*’, ‘consumption’, ‘habits’, ‘patterns’, ‘practices’, ‘dietetics’, ‘food habits’, ‘caloric intake’. The full electronic search strategy is presented in Supplementary Figure 1. The search strategy was complemented by a hand search of studies referenced in similar reviews and included studies. This systematic review was registered on Prospero ([CRD42018083952](https://doi.org/10.1017/S000714520003694)) and reported according to PRISMA guidelines <sup>(6)</sup>.

### *Eligibility Criteria*

Eligible study designs included randomised controlled trials (RCTs), quasi-experimental and pre-post intervention studies. Abstracts and studies not reported in English were excluded.

Studies of athletes, male and female aged between 12-65y from all sports, and athletic levels were eligible for inclusion. Interventions including individual/group counselling/education, in-person or virtual (for example online/DVD) modalities were eligible if the primary outcome, change in dietary intake (energy, macronutrients, micronutrients and/or food groups, diet quality/index), was reported quantitatively.

### *Selection of studies and data extraction*

After the search was conducted and duplicates removed, manuscript titles and abstracts were screened independently by two authors (AB, RT). Disagreements were resolved via discussion with a third author (GC). Full-texts of all potentially eligible studies were reviewed independently by two authors (AB, RT). After identification of eligible full-text articles, data were extracted in duplicate (AB, RT). When relevant, paper authors were contacted (AB) and requested to supply additional/missing information. A computer program (WebPlotDigitizer, Version 3.9, Austin, Texas, USA) was used to calculate the mean and standard deviation of data reported in figures <sup>(7)</sup>.

Nutritional information extracted pre- and post-intervention included mean energy (kcal), macronutrient (g), micronutrient (mg/ $\mu$ g), food group <sup>(7)</sup>, and KIDMED diet quality/index (score of +1 or -1 totalling to a maximum 12) <sup>(7)</sup>. All nutrients were converted to SI units if reported otherwise. To manage table size in this review, only the micronutrients iron and calcium were tabulated. Changes to other micronutrients were discussed only in the text. To compare changes, within- and when relevant, between-group (double-arm studies) effect size (ES) was calculated when relevant data (mean, SD or SEM) were provided. The Hedges g (random model) ES was calculated using extracted data (mean, SD, sample size) in the Comprehensive Meta-Analysis Version 2 software (Biostat, 2005, Englewood, IL, USA), and were considered trivial (0.0-0.19), small (0.2-0.49), moderate (0.5-0.79), or large ( $\geq$ 0.8) <sup>(8)</sup>.

*Calculation of energy, macronutrient and micronutrient values across studies*

To assess the adequacy of reported dietary intakes, the recommended food group or recommended dietary intake/recommended daily allowance (RDI/RDA) guidelines specific to the country were used, and for studies using diet quality/indexes, the recommended ranges for these parameters cited in the paper (a 16-item Mediterranean Diet Quality Index). To assess the adequacy of energy intake, the types of methodology used to determine the estimated energy requirement (EER) of the participants could range from published energy requirement algorithms, general population energy requirement recommendations for 'active' or 'very active' individuals, or literature reported values for the sport/activity.

The classification of adequacy for macronutrients varied depending on the specific guidelines used in the paper. Historically, scientific opinion on this ranged over the period of time the studies were conducted, initially being recommended as a proportion of daily energy (protein 10-15%; fat 20-30% and carbohydrate  $\geq$ 50-60% of daily energy intake), later recognition that g/kg/d recommendations were more appropriate (protein 1.0-2.0g/kg/d and carbohydrate 3-12 g/kg/d<sup>(9)</sup>). Where possible, g/kg/d calculations were undertaken if not reported in the paper to assist in evaluating the appropriateness of the older, percent of energy recommendations. Micronutrient adequacy was based on the respective country RDI/RDA values. Historically, mean intakes which did not meet the RDI/RDA, but were a set proportion ranging between 70-100% of the RDI/RDA for micronutrients were reported as 'likely' to be adequate in older nutrition literature, although this is not an established way of reporting dietary adequacy.

*Quality Assessment*

Study quality was independently assessed in duplicate by three researchers (AB all papers, HO and RT shared) using a modified version of the Downs and Black risk of bias rating tool<sup>(10)</sup>. The original tool consists of twenty-seven items that examine data reporting, external and

internal validity, including bias and statistical power. In single-arm study designs, 20 of the 27 items that logically applied were used. In double-arm study designs, 24 items were used. Item eighteen addressed the validity and reliability of the tools used to measure dietary intake. This item was assessed as two parts as previously described by Spronk *et al.* <sup>(5)</sup>, where one point was awarded for appropriate choice of dietary method and another point for appropriate application. This approach was adopted with consideration of items 7.2-7.5 in the Academy of Nutrition and Dietetics Quality Criteria Checklist to accompany the Downs and Black tool and allowing for clarification when assessing quality ratings. Downs and Black score ranges were provided with corresponding quality levels based on the scoring methodology of the Downs and Black checklist. Quality levels for single arm studies included; excellent (20-21), good (15-19), fair (12-14) and poor (<12) and for double arm studies excellent (23-25), good (18-22), fair (14-17) and poor (<14). The maximum scores were 21 and 25 for single-arm and double-arm studies, respectively. Disagreement between authors was discussed to achieve consensus.

## **Results**

### *Literature Search and Study Selection*

A total of 8004 articles were identified through the database search. After removal of duplicates, 6285 articles remained. Screening by title and abstract identified 36 articles for full-text review. After evaluation against inclusion and exclusion criteria, 21 articles were eligible for inclusion (Figure 1). One additional article was identified by hand-searching references of included papers.

### *Study Characteristics*

Study characteristics were summarised into single-arm (intervention group only) and double-arm (intervention and control groups) studies (Table 1). The sample size across all studies ranged from 7-210 athletes (80.5% female). Studies were conducted in the United States (US) (n=14), Europe (n=5), Iran (n=1), Malaysia (n=1) and Brazil (n=1). The majority of studies involved mixed sports (n=9), with one of these mixed sport studies involving two population groups (mixed sport and ballet dancers). Team sports were represented in seven studies including volleyball (n=3), soccer, baseball, softball, and handball (n=1 for all). Individual sports included swimming, ballet, track and field, canoeing, and wrestling (n=1 for all). The mean age of athletes across all studies was 19.8y. Athletic calibre included high school (n=3 studies), collegiate (n=4 studies), state (n=1 studies), national (n=10 studies), and international (n=4 studies) levels. Two studies involved athletes with physical disabilities <sup>(11; 12)</sup>.

### *Intervention Characteristics*

#### a) Nutrition Education Modality

Various nutrition education modalities were utilised across the 22 included studies. Face-to-face group lectures (n=8 studies) and individual nutrition counselling (n=6) were most commonly used. Other modalities included group workshops/activities (n=4) or mixed methods (n=5), including lectures and handouts or lectures and individual counselling.

#### b) Nutrition Education Topics

Half of the studies (n=11) incorporated a combination of nutrition topics including energy, macronutrient, micronutrient, and hydration principles; meal frequency and timing; and supplement use. Other topics incorporated food groups and dietary guidelines (n=2), nutrient recommendations (n=1), general sport nutrition principles (n=1), iron (n=1), Mediterranean

diet principles (n=1), and individual nutrition plans or weight control strategies (n=3). Self-efficacy, social cognitive or cognitive behavioural theory concepts to assist the athletes to make dietary change were included in 6/22 studies<sup>(13; 14)</sup>.

#### c) Duration and Frequency of Nutrition Education

The nutrition education interventions ranged from 2-39 weeks in duration, with two studies (one single-arm, one double-arm) incorporating a follow-up period (6-16 weeks) and reported retention of dietary changes<sup>(13; 14)</sup>. One study assessed athlete dietary intake changes across two seasons, with the education delivered during the second season, however failed to describe the duration and frequency of the education program<sup>(15)</sup>. Session number varied from three (3/22 studies), 4-7 (13/22 studies), to more than seven (6/22 studies) sessions across the intervention period. Session duration ranged from 10-120 minutes with total intervention time ranging from 60 to 720 minutes.

#### d) Nutrition Education Facilitator

Most of the interventions (17/22) were delivered by a qualified (or student) nutrition professional with expertise described as; 'dietitian' (n=10), 'sports or performance nutritionist/dietitian' (n=4), 'nutrition specialist/professional nutritionist' (n=2) and student dietitian (n=1). The remaining studies (5/22) failed to report facilitator expertise<sup>(12; 16; 17; 18)</sup>.

#### *Diet Methodology Characteristics*

Most studies (15/22) assessed dietary intake twice, immediately pre- and then post-intervention, with some (5/21) studies also conducting assessments during the intervention (data not extracted). Two studies did not disclose the number of dietary intake assessment sessions but did present pre- and post-intervention results<sup>(15; 19)</sup>. All studies utilised a valid diet method to collect nutrient intake, however only 10/22 studies appropriately applied the

methodology (i.e. appropriate sample size, population, duration, frequency, and nutrients assessed) <sup>(15; 19; 20; 21)</sup>.

### *Study Quality*

Methodological quality had a mean score of 12/21 (range 7-15) for single-arm studies and a mean score of 15/25 (range 11-20) for double-arm studies, representing poor study quality and fair study quality for single and double arm studies respectively (Supplementary Tables 1 and 2). Most reported their aims (21/22 studies), main outcomes (18/22 studies), described the main findings (19/22 studies), and used appropriate statistical tests (15/22 studies). The lowest ratings were for the recording of compliance to the education intervention and collection of dietary intake (5/22 studies). Ten of the studies appropriately applied the dietary methodology with limitations in application mostly due to small participant numbers and inappropriate measurement and number of collection days for specific micronutrients. For example, iron intake was reported after only 3-days of measurement in a food diary that requires up to 11 recording days <sup>(22)</sup>.

### *Single-Arm Studies Dietary Outcomes*

The single-arm studies (n=12) investigated the within-group changes in dietary intake pre- and post-nutrition intervention (Table 2a). Of these studies, 9/12 reported changes in energy intake and 10/12 changes in macronutrients using predominantly 3-day diet records (n=6 studies), with a smaller number using 7-day diet records (n=1 study), 24-hour recall (n=2 studies), or 72-hour recall (n=1 study). Micronutrient intake was reported in 4/12 studies using predominantly 3-day diet records (n=3 studies) and one using 24-hour recall (n=1 study). In 3/12 single-arm studies, food group intake was assessed <sup>(23)</sup>. Adherence to the Mediterranean Diet using a diet index (KIDMED) was assessed in 1/12 studies <sup>(23)</sup>.

Baseline dietary intakes did not always meet the nutrition targets identified by the researchers which were guided by their country's RDI/RDA or relevant sports nutrition literature at the time. In the case of energy, a range of methods were used to identify the appropriate EER for the athlete population investigated. This included use of general population-based energy requirements typically identifying a range of suitable energy intakes for 'active' or 'very active' individuals. Other studies used algorithms (e.g. Harris Benedict or Nelson Equations) or sport specific energy requirements reported in the literature to calculate the appropriate level of energy for the study population (based on age, sex, weight and height). Training loads of participants were only detailed in 2/9 studies assessing energy intake, contributing to the poor quality of studies since training information is essential to evaluate athlete energy requirements<sup>(9)</sup>.

Few (2/9) single-arm studies that assessed energy intake reported this to be within the researchers' targeted EER at baseline<sup>(21; 24)</sup>. Of those studies identified as having a low mean participant energy intake at baseline, 6/7 reported a significantly higher post-intervention energy intake (ES: 0.4-2.3;  $p \leq 0.05$ ), with only one<sup>(25)</sup> falling within the researchers' identified EER. The ES ranged from small to large. It is relevant to note that the study by Łagowska *et al.*<sup>(25)</sup> involved both ballet dancers and female-athlete sub-groups, and although both observed significant increases and large ES, only the female-athlete sub-group met energy intakes within researchers' range post-intervention (ES: 1.0-2.3;  $p \leq 0.001$ ). Conversely, the remaining study (1/7) with low baseline energy intake reported a close to significant decrease further below the researchers identified EER post-intervention (ES: -0.2;  $p = 0.05$ )<sup>(14)</sup>. All 7/9 studies where authors reported low baseline energy intakes assessed female athletes, with 1/7 including mixed-gender athletes<sup>(15)</sup>. In 2/9 studies where authors reported energy intake met the EER at baseline, both reported non-significant decreases which fell below the targeted EER post-intervention (ES: -0.4- -0.5)<sup>(21; 24)</sup>.

Carbohydrate intake was assessed in 10/12 single-arm studies. At baseline, mean carbohydrate intake was deemed to be appropriate by the researchers' (>50% of energy or 6-10g/kg/day) for the study population assessed in only 2/10 studies<sup>(14; 21)</sup>. In the 8/10 studies where carbohydrate intake was identified to be below requirements at baseline, the range in mean intake (when able to be calculated) was 3.1-5.2g/kg/d. Post-intervention, this increased to 3.8-7.0 g/kg/d, with 4/8 studies reporting significant increases in carbohydrate intake (ES: 0.7-2.4; p<0.05), yet still falling below the researchers' recommendations. In the 2/10 studies deemed to have adequate carbohydrate intake at baseline based on percent of energy, no significant changes in intake were observed post-intervention<sup>(14; 21)</sup>.

Protein intake was reported in 10/12 single-arm studies and was deemed adequate according to author recommendations in 5/10 studies at baseline (1.1-1.7g/kg/d, when able to be calculated)<sup>(15; 18)</sup>. In the remaining studies identified as having inadequate protein intake at baseline, three also reported protein as inadequate post-intervention (1.1-1.6g/kg/d), while the other two studies reported small to large increases to meet the targeted protein intake, ranging between 1.2-2.1g/kg/d (ES: 0.2-0.8)<sup>(15; 18)</sup>. In the 5/10 studies reporting adequate protein intake at baseline, 4/5 studies maintained an adequate protein intake post-intervention between 0.9-1.7g/kg/d, while one study reported a large reduction post-intervention, falling to the lower limit of the researchers' recommendations (1.0g/kg/d; ES: -0.8)<sup>(21)</sup>.

Dietary fat intake was reported in 9/12 single-arm studies. Of these, 5/9 were reported at baseline to exceed researchers' recommendations (20-30% of energy), with the remaining four meeting this recommendation at 23-28% of energy<sup>(14; 15; 21; 25)</sup>. Post-intervention, 3/5 studies that exceeded baseline recommendations reported non-significant decreases which fell within the recommended 20-30% of energy<sup>(12; 24; 26)</sup>, while the remaining two studies still exceeded 30% of energy post-intervention<sup>(18; 27)</sup>.

Iron and calcium were reported in 4/12 single-arm studies, with each of these meeting the researchers' targeted recommendation of >70% of the RDI/RDA at baseline. Two of these four studies reported non-significant increases for iron and calcium post-intervention (ES: iron 0.1-0.4; calcium 0.2-0.3) <sup>(12; 24)</sup>. The remaining 2/4 studies showed no change in iron intake <sup>(14; 27)</sup>, and a decrease in calcium intake to <70% of the RDI/RDA <sup>(14)</sup>. However, due to the fair quality of dietary methodology used for micronutrient assessment, these findings are questionable. Other micronutrients reported in 3/4 studies showed mostly adequate intakes at baseline and insignificant changes post-intervention <sup>(24; 27)</sup>. One study examined intake of dietary fibre, vitamin C and dietary cholesterol, with dietary cholesterol showing a significant reduction post-intervention <sup>(12)</sup>.

Change in food group consumption and adherence to the Mediterranean Diet was assessed in 3/12 and 1/12 single-arm studies, respectively. One of the studies assessing food group consumption <sup>(28)</sup>, reported athletes (adults and adolescents) as low, adequate or high at baseline and post-intervention based on the Brazilian Food Pyramid Guide. A significant reduction post-intervention in vegetable, fruit, and grain intake was reported in athletes classified as adequate at baseline for these food groups ( $p < 0.05$ ), falling below recommended intakes for vegetables. No significant changes were reported for dairy, or meat and egg intake. For athletes classified as low intake for vegetables, fruit, and dairy, significant increases were reported for these food groups ( $p < 0.05$ ), achieving the recommended intake for fruits. A significant reduction in meat and egg intake was observed in athletes classified as high intake at baseline ( $p < 0.05$ ), however these athletes remained above recommended intakes for this food group. No significant change in grain intake was observed in those classified as low at baseline <sup>(28)</sup>. Another study assessing food groups <sup>(21)</sup> reported participants met grains, vegetable and fruit groups at baseline, whilst dairy was reported as lower and meat serves higher than recommendations. Non-significant decreases were reported for vegetable, fruit,

and grain food groups post-intervention. Meat serves decreased to within the recommended intake (2.7 serves), while a large increase in milk intake to within the recommended range was observed post-intervention (2.5 serves; ES: 0.9)<sup>(21)</sup>. The remaining study assessing food groups reported low fruit and grain intake, and high meat, poultry and egg intake, pre-intervention. These food groups remained outside of recommended intakes post-intervention<sup>(27)</sup>. One study investigated adherence to the Mediterranean Diet, reporting a large increase in adherence post-intervention (ES: 1.0)<sup>(23)</sup>.

#### *Double-arm Studies Dietary Outcomes*

In the double-arm studies (n=10), energy intake was assessed in 7/10 studies, where 5/7 used 3-4-day food records, and 2/7 used 24-h recall (one study also used a 7-day food record). Macronutrients were measured in 8/10 studies (7/8 measuring carbohydrate and protein, and 8/8 dietary fat), where 4/8 used 3-4-day food records (one of these also used a 24-h recall), and 4/8 used 24-hour recalls (two of these combined this with a 4- or 7-day food dairy). One study measured food groups in addition to dietary fat using an FFQ<sup>(13)</sup>. Micronutrients were measured in 5/10 studies, using predominantly 3-day food records, with one study using a 7-day food record in addition to the 24h recall. Identification of nutrition targets was similar to the single-arm studies. Other micronutrients reported in 3/10 studies showed mostly adequate intakes at baseline and insignificant changes post-intervention.

The control groups in the double-arm studies received no placebo or sham intervention and generally only experienced minimal, non-significant within-group changes to dietary intake. There were generally more positive changes in dietary intake within the intervention groups, although not all changes were significant or in the direction targeted. Within- and between-group changes in dietary intake are presented in Table 2b. Sufficient information was

provided in 6/10 double-arm studies to perform between-group analyses <sup>(29; 30)</sup>, with 4/10 failing to provide sufficient data, reducing study quality <sup>(11; 13; 20; 31)</sup>.

Energy intake was compared between-groups in 4/10 studies. Energy intake was significantly increased relative to control in two studies (ES: 0.4-0.8;  $p < 0.05$ ) <sup>(29; 30)</sup>, while trivial, non-significant between-group differences in energy intake were observed post-intervention in two studies (ES: -0.2-0.1;  $p > 0.05$ ) <sup>(16; 32)</sup>.

Moderate to large increases in carbohydrate intake relative to control were reported in three (ES: 0.5-0.8;  $p > 0.05$ ) <sup>(16; 17; 32)</sup>, and a significant, large increase reported in two studies (ES: 1.4-1.8;  $p < 0.05$ ) <sup>(29; 30)</sup>. Similar variability was reported across studies with protein intake post-intervention. Two studies reported increases in intake relative to control (ES: 0.7-1.8) <sup>(29; 30)</sup>, two others reported small to moderate reductions relative to control (ES: -0.3- -0.7) <sup>(17; 32)</sup>. One study reported no between-group difference (ES: 0.0) <sup>(16)</sup>. A moderate increase in fat intake relative to control was reported in one study (ES: 0.4) <sup>(16)</sup>, while trivial to large reductions relative to control were found in four studies (ES: -0.1- -0.9) <sup>(17; 29; 30; 32)</sup>. Between-group differences in calcium intake were reported in 2/10 studies <sup>(16; 32)</sup>, with small increases reported in both studies (ES: 0.2-0.3). Small to moderate increases in iron intake compared to control were reported in three studies (ES: 0.3-0.7) <sup>(16; 32; 33)</sup>.

The remaining four double-arm studies did not conduct between-group analyses or did not provide sufficient information for ES to be calculated, contributing to the poor study quality and difficulty in drawing firm conclusions. Within-group analysis was performed in one of these studies, indicating a moderate reduction in energy intake (ES: -0.4), and a moderate increase in calcium intake (ES: 0.5), post-intervention <sup>(11)</sup>.

## Discussion

This is the first systematic review to evaluate the effectiveness of nutrition education programs on the dietary intake of athletes. Overall, the impact of nutrition education programs was varied. Given the range of intervention modalities and durations, the limitations in the dietary assessment methodologies employed, and the small number of studies conducting intervention-control comparison analyses, it is difficult to make firm conclusions as to the efficacy of the interventions or which interventions were best. Study quality was rated poor to fair-range, also indicating room for methodological improvement. Despite the significant investment in nutrition education of athletes, there is limited and generally low-quality evidence of the efficacy of interventions. Well designed and rigorous research application is needed in this area to inform future best practice.

Of the 22 studies, more than half (n=12) had a single-arm design which assessed dietary intake pre- and post-intervention (Table 1). While there was some evidence of intervention benefit, many of the ES were trivial or small and not statistically significant (Table 2a & 2b). Carbohydrate intake in particular often failed to meet the researchers' set targets, although with the age of the studies, many of these recommendations were outdated and likely inappropriate for the sports assessed <sup>(9)</sup>. Remarkably, intervention time was only able to be calculated for 3/12 studies and ranged from 180-300 min (3-5 h), across a 2- to 39-week duration.

In the double-arm studies (Table 1), none of the control arms used an alternative or 'sham' intervention to manage differences in group attention. Control groups generally experienced minimal changes, however, a limitation present in several studies was the failure to perform analysis between intervention and control groups (4/10 studies). Analysis was primarily conducted within-groups and rendered the use of the control group to qualitative comparison only. Across the double-arm studies, only 2/10 reported consistent, significant dietary improvements with ES in the large range <sup>(29; 30)</sup>. The remaining double-arm studies

demonstrated inconsistent dietary outcomes, with calculated ES varying in both direction and magnitude. Aligning with the single-arm study results, carbohydrate intake often failed to reach the researchers' targeted levels, although these may have been too high for the sports assessed<sup>(9)</sup>. In several instances, the control group outperformed the intervention group with respect to increasing nutrient intake. Most (9/10) double-arm studies provided sufficient detail on the intervention time which ranged from 60-720 min (1-12h) over 2-12 weeks.

Across all studies, most of the interventions focused on face-to-face group education, with some studies using resources such as handouts or emails to participants, and others using individual consults and meal plans. Most of the facilitators appeared to have training in nutrition/dietetics, although facilitator background/qualifications were not always provided. The heterogeneity of these factors in addition to the range in approaches used to assess dietary outcomes, makes it impossible to discern the overall effectiveness of interventions, nor which interventions are superior for improving dietary intake in athletes.

Given the heterogeneity and limited quality of the included studies in this review, few conclusions can be drawn as to the effect of nutrition education on dietary practices of athletes. A summary table in the form of a checklist has been constructed by authors to aid discussion of relative strengths and limitations of included studies (Table 3). This aims to succinctly critique the common flaws observed, while also guiding future nutrition education research. The checklist, Dietary Intake and Nutrition Education Reporting for Sports (DINERS), outlines factors which would inform stronger study design, methodology, and reporting. Appraisal of the literature included in this review identified four areas common to nutrition education intervention studies which require attention; participant characteristics, targeted dietary outcomes of the intervention (and underpinning rationale), intervention characteristics, and dietary methodology. These areas are described below, and in further detail in Supplementary Table 3.

### *Participant Characteristics*

Adequate participant description is necessary as different interventions may be more efficacious at different age stages or levels of athletic calibre. While most studies in this review reported age and variance (17/22), sex (21/22), sport (21/22) and athletic calibre (22/22), only two studies adequately described training characteristics (inclusive of frequency, intensity, duration and type)<sup>(9)</sup>. Detailed training information is necessary to evaluate athlete energy and nutrient requirements<sup>(9)</sup>. Moreover, to make sense of current nutrition guidelines, body weight and composition is important, especially how these may change over the intervention period. Only 10 and 13 studies reported on body composition and body weight, respectively. Detailed protocols and error (e.g. technical error of measurement) associated with the assessment of body composition is also an important inclusion in the methods of the paper. Only four papers in this review provided such information<sup>(9)</sup>.

### *Targeted Dietary Outcomes of the Intervention*

Defining intervention targets and desired dietary outcomes is essential for assessing intervention impact. Energy, macronutrient, and micronutrient targets were defined and justified by only three<sup>(9)</sup>, two<sup>(9)</sup>, and one<sup>(9)</sup> studies, respectively. Given athlete dietary goals can vary over a season or even a training period, the goals of the athletes may warrant lower or higher intake of energy, macronutrients or micronutrients, which needs to be explained with a clear underpinning rationale provided<sup>(34)</sup>.

### *Intervention Characteristics*

Intervention characteristics should be transparent. Details on the curriculum covered (17/22), the facilitator background and experience (12/22), modality of intervention used (individual or group, in-person or virtual) (19/22), as well as session duration, frequency and the total number of sessions (7/22) were not covered comprehensively across studies<sup>(34)</sup>. The total

amount of intervention minutes/hours should be provided. Participant attendance and compliance are also important and only half (11/22) of the studies examined these parameters. Underpinning behavioural theory and techniques are critical <sup>(35)</sup>. Clearly, there is a robust body of research outlining how behavioural support is needed to facilitate dietary change, yet many of the studies did not describe use of these methods. Process evaluation including how the participants perceived the intervention is important for determining how well the intervention was received, and this was only examined in two studies. Lastly, sustained dietary change is also relevant when evaluating intervention efficacy, and only three of the included studies assessed this <sup>(36; 37)</sup>. Clearly, financial or resource limitations may make follow-up after the intervention challenging.

### *Dietary Methodology*

A number of the included studies did not appropriately apply dietary methodology to their population, for example, 3-4-day food diaries are insufficient to examine intakes of micronutrients <sup>(38)</sup>. In this case, combining methods such as using a food frequency questionnaire (FFQ) for micronutrients with a food diary, which is better for quantifying energy and macronutrients, is recommended <sup>(36; 37)</sup>, although the level of agreement of FFQ is limited at the individual rather than group level <sup>(39)</sup>. The plausibility of data is also important and potential under-reporting should be assessed <sup>(22)</sup>. This requires capture of training loads; technology such as accelerometers, heart rate monitors or Global Positioning System trackers may be helpful as doubly-labelled water is unlikely to be available or affordable in most situations <sup>(22)</sup>. Another major flaw in included papers was the reporting of mean intakes of energy and macronutrients which may be skewed by athletes with extremely high or low intakes <sup>(40)</sup>. While mean intake can be useful, it is stronger to report the proportion of participants who reach the dietary targets as this facilitates a better assessment of intervention effectiveness in the cohort <sup>(40)</sup>. Similarly, in the case of micronutrients, it would be useful to check the proportion of participants meeting the Estimated Average Requirement and RDI/RDA for their age and sex but also report the proportion of participants meeting other sports nutrition specific targets which may be higher than those for the general population <sup>(9)</sup>. How the nutrient intake from dietary supplements contributes to intakes should also be detailed, and only two studies in this review reported on this <sup>(9)</sup>. Finally, the rigor of data capture and the background/experience of the researcher conducting the dietary analysis (reported in 5/22 studies) is also relevant to the quality of the dietary outcomes. Use of self-

report apps may make data capture easier but not necessarily accurate <sup>(41)</sup>.

Although the major limitation of this review is the quality of the literature that informs it, a major strength is the detailed synthesis of the studies and the construction of the DINERS checklist table (Table 3) which succinctly summarises the strengths and limitations of the included studies while also guiding future research practice. Nonetheless, the authors acknowledge some limitations of this review including studies excluded if published in languages other than English which poses a risk of publication bias. Further, no meta-analysis was performed due to the heterogeneity of included studies.

In conclusion, there is limited research informing the efficacy of nutrition education interventions in athletes, and what is available is of poor to fair quality, reporting varied outcomes. The findings of the review highlight 1) the requirement for ongoing nutrition education of athletes as they commonly report energy and carbohydrate intakes below recommendations, and 2) the importance of carefully planning interventions to ensure meaningful outcomes are aligned with sport specific nutritional requirements that can be clearly interpreted and subsequently reported by sports nutrition professionals and researchers. As nutrition education is a key strategy to enhance dietary intake in athletes and there is substantial investment in nutrition education interventions across the broader sporting context, there is a need for rigorous research in this area to inform best practice.

### **Acknowledgements and funding**

No funding was supplied in production of this manuscript. The authors declare no conflicts of interest. The study was designed by AB, GC, HOC; data were collected and analysed by AB, RT, HOC, LM; data interpretation and manuscript preparation was performed by all authors; all authors (except HOC) approved the final version of the manuscript.

This paper is dedicated to the memory of Associate Professor Helen O'Connor, who sadly passed during the preparation of this manuscript.

## References

1. Alaunyte I, Perry JL, Aubrey T (2015) Nutritional knowledge and eating habits of professional rugby league players: does knowledge translate into practice? *Journal of the International Society of Sports Nutrition* **12**, 18.
2. Heaney S, O'Connor H, Michael S et al. (2011) Nutrition Knowledge in Athletes: A Systematic Review. *International Journal of Sport Nutrition & Exercise Metabolism* **21**, 248-261.
3. Trakman GL, Forsyth A, Devlin BL et al. (2016) A Systematic Review of Athletes' and Coaches' Nutrition Knowledge and Reflections on the Quality of Current Nutrition Knowledge Measures. *Nutrients* **8**.
4. Tam R, Beck KL, Manore MM et al. (2019) Effectiveness of Education Interventions Designed to Improve Nutrition Knowledge in Athletes: A Systematic Review. *Sports Med* **49**, 1769-1786.
5. Spronk I, Kullen C, Burdon C et al. (2014) Relationship between nutrition knowledge and dietary intake. *The British journal of nutrition* **111**, 1713-1726.
6. Moher D, Shamseer L, Clarke M et al. (2015) Preferred reporting items for systematic review and meta-analysis protocols (PRISMA-P) 2015 statement. *Systematic Reviews* **4**, 1.
7. Rohatgi A (2015) WebPlotDigitizer.
8. Cohen J (1992) A power primer. *Psychol Bull* **112**, 155-159.
9. Thomas DT, Erdman KA, Burke LM (2016) American College of Sports Medicine Joint Position Statement. Nutrition and Athletic Performance. *Med Sci Sports Exerc* **48**, 543-568.
10. Downs SH, Black N (1998) The feasibility of creating a checklist for the assessment of the methodological quality both of randomised and non-randomised studies of health care interventions. *J Epidemiol Community Health* **52**, 377-384.
11. Rastmanesh R, Taleban FA, Kimiagar M et al. (2007) Nutritional Knowledge and Attitudes in Athletes With Physical Disabilities. *Journal of Athletic Training (National Athletic Trainers' Association)* **42**, 99-105.

12. Kandiah J (2000) Does Nutrition Education Help Lower Serum Lipid Levels in Athletes with Cerebral Palsy? *Palaestra* **16**, 44.
13. Doyle-Lucas AF, Davy BM (2011) Development and Evaluation of an Educational Intervention Program for Pre-professional Adolescent Ballet Dancers: Nutrition for Optimal Performance. *Journal of Dance Medicine & Science* **15**, 65-75.
14. Collison SB, Kuczmarski MF, Vickery CE (1996) Impact of nutrition education on female athletes. *American Journal of Health Behavior* **20**, 14-23.
15. Nowacka E, Leszczyńska T, Kopeć A et al. (2016) Nutritional behavior of Polish canoeist's athletes: The interest of nutritional education. *Science & Sports* **31**, e79-e91.
16. Abood DA, Black DR, Birnbaum RD (2004) Nutrition Education Intervention for College Female Athletes. *Journal of Nutrition Education & Behavior* **36**, 135-139.
17. Buffington BC, Melnyk BM, Morales S et al. (2016) Effects of an energy balance educational intervention and the COPE cognitive behavioral therapy intervention for Division I U.S. Air Force Academy female athletes. *Journal of the American Association of Nurse Practitioners* **28**, 181-187.
18. Rossi FE, Landreth A, Beam S et al. (2017) The Effects of a Sports Nutrition Education Intervention on Nutritional Status, Sport Nutrition Knowledge, Body Composition, and Performance during Off Season Training in NCAA Division I Baseball Players. *Journal of Sports Science & Medicine* **16**, 60-68.
19. Wenzel RK, Valliant MW, Chang Y et al. (2012) Dietary assessment and education improves body composition and diet in NCAA female volleyball players. *Topics in Clinical Nutrition* **27**, 67-73.
20. Chapman P, Toma RB, Tuveson RV et al. (1997) Nutrition knowledge among adolescent high school female athletes. *Adolescence* **32**, 437.

21. Wittkofski G (1996) Nutrition education using the food guide pyramid for female collegiate athletes. Ann Arbor, Mich.;; University Microfilms International.
22. Capling L, Beck KL, Gifford JA et al. (2017) Validity of Dietary Assessment in Athletes: A Systematic Review. *Nutrients* **9**.
23. Philippou E, Middleton N, Pistos C et al. (2017) The impact of nutrition education on nutrition knowledge and adherence to the Mediterranean Diet in adolescent competitive swimmers. *Journal of science and medicine in sport* **20**, 328-332.
24. Martinelli L (2013) The implementation and evaluation of a nutrition education programme for university elite athletes. *Progress in* **15**, 71-80.
25. Łagowska K, Kapczuk K, Jeszka J (2014) Nine Month Nutritional Intervention Improves Restoration of Menses in Young Female Athletes and Ballet Dancers. *Journal of the International Society of Sports Nutrition* **11**, 1-15.
26. Valliant MW, Emplaincourt HP, Wenzel RK et al. (2012) Nutrition education by a registered dietitian improves dietary intake and nutrition knowledge of a NCAA female volleyball team. *Nutrients* **4**, 506-516.
27. Molina-Lopez J, Molina JM, Chiroso LJ et al. (2013) Implementation of a nutrition education program in a handball team; consequences on nutritional status. *Nutricion hospitalaria* **28**, 1065-1076.
28. Nascimento M, Silva D, Ribeiro S et al. (2016) Effect of a Nutritional Intervention in Athlete's Body Composition, Eating Behaviour and Nutritional Knowledge: A Comparison between Adults and Adolescents. *Nutrients* **8**.
29. Garthe I, Raastad T, Refsnes PE et al. (2013) Effect of nutritional intervention on body composition and performance in elite athletes. *European Journal of Sport Science* **13**, 295-303.
30. Elias SSM, Saad HA, Taib MNM et al. (2018) Effects of sports nutrition education intervention on sports nutrition knowledge, attitude and practice, and dietary intake of Malaysian team sports athletes. *Malaysian journal of nutrition* **24**, 103-116.

31. Peitzmeier GA (2006) Impact of two nutrition interventions on dietary outcomes of female collegiate athletes. Doctor of Education, University of Kentucky.
32. Welch PK, Zager KA, Endres J et al. (1987) Nutrition Education, Body Composition, and Dietary Intake of Female College Athletes. *The Physician and sportsmedicine* **15**, 63-74.
33. Loprinzi MJ (1989) The effects of nutrition education sessions related to dietary iron on the dietary iron intake of high school wrestlers. Eugene, Ore.:: Microform Publications, College of Human Development and Performance, University of Oregon.
34. Jeukendrup AE (2017) Periodized Nutrition for Athletes. *Sports Med* **47**, 51-63.
35. Teixeira PJ, Marques MM (2017) Health Behavior Change for Obesity Management. *Obesity facts* **10**, 666-673.
36. Rumbold PL, St Clair Gibson A, Stevenson E et al. (2011) Agreement between two methods of dietary data collection in female adolescent netball players. *Appetite* **57**, 443-447.
37. Magkos F, Yannakoulia M (2003) Methodology of dietary assessment in athletes: concepts and pitfalls. *Curr Opin Clin Nutr Metab Care* **6**, 539-549.
38. Basiotis PP, Welsh SO, Cronin FJ et al. Number of days of food intake records required to estimate individual and group nutrient intakes with defined confidence. *J Nutr* **117**, 1638-1641.
39. Fogelholm M, Lahti-Koski M (1991) The validity of a food use questionnaire in assessing the nutrient intake of physically active young men. *Eur J Clin Nutr* **45**, 267-272.
40. Heaney S, O'Connor H, Gifford J et al. (2010) Comparison of Strategies for Assessing Nutritional Adequacy in Elite Female Athletes' Dietary Intake. *International Journal of Sport Nutrition & Exercise Metabolism* **20**, 245-256.
41. Chen J, Berkman W, Bardouh M et al. (2019) The use of a food logging app in the naturalistic setting fails to provide accurate measurements of nutrients and poses usability challenges. *Nutrition* **57**, 208-216.

Table 1. Participant, intervention, and dietary analysis characteristics of included studies. Table is presented as single-arm studies, followed by double-arm studies.

		Participants			Intervention			Dietary analysis	
Study	N (sex)	Age (yrs)	Population; calibre	Delivery style	Duration & frequency	Nutrition curriculum	Facilitator	Dietary analysis method	Times administered
Single-arm studies									
Collison et al., 1996	26 (F)	19.4 ± 1.2	Volleyball, field hockey, tennis; NR	Workshop	2/52 (1x weekly)	Energy; macro; micro; fluid; weight control; bone health; label reading	Dietitian	3-day diet record	Pre-intervention, Post-intervention, 3 months post-intervention (retention)
Kandiah, 2000	10 (M)	24 ± 2.12	Track & field; Elite	Weekly Newsletters, telephone counselling	16/52 (8x sessions)	Fat; cholesterol; influence of animal fat; macro & micro related to athletic performance; food sources; serving sizes; fluids	NR	3-day weighted food record	Random throughout intervention
Lagowska et al., 2014	52 (F)	17.1 ± 0.9 18.1 ± 2.6	Ballet, rowing, synchronised swimming, triathlon; High school	Individual nutrition counselling	9/12 (9x sessions)	Drinks; low-fat/calorie foods; supplements; shopping tips; food preparation; dining out; micro & deficiencies	Dietitian	7-day diet record	Post 3, 6, 9 months
Martinelli, 2013	7 F (4) M (3)	21.6 ± 2.4	Basketball, American football, shot put, water polo, rugby; Elite	Workshop	5/12 (6x sessions)	Fuelling; diet analysis; hydration; recovery; protein; supplements; self-efficacy	Sports nutritionist	3-day diet record	8x prior 8x post

Study	Participants			Intervention				Dietary analysis	
	N (sex)	Age (yrs)	Population; calibre	Delivery style	Duration & frequency	Nutrition curriculum	Facilitator	Dietary analysis method	Times administered
Molina-Lopez et al., 2013	14 (M)	22.9 ± 2.7	Handball; National	Activities	8/52 (3x 'ad hoc' sessions)	Nutrients, nutrition & physical activity, nutritional requirements, q & a	Nutrition specialist	72-h recall 3-day diet record	3x over 4/12 & weeks 0, 8, 16
Nascimento et al., 2016	ADOL M (15), F (6) Adult M (11)	15.4 23.7	Fighting, athletics, cycling, swimming, tennis, beach volleyball, surfing, rowing, sailing; State	Individual nutrition counselling; Lecture; Online Posts	4x 45-60 min, re-evaluations 60 days (6.4-8.5/52)	Hydration; meal frequency; diet quality; food guide; eating tip blog posts	Sports nutritionist	24-h recall	Pre-intervention, post-intervention
Nowacka et al., 2016	37 F (8), M (29)	21.5	Canoeing; National	Individual nutrition counselling, group consults	Over a 2-year period, intervention details NR.	Nutrition guidelines; nutritional mistakes	Nutritionist	24-h recall	3x days each season (x3) for two years
Phillipou et al., 2017	34 F (11), M (23)	15.2 ± 1.5	Swimming; National	Lecture; Shopping Tours	6/52 (2x half day sessions)	Nutrition issues; Mediterranean diet; dietary supplement use/misuse	Student dietitian	KIDMED Index	Pre-intervention, post-intervention
Rossi et al., 2017	15	19.31 ± 1.0	Baseball; Collegiate	Lecture	12/52 (initial 90-min session, then 45-min sessions every 3 weeks)	Energy; macro; micro; hydration; supplements; individual portion sizes; food timing	NR	3-day diet record	Pre-intervention, post-intervention
Valliant et al., 2012	11 (F)	19.5 ± 1.0	Volleyball; Collegiate	Individual nutrition counselling	4/12 (4x visits)	NR	Dietitian	3-day diet record	Pre-intervention & each month of intervention

Participants				Intervention				Dietary analysis	
Study	N (sex)	Age (yrs)	Population; calibre	Delivery style	Duration & frequency	Nutrition curriculum	Facilitator	Dietary analysis method	Times administered
Wenzel et al., 2012	11 (F)	19.8	Volleyball; Collegiate	Individual nutrition counselling	4/12 (3x sessions)	Selecting foods & beverages; personalised needs; practical strategies	Dietitian	3-day diet record	Pre-intervention & each month of intervention
Wittkofski, 1996	17 (F)	19 ± 1.3	Volleyball; Collegiate	Lecture, handout, activities	3x 30-min during in-season	Food groups; menu ordering; sport nutrition; SCT	Sports dietitian	24-h recall	Pre-intervention, post-intervention
Double-arm studies									
Abood et al., 2004	I: 15 (F) C: 15 (F)	19.6 ± 1.1	Soccer, Swimming; Collegiate	Lecture	8/52 (weekly, 1h)	Energy; macro; fluid; micro; nutrition principles; nutrition for travel; self-efficacy	NR	3-day diet record	Pre-intervention, 2 weeks post-intervention
Buffington et al., 2016	I: 27 (F) C: 20 (F)	22.5	Basketball, swimming, diving, volleyball, soccer, tennis, cheering, cross country, track, fencing; Collegiate	Online lecture	12/52 (weekly, 10-min sessions)	CBT in conjunction with energy balance education	NR	24-h recall	Week 1, post-intervention
Chapman et al., 1997	I: 37 (F) C: 35 (F)	16	Softball; High school	Lecture, handout	6/52 (weekly, 2x 45-min)	Ergogenic aids; dehydration; pre-competition meal; weight control	NR	24-h recall	Pre-intervention, post-intervention
Doyle-Lucas et al., 2011	I: 146 (M, F) C: 64	15.4 ± 0.2	Ballet; pre-Professional	Online Lecture	6/52 (3x 30-min)	Female athlete triad; macro; micro; hydration; healthy eating habits;	Dietitian	FFQ	Pre-intervention, post-intervention, 6 week post-

Study	Participants			Intervention			Dietary analysis		
	N (sex) (M, F)	Age (yrs)	Population; calibre	Delivery style	Duration & frequency	Nutrition curriculum	Facilitator	Dietary analysis method	Times administered
Elias et al., 2019	I: 52 (M) C: 53 (M)	Intervention: 18.69 Control: 23.26	Field hockey, football, cricket, rugby; National	Booklets, lecture, group consults	7/52 (7x 60-90min sessions)	Macro; micro; hydration; nutrition timing; energy balance; supplementation	Sports nutrition background	3-day diet record	Pre-intervention, 1 week post-intervention
Garthe et al., 2013	I: 21 (2 F) (19 M) C: 18 (1 F) (17 M)	19.1 ± 2.9	Rowing, kayak, soccer, volleyball, taekwondo, skating, ice hockey; National	Individual nutrition counselling	8-12/52, weekly (av. 9.5 weeks)	Basic nutrition; sports physiology; personalised dietary plans/weight regimen; self-efficacy	Dietitian, exercise physiologist, sports nutritionist	4-day weighed diet record (all) 24-h recall (control)	Pre-intervention, 2x mid + post
Loprinzi, 1989	I: 13 (M) C: 11 (M)	15.83 ± 2.69	Wrestling; High school	Lecture, handout	2/52 (3x 25-min sessions)	Food groups; iron deficiency, food sources; increasing iron; label reading; how iron is lost; vitamin supplements	Dietitian	3-day diet record	Pre-intervention, post-intervention
Peitzmeier, 2006 <sup>a</sup>	61 (F) I1: 17 (F) I2: 8 (F) C1: 20 (F) C2: 16 (F)	NR	Hockey & soccer; Collegiate/ National	Lecture, index cards	5/12 I1 - 1h session, questions, feedback I2 - 5x 20-min sessions, questions, feedback	Hydration; calories; macros; fast food; the timing of eating (breakfast, pre-exercise, recovery); personalised goals	Dietitian	3-day diet record	Week 1, post-intervention
Rastmanesh et al., 2007	I: 42 (NR) C: 30	30 ± 7.6	NR; Elite athletes with physical disabilities	Lecture, handout	1/12 (4x 3-h)	Food guide pyramid; nutrition & weight loss	Dietitian	3-day diet record	Pre-intervention, post-intervention

Study	Participants			Intervention			Dietary analysis		
	N (sex) (NR)	Age (yrs)	Population; calibre	Delivery style	Duration & frequency	Nutrition curriculum	Facilitator	Dietary analysis method	Times administered
Welch et al., 1987	I: 10 (F) C: 29 (F)	19.5	Basketball, cross country, field hockey, golf, swimming, softball, volleyball, tennis, track & field; Collegiate	Individual nutrition counselling	2-5x sessions	Food groups; dietary guidelines; dietary goals	Dietitian	7-d diet record 24-h recall	Pre-intervention, post-intervention

Age reported as mean  $\pm$  SD, where available. M, male; F, female; ADOL, adolescents; NR, not recorded; Macro, Macronutrients; Micro, Micronutrients; CBT, Cognitive Behavioural Theory; SCT, Social Cognitive Theory; FFQ, food frequency questionnaire; KIDMED Index (Mediterranean Diet Quality Index); I, intervention group; C, control group; <sup>a</sup>Peitzmeier (2006) assessed 2 intervention and 2 control groups.

Table 2a. Nutrient intake pre- and post-nutrition education for included studies. Table is presented as single-arm studies, followed by double-arm studies.

Study	Energy (kcal/d)		CHO (g/kg/d)		Protein (g/kg/d)		Fat (g/d)		Calcium (mg/d)		Iron (mg/d)	
	Pre	Post	Pre	Post	Pre	Post	Pre	Post	Pre	Post	Pre	Post
<b>Single-arm studies</b>												
Collison et al., 1996	1795±48 (-0.2)	1668±595	4.5	4	1.1	0.9	55.8	44.4	876	672	15.9	13.1
Kandiah, 2000	N/D	N/D	5.0±0.8	5.3±1.0 (0.0)	1.7±0.3	1.7±0.2 (0.0)	104±14	90±11* (-1.1)	988±76	1001±87 (0.2)	19.2±6.9	19.8±4.3 (0.1)
Lagowska et al., 2014 Athletes	2354±539	2800±321* (1.0)	5.2±1.3	7.0±1.0* (1.6)	1.3±0.2	1.6±0.2* (1.5)	92.2±7.5	87.4±11.1 (-0.5)	N/D	N/D	N/D	N/D
Lagowska et al., 2014 Ballet	1640±412	2368±182* (2.3)	4.2±1.1	6.3±0.6* (2.4)	1.1±0.3	1.6±0.2* (2.0)	57.0±18.9	73.2±15.9* (0.9)	N/D	N/D	N/D	N/D
Martinelli, 2013	2976±847	2652±740 (-0.4)	5.1±1.4	4.9±1.7 (0.1)	1.7±0.5	1.5±0.4 (-0.4)	99.2±38.2	77.1±23.1 (-0.7)	925±365	1113±826 (0.3)	12.8±2.8	14.5±6.2 (0.4)
Molina-Lopez et al., 2013	2975±211	3329±306* (1.3)	4.2±0.4	4.8±0.4* (1.5)	1.5±0.2	1.7±0.3* (0.8)	118.6±22.5	129.6±21.8 (0.5)	1251±338	1235±393 (0.0)	24.2±8.5	24.4±6.1 (0.0)
Nowacka et al., 2016 Female	1654±426	1807±373* (0.4)	3.8±1.0	4.0±0.7 (0.2)	1.1±0.2	1.2±0.4 (0.3)	52.0±16.5	58.4±16.3 (0.4)	N/D	N/D	N/D	N/D
Nowacka et al., 2016 Male	2372±379	2762±434* (1.0)	4.4±0.9	5.1±1.0* (0.7)	1.4±0.2	1.5±0.2 (0.5)	78±19.5	91.0±16.9 (0.7)	N/D	N/D	N/D	N/D
Rossi et al., 2017	2878±443	3366±451* (1.1)	3.6±0.9	3.8±0.7 (0.2)	1.8±0.3	2.1±0.4* (0.8)	129±21	162±37* (1.1)	N/D	N/D	N/D	N/D
Valliant et al., 2012	1756±558	2178±492* (0.8)	3.1±1.1	4.2±1.3* (0.9)	0.9±0.3	1.1±0.3* (0.7)	67.4±27.8	69.0±24.8 (0.1)	N/D	N/D	N/D	N/D
Wenzel et al., 2012	56.0% <sup>§</sup>	70.0% <sup>§</sup>	3.1	4.1	1.0	1.1	N/D	N/D	N/D	N/D	N/D	N/D
Wittkofski, 1996	2524±655	2256±516 (-0.5)	5.4±0.7	4.9±0.9	1.3±0.4	1.0±0.4 (-0.8)	64.5±16.8	65.2±25.1 (0.0)	N/D	N/D	N/D	N/D

Study	Energy (kcal/d)		CHO (g/kg/d)		Protein (g/kg/d)		Fat (g/d)		Calcium (mg/d)		Iron (mg/d)	
	Pre	Post	Pre	Post	Pre	Post	Pre	Post	Pre	Post	Pre	Post
<b>Double-arm studies</b>												
Abood et al., 2014	1969±414	1974±473	4.2±0.7	4.8±0.8	1.0±0.2	1.2±0.2	52.5±15.3	50.3±13.6	659±229	309±344	18±12	21±16
Intervention		(0.0)		(0.8)		(1.0)		(-0.2)		(0.5)		(0.2)
Control	2240±366	1947±515	5.5±0.6	4.4±0.7*	1.2±0.6	1.2±0.5	57.2±14.9	45.4±8.7	918±305	695±329	15±4	13±5
Buffington et al., 2016		(-0.7;0.1)		(-1.7;0.5)		(0.0;0.0)		(-1.0;0.4)		(-0.7; 0.3)		(-0.4;0.7)
Intervention (%E)	N/D	N/D	46.6±15.9	57.0±11.4*	17.2±7.2	16.5±4.9	33.6±14.0	25.4±8.8	N/D	N/D	N/D	N/D
Control (%E)			47.5±12.6	51.0±12.7	15.6±5.0	19.7±4.8*	34.9±8.6	29.1±10.6				
				(0.8;0.5)		(-0.1;- 0.7)		(-0.7;-0.4)				
								(-0.6)				
Chapman et al., 1997	2054	1892	48.5	51.0	13.0	12.0	39.5	34.0	N/D	N/D	N/D	N/D
Intervention (%E)	1683	1793	47.0	48.2	9.0	9.5	39.0	41.0				
Control (%E)												
Doyle-Lucas et al., 2011 <sup>c</sup>												
Intervention	N/D	N/D	N/D	N/D	N/D	N/D	95.2±1.8	88.2±1.6*	N/D	N/D	N/D	N/D
Control							91.7±2.8	89.3±2.5				
Elias et al., 2019	2478±364	2879±385* <sup>†</sup>	5.3±1.1	5.8±1.2* <sup>†</sup>	1.7±0.4	2.1±0.5* <sup>†</sup>	75.9±16.2	94.3±19.8* <sup>†</sup>	N/D	N/D	N/D	N/D
Intervention		(1.1; 0.4)		(0.4;1.8)		(0.9;0.7)		(1.0;-0.1)				
Control	2801±541	2697±600*	4.4±0.9	4.1±0.6*	1.8±0.4	1.8±0.4	101.7±26.4	96.2±26.8				
Garthe et al., 2013		(-0.2)		(-0.4)		(0.0)		(-0.2)				
Intervention	3041±578	3585±600* <sup>†</sup>	5.4±1.1	6.8±1.3* <sup>†</sup>	1.8±0.4	2.4±0.4* <sup>†</sup>	99.0±28.3	99.0±21.2 <sup>†</sup>	N/D	N/D	N/D	N/D
Control	3032±771	2964±884	5.4±1.7	4.5±1.9	1.7±0.5	1.7±0.4	104.9±44.8	112.4±30.0				
Loprinzi, 1989		(-0.1)		(-0.5)		(0.0)		(0.2)				
	N/D	N/D	N/D	N/D	N/D	N/D	N/D	N/D	N/D	N/D	7.6±2.9	11.3±2.5*

Study	Energy (kcal/d)		CHO (g/kg/d)		Protein (g/kg/d)		Fat (g/d)		Calcium (mg/d)		Iron (mg/d)	
	Pre	Post	Pre	Post	Pre	Post	Pre	Post	Pre	Post	Pre	Post
Intervention												(1.4;0.3)
Control												10.8±3.8*
Peitzmeier, 2006	2034	2103	4.5	4.6	0.8	1.4	67.5	66.3	797	879	15.9	15.3
Intervention 1	2139	2323	4.9	5.2	1.3	1.4	62.7	71.4	1041	974	17.0	16.6
Control 1	1626	1647	3.4	3.5	1.0	1.1	51.3	48.3	642	688	11.5	13.4
Intervention 2	1409	1356	3.1	3.2	1.1	0.9	34.7	32.3	646	531	11.7	12.9
Control 2												
Rastmanesh et al., 2017												
Intervention	1690±490	1520±450* (-0.4)	N/D	N/D	N/D	N/D	N/D	N/D	715±250	840±300* (0.5)	N/D	N/D
Control <sup>d</sup>	1670±430	-							690±270			
Welch et al., 1987	1722±1052	1771±852 (0.1;-0.2)	3.5±0.6	4.3±0.8* (1.1;0.8)	0.7±0.1	0.9±0.3 (0.9;-0.3)	80.4±15.3	57.1±17.7* (-1.4;-0.9)	610±214	822±323 (0.8;0.2)	6.0±4.1	15.4±13.3 (1.0;0.4)
Intervention	1788±654	1893±648 (0.2)	3.6±0.7	3.7±0.8 (0.1)	1.1±0.4	1.0±0.3 (-0.3)	69.5±17.9	73.6±18.9 (0.2)	688±447	731±384 (0.1)	8.9±3.2	12.4±4.9* (0.8)
Control												

Results presented as mean ± SD (Within-group ES; Between-group ES), where available. \*, significant within-group (pre-post) difference ( $p < 0.05$ ); †, significant between-group (post-intervention) difference ( $p < 0.05$ ); <sup>c</sup>, presented as mean ± standard error of the mean; <sup>d</sup>, no post-intervention assessment conducted.

Table 2b. Diet quality pre- and post-nutrition education for included single-arm studies, followed by double-arm studies.

	Vegetables (portions/d) <sup>a</sup>		Fruit (portions/d) <sup>a</sup>		Grains (portions/d) <sup>a</sup>		Dairy (portions/d) <sup>a</sup>		Meat, poultry, eggs (portions/d) <sup>a</sup>	
	Pre	Post	Pre	Post	Pre	Post	Pre	Post	Pre	Post
<b>Single-arm studies</b>										
Molina-Lopez et al., 2013	2.2 (1.7-2.8)	2.1 (1.6-2.6)	1.7 (0.7-2.7)	1.1 (0.7-1.6)	3.6 (2.8-4.4)	3.4 (2.9-3.9)	2.7 (1.8-3.6)	2.7 (2.2-3.3)	4.5 (6.3-8.6)	6.8 (5.8-7.8)
Nascimento et al., 2016 <sup>a,b</sup> (Adequate)	6.3 (2.5-16.0)	2.5 (1.4-4.8)*	6.6 (5.0-8.7)	4.8 (2.6-8.6)*	9.8 (6.7-14.0)	6.1 (3.5-10.0)*	5.0 (3.8-6.8)	3.3 (1.4-8.0)	2.1 (1.6-3.0)	2.8 (1.8-4.0)
Nascimento et al., 2016 <sup>a,b</sup> (Low)	1.6 (1.6-3.1)	2.2 (0.8-7.8)*	2.4 (1.2-5.0)	4.6 (1.6-12.0)*	3.0 (1.7-5.5)	3.8 (2.1-6.8)	1.8 (1.1-2.9)	2.5 (1.7-3.7)*	4.0 (3.0-5.0)	2.8 (1.7-4.0)*
Wittkofski, 1996	3.1±2.3	2.8±2.8 (-0.1)	2.5±2.7	1.4±1.5 (-0.5)	9.6±6.0	7.6±2.6 (-0.4)	4.1±1.8	2.7±1.5 (-0.8)	1.3±1.1	2.5±1.5 (0.9)
<b>Adherence to Mediterranean Diet (KIDMED Score)<sup>a</sup></b>										
	Pre	Post								
Phillipou et al., 2017 <sup>c</sup>	5.7±2.1	7.6±1.7 (1.0)								
<b>Double-arm studies</b>										
	Fruit & Vegetables (portions/d) <sup>d</sup>		Grains (portions/d) <sup>d</sup>		Dairy (portions/d) <sup>d</sup>		Meat, poultry, eggs (portions/d) <sup>d</sup>			
	Pre	Post	Pre	Post	Pre	Post	Pre	Post		
Doyle-Lucas et al., 2011			N/D	N/D			N/D	N/D		
Intervention	4.9±0.1	4.5±0.1*			2.5±0.1	2.7±0.1				
Control	4.9±0.2	4.4±0.2			2.5±0.2	2.4±0.2				

Results presented as mean ± SD (Within-group ES; Between-group ES), where available. N/D, not defined; \*, significant within-group (pre-post) difference ( $p < 0.05$ ); <sup>a</sup>, presented as mean (95% CI); <sup>b</sup>, Nascimento et al 2016 classified participants as adequate or low based on meeting recommended food portion number (or adequate and high for meat and eggs); <sup>c</sup>, Adherence to the MD is described as poor (score: 0–3), medium (score: 4–7) or good (score: 8–12); <sup>d</sup>, presented as mean ± standard error of the mean.

Table 3. Study design, methodology and reporting summary checklist of included studies (n=22), Dietary Intake and Nutrition Education Reporting for Sports (DINERS) checklist.

Item	Yes	No	Unable to determine/not applicable
<b>Participant characteristics</b>			
Age	17	5	0
Sex	21	1	0
Athletic calibre	22	0	0
Sport type	21	1	0
Representative sample recruited	6	15	1
Training characteristics (type, frequency, intensity, duration)	2	20	0
Body composition assessed at baseline	10	12	0
Body composition assessed at post-intervention	10	12	0
Body composition assessment protocols clearly defined	4	5	0
Weight change assessed pre- and post-diet	13	9	0
<b>Targeted dietary outcomes</b>			
Energy intake targets defined and justified	3	17	2
Macronutrient targets defined and justified	2	18	2
Micronutrient targets defined and justified	1	16	5
Fluid or other dietary outcomes justified and defined	0	15	7
Dietary outcomes linked to athlete goals	3	18	1
<b>Intervention characteristics</b>			
Curriculum covered is clearly described	17	5	0
Instructor background/qualifications described	12	10	0
Modality (electronic, face to face, individual, group) described	19	2	0
Session characteristics (individual session duration, number, and frequency) described	8	14	0
Duration of the intervention	18	4	0
Compliance/attendance	11	10	1
Behavioural Theory clearly defined and justified	5	16	1
Previous dietary education outlined	6	16	0
Process evaluation by participants	2	20	0
Follow-up or retention outlined if applicable	3	10	9

**Dietary methodology**

Method used to assess nutrients is appropriate (e.g. food diary, FFQ)	10	12	0
Dietary collection methods well described	20	2	0
Participant <i>n</i> adequate for dietary methods used	20	2	0
Number of collection days and description clearly defined	22	0	0
Number of days collected for primary nutrient justified	12	8	0
Participant instruction clearly described	9	11	2
Checking and cleaning of diet data well described	11	8	2
Dietary analysis software well described and appropriate	17	3	2
Background/qualifications of researcher conducting diet analysis is appropriate	5	14	3
Energy expenditure empirically measured or prediction provided	7	15	0
PAL level/under-reporting assessed	0	22	0
Revised nutrient requirements addressed - iron altitude, amenorrhoea	1	21	0
Energy reported as kJ/kg	3	17	2
Carbohydrate expressed as g/kg/d	6	14	2
Protein expressed as g/kg/d	7	13	2
Fat (% of energy reported)	13	7	2
Fat types reported when relevant	2	7	13
EAR/RDA and relevant sports guidelines used for micronutrients	5	13	4
Proportion of participants meeting or under recommendations reported	3	18	1
Contribution of diet supplements/sports foods to intake well described	2	20	0

---

PAL, physical activity level.

Figure 1. PRISMA flowchart.

