

Bond University
Research Repository



Optimising nutrition in residential aged care: A narrative review

Agarwal, Ekta; Marshall, Skye; Miller, Michelle; Isenring, Elisabeth

Published in:
Maturitas

DOI:
[10.1016/j.maturitas.2016.06.013](https://doi.org/10.1016/j.maturitas.2016.06.013)

Licence:
CC BY-NC-ND

[Link to output in Bond University research repository.](#)

Recommended citation(APA):
Agarwal, E., Marshall, S., Miller, M., & Isenring, E. (2016). Optimising nutrition in residential aged care: A narrative review. *Maturitas*, 92, 70-78. <https://doi.org/10.1016/j.maturitas.2016.06.013>

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

For more information, or if you believe that this document breaches copyright, please contact the Bond University research repository coordinator.

Title: Optimising nutrition in residential aged care: a narrative review

Authors:

Agarwal Ekta¹, Marshall Skye², Miller Michelle³, Isenring Elisabeth²

¹School of Exercise and Nutrition Sciences, Queensland University of Technology, Brisbane, Australia. Email: ekta.agarwal@hotmail.com

²Faculty of Health Sciences and Medicine, Bond University, Robina, Australia.

Email: skye_marshall@bond.edu.au; lisenrin@bond.edu.au

³School of Health Sciences, Flinders University, Adelaide, Australia.

Email: michelle.miller@flinders.edu.au

Corresponding Author:

Professor Elisabeth Isenring

Faculty of Health Sciences and Medicine

Level 2, Building 18, Bond Institute of Health and Sport

Bond University

Qld 4229, Australia

Tel: +61755953337; Email: lisenrin@bond.edu.au

Abstract

In developed countries the prevalence of protein-energy malnutrition increases with age and multi-morbidities increasing nutritional risk in aged care residents in particular. The purpose of this paper was to examine, synthesise and develop a narrative review of the current literature on the identification, prevalence, associated risk factors, consequences, and management of malnutrition in the residential aged care (RAC) setting. We performed searches of English language publications in Medline, PubMed, Ovid and the Cochrane Library from January 1 1990- November 25 2015. We found that, on average, half of all residents in aged care are malnourished as a result of factors affecting appetite, dietary intake and nutrient absorption. Malnutrition is associated with a multitude of adverse outcomes including increased risk of infections, falls, pressure ulcers and hospital admissions, all of which can lead to increased health care costs and poorer quality of life. A number of food and nutrition strategies have demonstrated positive nutritional and clinical outcomes in the RAC setting. These strategies extend beyond simply enhancing the nutritional value of foods and hence necessitate the involvement of a range of committed stakeholders. Implementing a nutritional protocol in RAC facilities that comprises of routine nutrition screening, assessment, appropriate nutrition intervention, including attention to food service systems, and monitoring by a multidisciplinary team can help prevent decline in residents' nutritional status. Food and nutritional issues should be identified early and managed on admission and regularly in the RAC setting.

Keywords: Malnutrition, Aged, Nutrition Assessment, Nutritional Support

1. Introduction

The proportion of the world's population aged 60 years and over is rapidly increasing (1). Whilst this population has doubled from 12% in 1950 to 23% in 2013, it is projected to triple to 32% by 2050 (1, 2). In developed countries improvements in healthcare have led to declines in mortality from conditions such as stroke and heart disease (3). However, increased longevity has resulted in an increased prevalence of chronic illness and multi-morbidities (defined as the presence of two or more diseases at the same time); along with a burgeoning demand for residential aged care (RAC) services to offer accommodation and support for older people with complex needs who can no longer live independently in their home (4, 5).

Malnutrition (or inadequate nutritional status) in older people is defined as undernourishment due to poor dietary intake consequently leading to involuntary weight loss and muscle wasting (6). Although frail older persons are at an increased risk of malnutrition, research indicates that nutrition often ranks low on the list of care priorities in RAC (7). Lack of knowledge and awareness, and the inability to identify malnutrition by care staff; rushed and task-focussed eating environments that are often accompanied by little communication and conversation between residents and staff assisting with feeding; and residents' own physical, clinical, and psychosocial issues have been identified as major barriers against the prevention and management of malnutrition in the elderly (7-12). In a recent poll conducted by the International Association of Gerontology and Geriatrics, RAC experts from eight countries identified "improving nutrition" as one of the top five international research priorities in this setting (13). To the best of our knowledge, previously published reviews in the RAC setting have summarised evidence regarding malnutrition prevalence (14), associated risk factors (15), and effective nutritional interventions (16-18). The purpose of this paper is to review and summarise existing evidence on these aspects of malnutrition in the RAC setting, along with including a synopsis of evidence-based methods to identify malnutrition in the RAC setting, highlight the consequences of malnutrition in this population and provide future directions for practice and research.

2. Methods

In the literature, depending on the geographical location, low care settings may also be referred to as “hostels”, “residential care homes”, or “assisted care”; whereas high care settings may also be referred to as “nursing homes”. For the purpose of this paper, the term RAC will encompass both low and high care settings. References for this narrative review were identified through searches of publications listed in electronic databases including Medline, PubMed, Ovid, and the Cochrane Library. The searches were limited to full-length articles, in the English language, and publications dated from 1 January 1990 to 25 November 2015. The titles and abstracts of articles were examined using the following keywords: “nursing homes”, “residential aged care facility”, “elderly”, “older adults”, “malnutrition”, “nutritional status”, “multicentre”, “prevalence”, “aetiology”, “consequences”, “health-related outcomes”, “nutritional interventions”, “dietary intake”, and “enteral nutrition”. Retrospective searches were also conducted from relevant review articles, book chapters and relevant evidence-based guidelines. Further details of the search methods are available from the corresponding author.

3. Identification of malnutrition in RAC

Although there is no gold standard for diagnosing malnutrition, it is generally accepted that the condition is characterised by unintentional weight loss with suboptimal dietary intake, thereby resulting in muscle wasting and/or loss of subcutaneous fat; or low BMI ($\leq 18.5\text{kg/m}^2$) (19). However, in a recent meta-analysis evaluating the association of BMI and all-cause mortality in older adults, Winter and colleagues observed that when compared with a BMI reference range of 23.0-23.9 kg/m^2 , a BMI of 21.0-21.9 kg/m^2 and a BMI of $\leq 20.0\text{kg/m}^2$ demonstrated a 12% and 28% greater risk of all-cause mortality respectively (20). This meta-analysis indicates that the healthy BMI range for adults (18.5-24.9 kg/m^2) as established by the World Health Organisation is not suitable for older adults and that individuals with a BMI of $<23.0\text{kg/m}^2$ should be monitored for changes in weight and risk of malnutrition (20). Research also indicates that a moderately higher BMI is protective against morbidity and mortality (21, 22).

It is important to remain mindful that just like some laboratory parameters (albumin, transferrin, and lymphocyte count) are more indicative of illness rather than nutritional deficits, body weight is also affected by health status rather than nutrition exclusively (23). Therefore, rather than relying on a single parameter to identify malnutrition, it is believed that nutrition screening and assessment tools are more suitable to identify malnutrition risk and malnutrition as they take into account its complex multifactorial aetiology including biological, iatrogenic, socioeconomic, psychological and physical risk factors (24). Of the several nutrition screening tools available in the literature, many have been developed for older adults (25-31) and some have been specifically developed for use in the RAC setting (32-35). However, recent systematic reviews concluded that few tools have undergone rigorous reliability and validity testing; the ability of nutrition screening tools to screen nutritional status and predict outcomes in RAC residents ranges from fair-to-moderate; and that more items that are specific to development of malnutrition in this population (dietary factors, feeding dependence, oral health, environment, comorbidities) need to be included in the tools (24, 36, 37). White et al have proposed that since no single parameter can define malnutrition, the diagnosis of malnutrition should be made with the identification of two or more of the following characteristics rather than depend on any particular nutrition screening tool (38):

- Inadequate energy intake: defined as consuming $\leq 75\%$ of estimated energy requirements for at least one month;
- Weight loss: defined as unintended loss of at least 5% usual body weight within one month and/or at least 10% loss in usual body weight within six months;
- Loss of muscle mass: defined as mild-to-severe visible wasting of the temples, clavicles, shoulders, interosseous muscles, scapula, thigh and calf;
- Loss of subcutaneous fat: defined as mild-to-severe visible wasting orbital, triceps, and fat overlying the ribs;
- Localised or generalised fluid accumulation: in the extremities, vulvar/scrotal oedema, and ascites;

- Reduced functional capacity: determined by handgrip strength.

However, this proposed consensus definition for malnutrition is still to be tested for validity and is a work in progress. Comprehensive assessment of nutritional status by trained professionals, such as dietitians, is essential once malnutrition risk has been identified. Nutrition assessment uses a combination of medical history, nutritional data, physical examination, anthropometric measurements, and biochemical parameters to determine the individual's nutritional status (38). The Subjective Global Assessment (SGA) (39), scored Patient-Generated Subjective Global Assessment (PG-SGA) (40), and Mini-Nutritional Assessment (41) are multidimensional nutrition assessment tools validated for the RAC setting (42-45) and include a range of parameters to make a nutritional diagnosis and initiate nutritional management in older residents.

4. Prevalence of malnutrition in RAC

Bell et al recently published a systematic review on malnutrition prevalence in the RAC setting (14). The review included studies that were published from 1999-2011, single and multicentre studies, and cohorts ranging from 50–11,902 RAC residents (14). Various methods such as nutrition screening and assessment tools, body mass index (BMI), weight loss percentage, anthropometric measurements, and weight were utilised to determine malnutrition prevalence (14). Malnutrition risk and prevalence were estimated to range from 29-97% and 4-71% respectively, with the large variation attributed to selection bias and research methodology to define malnutrition (14). However, in most studies malnutrition risk and prevalence were reportedly between 47-62% and 20-39% respectively (14). The findings of this review are consistent with results from other multicentre studies since published from regions including Iran, Italy, Austria and Finland (46-50).

5. Aetiology of malnutrition in RAC

The aetiology of malnutrition in older adults is complex and multifactorial. Inadequate dietary intake is the leading cause of malnutrition in older adults (51). A physiological reduction in appetite and energy intake, referred to as the “anorexia of ageing”, occurs naturally with progressive aging and usually exceeds the reduction in energy expenditure, resulting in unintentional weight loss mainly in the form of lean body mass (52). Additionally, a multitude of factors often referred to as the “nine D’s” (Figure 1) are commonly observed in older adults and are associated with difficulty with self-feeding; social and financial problems; impaired digestion, absorption, and excretion of nutrients; and gastrointestinal symptoms (such as nausea, constipation, vomiting, diarrhoea, loss of appetite, and early satiety); resulting in decreased food consumption and exacerbation of malnutrition (12, 15, 53-61). The use of plastic dishware has been associated with malnutrition due to its potential influence on taste of food items and residents’ perception of being in hospital (62). Although not well explored in the literature, food insecurity (or the limited ability or uncertainty in accessing adequate, safe, and nutritious food that meets one’s taste and cultural preferences in a socially acceptable manner) has also been implicated in the aetiology of malnutrition in the RAC setting (63, 64). RAC residents are particularly vulnerable to food insecurity due to their increasing age, presence of debilitating diseases, lack of control over food intake and preferences, declining physical and psychological health, and limited social and financial capacity (62, 64). Evidence also indicates that the inability of nursing staff to identify and recognise malnutrition as a formidable problem in the RAC setting results in them not prioritising nutrition care for the residents thereby adding to risk of RAC residents developing malnutrition (7, 10, 65).

6. Consequences of malnutrition in RAC

Malnutrition bears significant and diverse consequences including physiological, psychosocial and healthcare costs-related consequences.

6.1. *Physiological consequences:* The breakdown of liver protein during catabolism in malnourished older adults inhibits hepatic protein synthesis leading to a decrease in immune cells and serum proteins (66). This can cause anaemia, oedema, poor wound healing and impaired immunity with the consequence of increased susceptibility to infections and pressure ulcers. The loss of cardiac muscle decreases cardiac output thereby compromising heart function (67). The ability to control breathing becomes impaired due to the loss of respiratory muscle mass and strength as well as decreased production of surfactant (68). Malnutrition further affects lung structure, elasticity and immune defence and pulmonary oedema can result due to a decrease in colloidal osmolarity secondary to hypoproteinaemia (68). The combination of sarcopenia, loss of fat, impaired cardiac and respiratory function and impaired thermoregulation contribute to a significantly increased risk of falls (69). Malnutrition-related decreasing physical function in multiple organ systems is also associated with an increased risk of mortality (70). Malnutrition was found to be independently associated with a 3- to 6-fold increased risk of mortality in RAC (71-73). Furthermore, significant weight loss within six months and/or eating problems in RAC residents has also been found to increase the risk of 3 month mortality by 20% (74).

6.2. *Psychosocial consequences:* Depression is prevalent in older adults, with rates of up to 45% in the institutionalised elderly (75, 76). Depression is associated with weight loss and although the association between depression and nutritional status is multifactorial it is unclear if it is a cause and/or consequence of malnutrition in older residents (77-81). As a result of these physiological and psychosocial outcomes, malnutrition in the individual leads to decreased physical functioning and quality of life (82).

6.3. *Healthcare costs-related consequences:* Beyond the burden of malnutrition to the resident, malnutrition also contributes significantly towards the economic burden in RAC facilities. While

the additional cost of malnutrition in the RAC setting has been estimated in excess of €450 million (USD\$489 million) (83), the cost of treating malnourished patients in RAC in the UK has been estimated at £2.6 billion (USD\$3.9 billion) (84).

Based on the foregoing it can be inferred that early identification and management of malnutrition is imperative not only to manage the associated physiological and psychosocial consequences for residents but also to minimise associated health-related costs.

7. Nutritional management of malnutrition in RAC

Malnutrition is preventable and treatable, and in some instances completely reversible. The goals of managing the nutritional status of long term RAC residents include not only the treatment of malnutrition but also the prevention of decline in nutritional status (85, 86). Although mealtimes are recognized as ‘the highlight of the day’ for many RAC residents (87) admission to RAC is known to come with challenges for residents including adjusting to specific meal times and choices, and dining companions and environments; and depending on RAC menus solely for their nutritional intake (88). It has been suggested that malnutrition may be caused iatrogenically by dietary accreditation standards that are either inadequate at meeting nutritional requirements of elderly residents (89) or extremely limited in offered choices related to menu selection (90, 91). Therefore, to alleviate the issue of malnutrition in the RAC setting, there is increasing recognition for (a) revising existing menu guidelines to address nutritional requirements of elderly residents likely to have limited food intake (89); (b) accommodating residents’ preferences based on cultural diversity and special dietary requirements (92); (c) improving mealtime choices and variety offered to residents (88). Furthermore, staff education, regular monitoring for nutritional risk and weight, with prompt and individualised triage for malnourished residents is crucial in the management of malnutrition (93-95). In addition, the training and appointment of a staff member as a nutrition coordinator or champion has also demonstrated positive results in managing malnutrition in RAC (96). Evidence highlighting the use of various strategies to improve residents’ nutritional status is as below:

7.1 Evidence for the use of modifications to food and mealtimes in the RAC setting:

Since inadequate food intake is the most prominent risk factor contributing to malnutrition in the elderly it is logical to implement strategies that improve food intake in the treatment of malnutrition.

(i) Meal enrichment: The provision of energy- and protein-enriched meals and snacks is a low-cost, ‘food first’ approach to improving nutrient intake without increasing the amount of food offered,

and may prevent decline in nutritional and functional status for at risk populations (97-99). Therefore, the provision of a high energy-high protein diet, such as through the addition of milk powder, egg yolk, nuts, oil and cream, as well as dairy-based mid-meals, should be provided to older RAC residents who are found to be at risk of malnutrition (98, 100-103). However, research also suggests that food fortification is unlikely to provide sufficient nutritional support to treat residents with overt malnutrition or significantly improve physical function (100-102, 104). This may be due to the poor intervention adherence of residents and staff, short intervention periods, insufficient nutrition support, and insufficient physical activity in a largely frail and immobile population (100-102, 105, 106).

(ii) Therapeutic diets: An Australian study comprising of 199 residents from 18 RAC facilities found that whilst residents' nutritional intake was inadequate in meeting their energy and protein requirements, their sodium intake was three times higher than the recommendation (106). Sugar intake was also high and mainly contributed by excessive serves of 'extras' (including cakes, biscuits, confectionery, soft drinks, and ice-cream) (106). Research demonstrates that prescriptive therapeutic diets, often used in aged care for conditions such as diabetes and cardiovascular diseases, may increase the risk of malnutrition due to their restrictive nature (12, 107). To prevent excessive intake of sodium, the use of herbs and spices as flavour enhancers is recommended to improve flavour perception that may be affected by age-related sensory deficits. Regular physical activity is also a beneficial alternative to food restriction due to its multifactorial effect including increased insulin sensitivity, improved cardiovascular profile, improved muscle mass, and enhanced quality of life.

(iii) Texture modified diets: It is estimated that over half the RAC population is affected by degenerative conditions such as dementia, stroke, Parkinson's disease and Alzheimer's disease that commonly results in dysphagia, or a dysfunctional swallow (108). Texture modified diets are therefore often prescribed to residents with dysphagia and/or poor dentition (108, 109). However, texture modified diets are known to be nutritionally dilute due to the addition of fluid to achieve

desired consistency, are limited in choice, and visually unappetizing thereby contributing towards reduced intake and thus malnutrition (88, 109-111). To manage these issues, Germain et al used a technology that reformed the appearance of smooth homogenous pureed food to look like its natural form, without altering the nutritional content of the food (108). When malnourished residents with dysphagia were offered this food over a 12-week period a significant improvement was observed in their energy intake (+ 600kcal) and weight (+ nearly 4kg) from baseline (108). This study demonstrated that with careful planning and execution, texture modified foods can have an improved appearance, which can result in better nutritional intake by residents in RAC facilities.

(iv) Mealtime ambience: In the RAC setting, meal ambience, delivery method, choice over the meals and overall diet liberalisation may promote appetite and thereby have a significant impact upon the total dietary intake of residents (14, 90, 112). A Dutch cluster randomised, controlled trial revealed that when meals were provided “family style”, energy intake improved by 991kJ (95%CI: 504-1479kJ), which was both statistically and clinically significant, leading to an improvement in global nutritional status (112). Other successful strategies to promote mealtime ambience include restaurant style seating and involving the residents in cooking the meals on the ward (51).

7.2 Evidence for the use of oral nutritional supplements in the RAC setting

The provision of ONS has shown to significantly improve energy and protein intake beyond that provided by a regular or high energy-high protein diet (113, 114). Although ONS consumption may decrease the total amount of energy and protein consumed by meals and snacks, the supplements still result in an overall increased energy and protein intake in malnourished RAC residents (114, 115).

The provision of ONS in the RAC setting has shown to be effective in promoting weight gain and mid-arm and calf circumference and health-related quality of life in residents at risk or malnourished, and therefore may improve nutrition status in malnourished groups (17, 116-119). Only one study reported an improvement in global nutrition status with the provision of ONS (119); however, it must be acknowledged that very few studies have included global nutrition status as an

outcome (17). Although some ONS interventions have shown improvements in physical function (17, 119, 120), others have shown no improvement in physical function unless ONS was combined with resistance training (116, 121, 122). Physical activity combined with ONS and oral care was also shown to have additional beneficial effects on social engagement in one study (122).

Resident adherence in consuming ONS is positively associated with improvements in nutritional parameters, such as weight and mid-arm circumference (118, 123). Adherence has been found to improve when ONS is provided as a nutrient dense, low volume option, of approximately 50 - 125ml with accompanying dietary counselling (116, 118, 123). Adherence is also shown to be higher in residents who are malnourished or have chewing difficulties, but lower in residents who are depressed, are immobile or report gastrointestinal distress, indicating residents with these characteristics will require increased individualised support (118).

Residents with dementia present a unique challenge in the treatment of malnutrition due to increased difficulty with ONS adherence. A 2010 systematic review found that residents with dementia had increased levels of adherence when they received the ONS as part of medication rounds, between meals, and when they received encouragement by carers (124). However, carer compliance with providing the prescribed ONS was still found to be a significant problem (124). Although there is limited research into the health-related outcomes of ONS prescription in RAC residents with dementia, some studies have shown an improvement in weight status and nutrition status (124), which aligns with findings in the general RAC population.

Overall, evidence suggests that individualised ONS in low volumes (50-125ml) may be effective in treating malnutrition in residents by promoting weight gain and resident adherence (116, 118). Oral nutrition supplements may prevent a decrease in physical function, but should be used in combination with resistance exercise in frail residents (17, 115, 119-121). Measures to increase adherence will significantly improve the resident outcomes as well as decrease waste, especially in residents with dementia (118, 120, 124). These findings further emphasise the need for policy

which promotes regular malnutrition screening and nutrition assessment of residents, so that intervention can be needs-based.

7.3 Evidence for the use of enteral and parenteral nutrition in the RAC setting

Few studies have investigated the efficacy of enteral nutrition in the RAC setting for the purpose of preventing or treating malnutrition, despite a large number of residents having a feeding tube (125). One study found that dysphagic residents (predominately with dementia) who were fed by percutaneous endoscopic gastrostomy (PEG) were more likely to have a low BMI but failed to have anthropometric improvement with PEG feeding despite increased energy and protein intake compared with orally-fed dysphagic residents (126). Although biochemical nutritional parameters improved, the findings suggest the need for enteral nutrition to be combined with physical activity in order to improve body composition (126). Conversely, an earlier study showed that residents with dementia who received enteral nutrition had a significant improvement in weight; however, were also more likely to have aspiration pneumonia (127). No intervention studies were identified which examined the role of parental nutrition for the prevention or treatment of malnutrition in the RAC setting.

It can be argued that all older adults admitted to RAC facilities are at risk of malnutrition in the long term, and there may be a cost-benefit to the RAC facilities to provide a liberalised, prophylactic, high energy-high protein diet as the standard menu, as this may prevent the significant costs associated with malnutrition and/or frailty (107, 128, 129). Although little research has been done into the cost-effectiveness of nutritional interventions in RAC, a Dutch study showed that the cost of managing a resident at risk of malnutrition was an additional €8,200 (USD\$8,800), and an additional €10,500 (USD\$11,300) for a malnourished resident per year (130). Conversely, the provision of a high energy-high protein diet via food fortification was found to cost €40 (USD\$43) per resident per year in Sweden (104).

8. Implications for practice and future research

A collaborative multidisciplinary approach to developing effective and systematic strategies that help with identifying, diagnosing and treating malnutrition is imperative to the management of malnutrition in the RAC setting (131). Adequate staffing along with provision of nutrition education, support and adequate time allowance for offering daily care must be given due consideration by policy makers and government authorities (9).

It is imperative to have a consistent approach when defining malnutrition in the RAC setting as this will prevent the issue of potential misdiagnosis and facilitate improved correlation between effective nutritional interventions and health-related outcomes thereby influencing advocacy for the development and implementation of suitable policies, regulations and legislations (38). Residents found to be at risk of malnutrition should undergo comprehensive nutrition assessment by a dietitian or physician. A dietitian is ideally placed to provide nutrition assessment and individualised nutrition intervention due to not only their expertise in malnutrition, but also because many physicians in the RAC setting have insufficient time to dedicate solely to developing and implementing a nutrition care plan. However, in an RAC setting, the nutrition care plan involving a multidisciplinary team is critical for resident improvement, and should involve the dietitian, physician, carers and physical therapists.

There is evidence to suggest that in RAC residents, although the volume of intake at main meals remains unchanged regardless of menu type, energy intake is 50% higher with enriched meals Odlund Olin (104). To increase the energy density of meals, fat is usually added for its higher per gram energy content (37kJ/g) in comparison to protein (17kJ/g) and carbohydrate (16kJ/g). While fat has a lower degree for satiety per calorie, it has the ability to improve food palatability and acceptance, reinforcing the idea that low-volume, energy dense foods are a valid strategy to increase energy intake without affecting appetite in older adults. The provision of a high energy and high protein liberalised diet as the standard menu is likely to improve the nutrition status and quality of life of the resident. However, these enriched meals should be provided in a manner which increases

meal ambience, meal quality and resident choice in order to increase dietary intake and reduce waste, thereby also contributing to the prevention and possibly treatment of malnutrition. Oral nutritional supplements will help to improve the weight status and health-related quality of life in some malnourished residents, but should be combined with resistance exercise in frail older adults to improve physical function. Furthermore, strategies should be employed to increase staff and resident adherence with ONS, which will improve resident outcomes as well as cost-effectiveness. The evidence for enteral nutrition as a method of malnutrition treatment in RAC residents is controversial and insufficient, and no studies have examined the role of parental nutrition.

Since malnutrition is associated with inadequate nutritional intake, and given its high prevalence in the RAC setting, it can be concurred that food availability in RAC facilities is not synonymous with food security for RAC residents (64). However, this aspect has not been well documented in the literature and offers scope for further investigation in the RAC setting, particularly in residents with various forms of cognitive impairment.

Further research is required to examine the role of residents and their family members/carers having a greater input into food, improved meal time ambience combined with a prophylactic high energy-high protein liberalised diet and a nutrition coordinator in the prevention and treatment of malnutrition. Evidence is also required for the cost-effectiveness and acceptability of nutrition interventions, with an emphasis on improving compliance and minimising food insecurity. Furthermore, to improve the quality of evidence available, nutrition interventions should report outcomes on global nutrition status as determined by a validated method of nutrition assessment in this setting.

9. Conclusion

Unfortunately malnutrition remains unacceptably high in the RAC setting, with at least half the RAC population at risk of developing malnutrition and approximately one-in-three residents assessed as malnourished. Identifying and managing malnutrition in the RAC setting will not only

likely improve associated health-related outcomes for individual residents but also possibly extend to benefitting the institutions and health care system economically. Although nutritional interventions which modify the method of food provision and/or provide ONS have demonstrated better outcomes for at risk and malnourished aged care residents, further research is required to improve the evidence base and inform aged care policy.

Contributors

All co-authors have contributed and agree to the content of the paper.

Conflict of interest

None

Funding

No funding was received for this manuscript. Co-authors contributed during their academic activities from Queensland University of Technology, Flinders University, and Bond University.

References

1. United Nations, Department of Economic and Social Affairs, Population Division (2013). World Population Ageing 2013; Available from: <http://www.un.org/en/development/desa/population/publications/pdf/ageing/WorldPopulationAgeing2013.pdf>. Accessed 10th July 2015.
2. WHO. Interesting facts about ageing. 2012; Available from: www.who.int/ageing/about_facts_en_index.html.pdf. Accessed 13th June 2013.
3. Lopez A, Mathers C, Ezzati M, Jamison D, Murray C. Global and regional burden of disease and risk factors, 2001: systematic analysis of population health data. *Lancet*. 2002;367(9524):1747-57.
4. Department of Health and Ageing. Residential Care Services. 2008; Available from: <http://www.health.gov.au/internet/main/publishing.nsf/Content/ageing-rescare-index.htm>. Accessed 30th August 2013.
5. AIHW 2014. Australia's health 2014. Canberra: Australian Institute of Health and Welfare. Report No.: Australia's health series no. 14. Cat. no. AUS 178. Available from: <http://www.aihw.gov.au/publication-detail/?id=60129547205>. Accessed 17th November 2015.
6. Chen CC-H, Schilling LS, Lyder CH. A concept analysis of malnutrition in the elderly. *J Adv Nurs*. 2001;36(1):131-42.
7. Beattie E, O'Reilly M, Strange E, Franklin S, Isenring E. How much do residential aged care staff members know about the nutritional needs of residents? *Int J Older People Nurs*. 2013;9(1):54-64.
8. Suominen M, Sandelin E, Soini H, Pitkala K. How well do nurses recognize malnutrition in elderly patients? . *Eur J Clin Nutr*. 2007;63(2):292-6.
9. Gaskill D, Black L, Isenring E, Hassall S, Sanders F, Bauer J. Malnutrition prevalence and nutrition issues in residential aged care facilities. *Australas J Ageing* 2008;27(4):189-94.
10. Geurden B, Franck E, Lopez Hartmann M, Weyler J, Ysebaert D. Prevalence of 'being at risk of malnutrition' and associated factors in adult patients receiving nursing care at home in Belgium. *Int J Nurs Pract*. 2015;21:635-44.
11. Bauer S, Halfens R, Lohrmann C. Knowledge and attitudes of nursing staff towards malnutrition care in nursing homes: a multicentre cross-sectional study. *J Nutr Health Aging*. 2015;19(7):734-40.
12. Sloane P, Ivey J, Helton M, Barrick A, Cerna A. Nutritional Issues in Long-Term Care. *J Am Med Dir Assoc*. 2008;9(7):476-85.
13. Morley J, Caplan G, Cesari M, Dong B, Flaherty J, Grossberg G, et al. International Survey of Nursing Home Research Priorities. *J Am Med Dir Assoc*. 2014;15(5):309-12.
14. Bell C, Tamura B, Masaki K, Amella E. Prevalence and Measures of Nutritional Compromise Among Nursing Home Patients: Weight Loss, Low Body Mass Index, Malnutrition, and Feeding Dependency, A Systematic Review of the Literature. *J Am Med Dir Assoc*. 2013;14(2):94-100.
15. Moreira NCF, Krausch-Hofmann S, Matthys C, Vereecken C, Vanhauwaert E, Declercq A, et al. Risk Factors for Malnutrition in Older Adults: A Systematic Review of the Literature Based on Longitudinal Data. *Adv Nutr*. 2016;7(3):507-22.
16. Abbott RA, Whear R, Thompson-Coon J, Ukoumunne OC, Rogers M, Bethel A, et al. Effectiveness of mealtime interventions on nutritional outcomes for the elderly living in residential care: A systematic review and meta-analysis. *Ageing Res Rev*. 2013;12(4):967-81.
17. Beck A, Wijnhoven H, Lassen K. A review of the effect of oral nutritional interventions on both weight change and functional outcomes in older nursing home residents. *E Spen Eur E J Clin Nutr Metab*. 2011;6(3):e101-e5.
18. Elia M, Normand C, Laviano A, Norman K. A systematic review of the cost and cost effectiveness of using standard oral nutritional supplements in community and care home settings. *Clin Nutr*. 2016;35(1):125-37.
19. National Centre for Classification in Health. The international statistical classification of diseases and related health problems, 10th revision, Australian modification (ICD-10-AM):

New South Wales, Australia: National Centre for Classification in Health, Faculty of Health Sciences, The University of Sydney; 2010.

20. Winter JE, MacInnis RJ, Wattanapenpaiboon N, Nowson CA. BMI and all-cause mortality in older adults: a meta-analysis. *Am J Clin Nutr.* 2014; 99(4):875-90.
21. Donini L, Savina C, Gennaro E, De Felice MR, Rosano A, Pandolfo MM, et al. A systematic review of the literature concerning the relationship between obesity and mortality in the elderly. *J Nutr Health Aging.* 2012;16(1):89-98.
22. Veronese N, Cereda E, Solmi M, Fowler S, Manzato E, Maggi S, et al. Inverse relationship between body mass index and mortality in older nursing home residents: a meta-analysis of 19,538 elderly subjects. *Obes Rev.* 2015;16(11):1001-15.
23. Volkert D. Malnutrition in Older Adults - Urgent Need for Action: A Plea for Improving the Nutritional Situation of Older Adults. *Gerontology.* 2013;59(4):328-33.
24. Green SM, Watson R. Nutritional screening and assessment tools for older adults: literature review. *J Adv Nurs.* 2006;54(4):477-90.
25. Rubenstein L, Harker J, Salva A, Guigoz Y, Vellas B. Screening for undernutrition in geriatric practice: Developing the Short-Form Mini Nutritional Assessment (MNA-SF). *J Gerontol.* 2001;56A:M366-M72.
26. White J, Dwyer J, Posner B, Ham R, Lipschitz D, Wellman N. Nutrition Screening Initiative: development and implementation of the public awareness checklist and screening tools. *J Am Diet Assoc* 1992;92(2):163-7.
27. Bouillanne O, Morineau G, Dupont C, Coulombel I, Vincent J-P, Nicolis I, et al. Geriatric Nutritional Risk Index: a new index for evaluating at-risk elderly medical patients. *Am J Clin Nutr.* 2005;82(4):777-83.
28. Wolinsky F, Coe R, McIntosh W, Kubena K, Prendergast J, Chavez M, et al. Progress in the Development of a Nutritional Risk Index. *J Nutr.* 1990;120(11 Suppl):1549-53.
29. Elia M. The 'MUST' report. Nutritional screening for adults: a multidisciplinary responsibility. Development and use of the 'Malnutrition Universal Screening Tool' (MUST) for adults: British Association for Parenteral and Enteral Nutrition (BAPEN); 2003.
30. Söderhamn U, Söderhamn O. Developing and testing the Nutritional Form For the Elderly. *Int J Nurs Pract.* 2001;7(5):336-41.
31. Laporte M, Villalon L, Payette H. Simple nutrition screening tools for healthcare facilities: development and validity assessment. *Can J Diet Pract Res.* 2001;62(1):26.
32. Blaum CS, O'Neill EF, Clements KM, Fries BE, Fiatarone MA. Validity of the minimum data set for assessing nutritional status in nursing home residents. *Am J Clin Nutr.* 1997;66(4):787-94.
33. Woo J, Chumlea W, Sun S, Kwok T, Lui H, Hui E, et al. Development of the Chinese Nutrition Screen (CNS) for use in institutional settings. *J Nutr Health Aging.* 2005;9(4):203-10.
34. Wilson M, Thomas D, Rubenstein L, Chibnall J, Anderson S, Baxi A, et al. Appetite assessment: simple appetite questionnaire predicts weight loss in community-dwelling adults and nursing home residents. *Am J Clin Nutr.* 2005;82(5):1074-81.
35. Kruizenga HM, De Vet HCW, Van Marissing CME, Stassen EEP, Strijk JE, Van Bokhorst-De Van Der Schueren MAE, et al. The SNAQRC, an easy traffic light system as a first step in the recognition of undernutrition in residential care. *J Nutr Health Aging.* 2010;14(2):83-9.
36. Dent E, Visvanathan R, Piantadosi C, Chapman I. Nutritional Screening Tools as Predictors of Mortality, Functional Decline, and Move to Higher Level Care in Older People: A Systematic Review. *J Nutr Gerontol Geriatr.* 2012;31(2):97-145.
37. van Bokhorst-de van der Schueren M, Guaitoli P, Jansma E, de Vet H. A systematic review of malnutrition screening tools for the nursing home setting. *J Am Med Dir Assoc.* 2014;15(3):171-84.
38. White J, Guenter P, Jensen G, Malone A, Schofield M, Academy Malnutrition Work Group, et al. Consensus Statement: Academy of Nutrition and Dietetics and American Society for Parenteral and Enteral Nutrition. *JPEN J Parenter Enteral Nutr* 2012;36(3):275-83.
39. Detsky A. What is subjective global assessment of nutritional status? *JPEN J Parenter Enteral Nutr* 1987;11(1):8-13.
40. Byles J, Perry L, Parkinson L, Bellchambers H, Moxey A, Howie A, et al. Encouraging best practice nutrition and hydration in residential aged care. Final report: Australian

Government Department of Health & Ageing 2009. Available from:
https://www.newcastle.edu.au/data/assets/pdf_file/0017/34037/EBPRAC-report.pdf.
Accessed 10th November 2015.

41. Guigoz Y, Vellas B, Garry P. Assessing the Nutritional Status of the Elderly: The Mini Nutritional Assessment as part of the Geriatric Evaluation. *Nutr Rev.* 1996;54:S59-S65.
42. Kuzuya M, Kanda S, Koike T, Suzuki Y, Satake S, Iguchi A. Evaluation of mini-nutritional assessment for Japanese frail elderly. *Nutrition.* 2005;21(4):498-503.
43. Sacks GS, Dearman K, Replogle WH, Cora VL, Meeks M, Canada T. Use of subjective global assessment to identify nutrition-associated complications and death in geriatric long-term care facility residents. *J Am Coll Nutr.* 2000;19(5):570-7.
44. Compan B, Di Castri A, Plaze J, Arnaud-Battandier F. Epidemiological Study of Malnutrition in Elderly Patients in Acute, Sub-Acute and Long-Term Care Using the MNA [R]. *J Nutr Health Aging.* 1999;3:146-51.
45. Saletti A, Lindgren EY, Johansson L, Cederholm T. Nutritional status according to mini nutritional assessment in an institutionalized elderly population in Sweden. *Gerontology.* 2000;46(3):139-45.
46. Pezzana A, Cereda E, Avagnina P, Malfi G, Paiola E, Frighi Z, et al. Nutritional care needs in elderly residents of long-term care institutions: Potential implications for policies. *J Nutr Health Aging.* 2015;19(9):947-54.
47. Nazemi L, Skoog I, Karlsson I, Hosseini S, Mohammadi MR, Hosseini M, et al. Malnutrition, Prevalence and Relation to Some Risk Factors among Elderly Residents of Nursing Homes in Tehran, Iran. *Iran J Public Health.* 2015;44(2):218-27.
48. Bonaccorsi G, Collini F, Castagnoli M, Di Bari M, Cavallini MC, Zaffarana N, et al. A cross-sectional survey to investigate the quality of care in Tuscan (Italy) nursing homes: the structural, process and outcome indicators of nutritional care. *BMC Health Serv Res.* 2015;15(1):1-10.
49. Suominen M, Muurinen S, Routasalo P, Soini H, Suur-Uski I, Peiponen A, et al. Malnutrition and associated factors among aged residents in all nursing homes in Helsinki. *Eur J Clin Nutr.* 2005;59(4):578-83.
50. Schönherr S, Halfens R, Meijers J, Schols J, Lohrmann C. Structural and process indicators of nutritional care: A comparison between austrian hospitals and nursing homes. *Nutrition.* 2012;28(9):868-73.
51. Nijs K, de Graaf C, van Staveren WA, de Groot LCPGM. Malnutrition and Mealtime Ambiance in Nursing Homes. *J Am Med Dir Assoc.* 2009;10(4):226-9.
52. Morley J. Anorexia of aging: Physiologic and pathologic. *Am J Clin Nutr.* 1997;66(4):760-73.
53. Robbins L. Evaluation of weight loss in the elderly. *Geriatrics.* 1989;44(4):31-7.
54. Australian and New Zealand Society for Geriatric Medicine (Position Statement No. 6). Under-nutrition and the Older Person. *Australas J Ageing* 2009;28(2):99-105.
55. Forster S, Gariballa S. Age as a determinant of nutritional status: A cross sectional study. *Nutrition.* 2005;4:28.
56. Perera R, Ekanayake L. Relationship between nutritional status and tooth loss in an older population from Sri Lanka. *Gerodontology.* 2012;29(2):e566-e70.
57. Stanga Z. Basics in clinical nutrition: Nutrition in the elderly. *E Spen Eur E J Clin Nutr Metab.* 2009;4(6):e289-e99.
58. Kuczmarski M, Weddle D. Position paper of the American Dietetic Association: Nutrition across the spectrum of aging. *J Am Diet Assoc* 2005;105(4):616-33.
59. Namasivayam A, Steele C. Malnutrition and Dysphagia in long-term care: a systematic review. *J Nutr Gerontol Geriatr.* 2015;34(1):1-21.
60. Tamura B, Bell C, Masaki K, Amella E. Factors associated with weight loss, low BMI, and malnutrition among nursing home patients: a systematic review of the literature. *J Am Med Dir Assoc.* 2013;14(9):649-55.
61. Beck A. Weight loss, mortality and associated potentially modifiable nutritional risk factors among nursing home residents- A Danish follow-up study. *J Nutr Health Aging.* 2015;19(1):96-101.
62. Carrier N, Ouellet D, West G. Nursing home food services linked with risk of malnutrition. *Can J Diet Pract Res.* 2007;68(1):14-20.

63. Power E. Individual and household food insecurity in Canada: position of Dietitians of Canada. *Can J Diet Pract Res.* 2005;66(1):43.
64. Vahabi M, Martin LS. Food security: Who is being excluded? A case of older people with dementia in long-term care homes. *J Nutr Health Aging.* 2014;18(7):685-91.
65. Bauer S, Halfens RJG, Lohrmann C. Knowledge and attitudes of nursing staff towards malnutrition care in nursing homes: A multicentre cross-sectional study. *J Nutr Health Aging.* 2015;19(7):734-40.
66. Pleuss J. Alterations in nutritional status. In: Porth C, editor. *Pathophysiology, concepts of altered health states.* 7th ed. Philadelphia: Lippincott Williams & Wilkins; 2005.
67. Marzetti E, Leeuwenburgh C. Skeletal muscle apoptosis, sarcopenia and frailty at old age. *Exp Geront.* 2006;41(12):1234-8.
68. Mueller DH. Medical nutrition therapy for pulmonary disease. In: Mahan LK, Escott-Stump S, editors. *Krouse's food & nutrition therapy.* Missouri: Saunders Elsevier; 2008. p. 899 - 920.
69. Landi F, Liperoti R, Russo A, Giovannini S, Tosato M, Capoluongo E, et al. Sarcopenia as a risk factor for falls in elderly individuals: Results from the iLSIRENTE study. *Clin Nutr.* 31(5):652-8.
70. Hickson M. Malnutrition and ageing. *Postgrad Med J.* 2006;82(963):2-8.
71. Diekmann R, Winning K, Uter W, MJ K, Sieber C, Volkert D, et al. Screening for malnutrition among nursing home residents -- a comparative analysis of the Mini Nutritional Assessment, the Nutritional Risk Screening, and the Malnutrition Universal Screening Tool. *J Nutr Health Aging.* 2013;17(4):326-31.
72. Lilamand M, Kelaiditi E, Demougeot L, Rolland Y, Vellas B, Cesari M. The Mini Nutritional Assessment-Short Form and mortality in nursing home residents — Results from the INCUR study. *J Nutr Health Aging.* 2015;19(4):383-8.
73. Chan M, Lim Y, Ernest A, Tan T. Nutritional assessment in an Asian nursing home and its association with mortality. *J Nutr Health Aging.* 2010;14(1):23-8.
74. Hanson L, Ersek M, Lin F, Carey T. Outcomes of Feeding Problems in Advanced Dementia in a Nursing Home Population. *J Am Geriatr Soc.* 2013;61(10):1692-7.
75. Jongenelis K, Pot A, Eisses A, Beekman A, Kluiters H, Ribbe M. Prevalence and risk indicators of depression in elderly nursing home patients: the AGED study. *J Affect Disord.* 2004;83(2-3):135-42.
76. Webber AP, Martin JL, Harker JO, Josephson KR, Rubenstein LZ, Alessi CA. Depression in older patients admitted for postacute nursing home rehabilitation. *J Am Geriatr Soc.* 2005;53(6):1017-22.
77. Blaum CS, Fries BE, Fiatarone MA. Factors Associated With Low Body Mass Index and Weight Loss in Nursing Home Residents. *J Gerontol A Biol Sci Med Sci.* 1995;50(3):M162-M168.
78. Smoliner C, Norman K, Wagner K-H, Hartig W, Lochs H, Pirlich M. Malnutrition and depression in the institutionalised elderly. *Br J Nutr.* 2009;102(11):1663-7.
79. Cabrera MAS, Mesas AE, Garcia ARL, de Andrade SM. Malnutrition and Depression among Community-dwelling Elderly People. *J Am Med Dir Assoc.* 2007;8(9):582-4.
80. German L, Feldblum I, Bilenko N, Castel H, Harman-Boehm I, Shahar DR. Depressive symptoms and risk for malnutrition among hospitalized elderly people. *J Nutr Health Aging.* 2008 2008/05/01;12(5):313-8.
81. Morley J, Kraenzle D. Causes of Weight Loss in a Community Nursing Home. *J Am Geriatr Soc.* 1994;42(6):583-5.
82. Rasheed S, Woods RT. An investigation into the association between nutritional status and quality of life in older people admitted to hospital. *J Hum Nutr Diet.* 2014;27(2):142-51.
83. Freijer K, Tan S, Koopmanschap M, Meijers J, Halfens R, Nuijten M. The economic costs of disease related malnutrition. *Clin Nutr.* 2013;32(1):136-41.
84. Elia M, Stratton R, Russell C, Green C, Pang F. The cost of disease-related malnutrition in the UK and economic considerations for the use of oral nutritional supplements (ONS) in adults. Redditch, UK: British Association for Parenteral and Enteral Nutrition 2006. Available from: http://www.bapen.org.uk/pdfs/health_econ_exec_sum.pdf. Accessed on 10th September 2012.

85. Watterson C, Fraser A, Banks M, Isenring E, Miller M, Silvester C, et al. Evidence based practice guidelines for the nutritional management of malnutrition in patients across the continuum of care. *Nutr Diet*. 2009;66(Suppl 3):S1 - S34.
86. Marshall S, Bauer J, Capra S, Isenring E. Are informal carers and community care workers effective in managing malnutrition in the older adult community? A systematic review of current evidence. *J Nutr Health Aging*. 2013; 17(8):645-51.
87. National Collaborating Centre for Acute Care (UK). Nutrition support for adults: oral nutrition support, enteral tube feeding and parenteral nutrition. London: National Collaborating Centre for Acute Care (UK); 2006 Feb. (NICE Clinical Guidelines, No. 32.). Available from: <http://www.ncbi.nlm.nih.gov/books/NBK49269/>. Accessed on July 14th 2011.
88. Abbey K, Wright O, Capra S. Menu Planning in Residential Aged Care—The Level of Choice and Quality of Planning of Meals Available to Residents. *Nutrients*. 2015;7(9):7580-92.
89. Wendland BE, Greenwood CE, Weinberg I, Young KWH. Malnutrition in Institutionalized Seniors: The Iatrogenic Component. *J Am Geriatr Soc*. 2003;51(1):85-90.
90. Ball MM, Whittington FJ, Perkins MM, Patterson VL, Hollingsworth C, King SV, et al. Quality of Life in Assisted Living Facilities: Viewpoints of Residents. *J Appl Gerontol*. 2000;19(3):304-25.
91. Wright ORL, Connelly LB, Capra S, Hendrikz J. Determinants of foodservice satisfaction for patients in geriatrics/rehabilitation and residents in residential aged care. *Health Expect*. 2013;16(3):251-65.
92. Ducak K, Keller H. Menu planning in long-term care: Toward resident-centred menus. *Can J Diet Pract Res*. 2011;72(2):e126-e33
93. Kuosma K, Hjerrild J, Pedersen PU, Hundrup YA. Assessment of the nutritional status among residents in a Danish nursing home - health effects of a formulated food and meal policy. *J Clin Nurs*. 17(17):2288-93.
94. Schonherr S, Halfens RJ, Lohrmann C. Development and psychometric evaluation of the Knowledge of Malnutrition - Geriatric (KoM-G) questionnaire to measure malnutrition knowledge among nursing staff in Austrian nursing homes. *Scand J Caring Sci*. 2015;29(1):193-202.
95. Bell CL, Lee AS, Tamura BK. Malnutrition in the nursing home. *Curr Opin Clin Nutr Metab Care*. 2015;18(1):17-23.
96. Gaskill D, Isenring EA, Black LJ, Hassall S, Bauer JD. Maintaining nutrition in aged care residents with a train-the-trainer intervention and nutrition coordinator. *J Nutr Health Aging*. 2009;13(10):913-7.
97. Lorefalt B, Wissing U, Unosson M. Smaller but energy and protein enriched meals improve energy and nutrient intakes in elderly patients. *J Nutr Health Aging*. 2005;9(4):243-7.
98. Mila R, Abellana R, Padro L, Basulto J, Farran A. High consumption foods and their influence on energy and protein intake in institutionalized older adults. *J Nutr Health Aging*. 2012;16(2):115-22.
99. Lam ITY, Keller HH, Pfisterer K, Duizer L, Stark K, Duncan AM. Micronutrient Food Fortification for Residential Care: A Scoping Review of Current Interventions. *J Am Med Dir Assoc*. In Press.
100. Odlund Olin A, Armyr I, Soop M, Jerstrom S, Classon I, Cederholm T, et al. Energy-dense meals improve energy intake in elderly residents in a nursing home. *Clin Nutr*. 2003 Apr;22(2):125-31.
101. Leslie W, Woodward M, Lean M, Theobald H, Watson L, Hankey C. Improving the dietary intake of under nourished older people in residential care homes using an energy-enriching food approach: a cluster randomised controlled study. *J Hum Nutr Diet*. 2013; 26(4):387-394.
102. Smoliner C, Norman K, Scheufele R, Hartig W, Pirlich M, Lochs H. Effects of food fortification on nutritional and functional status in frail elderly nursing home residents at risk of malnutrition. *Nutrition*. 2008; 24(11-12):1139-44.
103. Iuliano S, Woods J, Robbins J. Consuming two additional serves of dairy food a day significantly improves energy and nutrient intakes in ambulatory aged care residents: A feasibility study. *J Nutr Health Aging*. 2013;17(6):509-13.

104. ÖDlund Olin A, Armyr I, Soop M, Jerström S, Classon I, Cederholm T, et al. Energy-dense meals improve energy intake in elderly residents in a nursing home. *Clin Nutr.* 2003;22(2):125-31.
105. Beck AM, Damkjaer K, Tetens I. Lack of compliance of staff in an intervention study with focus on nutrition, exercise and oral care among old (65+ yrs) Danish nursing home residents. *Aging Clin Exp Res.* 2009;21(2):143-9.
106. Iuliano S, Olden A, Woods J. Meeting the nutritional needs of elderly residents in aged-care: Are we doing enough? *J Nutr Health Aging.* 2013;17(6):503-8.
107. Darmon P, Kaiser MJ, Bauer JM, Sieber CC, Pichard C. Restrictive diets in the elderly: Never say never again? *Clin Nutr.* 2010;29(2):170-4.
108. Germain I, Dufresne T, Gray-Donald K. A Novel Dysphagia Diet Improves the Nutrient Intake of Institutionalized Elders. *J Am Diet Assoc.* 2006;106(10):1614-23.
109. Bannerman E, McDermott K. Dietary and Fluid Intakes of Older Adults in Care Homes Requiring a Texture Modified Diet: The Role of Snacks. *J Am Med Dir Assoc.* 2011;12(3):234-9.
110. Massoulard A, Bonnabau H, Gindre-Poulvelarie L, Baptistev A, Preux P-M, Villemonteix C, et al. Analysis of the food consumption of 87 elderly nursing home residents, depending on food texture. *J Nutr Health Aging.* 2011;15(3):192-5.
111. Keller H, Chambers L, Niezgodá H, Duizer L. Issues associated with the use of modified texture foods. *J Nutr Health Aging.* 2012;16(3):195-200.
112. Nijs KA, de Graaf C, Siebelink E, Blauw YH, Vanneste V, Kok FJ, et al. Effect of family-style meals on energy intake and risk of malnutrition in dutch nursing home residents: a randomized controlled trial. *J Gerontol A Biol Sci Med Sci.* 2006;61(9):935-42.
113. Collins J, Porter J. The effect of interventions to prevent and treat malnutrition in patients admitted for rehabilitation: a systematic review with meta-analysis. *J Hum Nutr Diet.* 2015;28(1):1-15.
114. Beck AM, Ovesen L, Schroll M. Home-made oral supplement as nutritional support of old nursing home residents, who are undernourished or at risk of undernutrition based on the MNA. A pilot trial. *Mini Nutritional Assessment.* *Aging Clin Exp Res.* 2002;14(3):212-5.
115. Manders M, Groot CP, Blauw YH, Dhonukshe-Rutten RA, Hoeckel-Prüst L, Bindels JG, et al. Effect of a nutrient-enriched drink on dietary intake and nutritional status in institutionalised elderly. *Eur J Clin Nutr.* 2009; 63(10): 1241–1250.
116. Stange I, Bartram M, Liao Y, Poeschl K, Kolpatzik S, Uter W, et al. Effects of a low-volume, nutrient- and energy-dense oral nutritional supplement on nutritional and functional status: a randomized, controlled trial in nursing home residents. *J Am Med Dir Assoc.* 2013;14(8):628.e1-8.
117. Pouyssegur V, Brocker P, Schneider SM, Philip JL, Barat P, Reichert E, et al. An innovative solid oral nutritional supplement to fight weight loss and anorexia: Open, randomised controlled trial of efficacy in institutionalised, malnourished older adults. *Age Ageing.* 2015;44(2): 245-251.
118. Jobse I, Liao Y, Bartram M, Delantonio K, Uter W, Stehle P, et al. Compliance of nursing home residents with a nutrient- and energy-dense oral nutritional supplement determines effects on nutritional status. *J Nutr Health Aging.* 2015;19(3):356-64.
119. Rondanelli M, Opizzi A, Antonello N, Boschi F, Iadarola P, Pasini E, et al. Effect of essential amino acid supplementation on quality of life, amino acid profile and strength in institutionalized elderly patients. *Clin Nutr.* 2011; 30(5): 571-577.
120. Lee LC, Tsai AC, Wang JY. Need-based nutritional intervention is effective in improving handgrip strength and Barthel Index scores of older people living in a nursing home: a randomized controlled trial. *Int J Nurs Stud.* 2015;52(5):904-12.
121. Fiatarone M, O'Neill E, Ryan N, Clements K, Solares G, Nelson M, et al. Exercise training and nutritional supplementation for physical frailty in very elderly people. *N Engl J Med.* 1994;330(25):1769-75.
122. Beck AM, Damkjaer K, Sørbye LW. Physical and social functional abilities seem to be maintained by a multifaceted randomized controlled nutritional intervention among old (>65 years) Danish nursing home residents. *Arch Gerontol Geriatr.* 2010;50(3): 351-55.

123. Lee LC, Tsai AC, Wang JY, Hurng BS, Hsu HC, Tsai HJ. Need-based intervention is an effective strategy for improving the nutritional status of older people living in a nursing home: a randomized controlled trial. *Int J Nurs Stud*. 2013;50(12):1580-8.
124. Hines S, Wilson J, McCrow J, Abbey J, Sacre S. Oral liquid nutritional supplements for people with dementia in residential aged care facilities: a systematic review. *JBI Database System Rev Implement Rep*. 2010;8(1):1-52.
125. Mitchell S, Teno J, Roy J, Kabumoto G, Mor V. Clinical and organizational factors associated with feeding tube use among nursing home residents with advanced cognitive impairment. *JAMA*. 2003;290(1):73-80.
126. Kimyagarov S, Turgeman D, Fleissig Y, Klid R, Kopel B, Adunsky A. Percutaneous endoscopic gastrostomy (PEG) tube feeding of nursing home residents is not associated with improved body composition parameters. *J Nutr Health Aging*. 2013;17(2):162-5.
127. Peck A, Cohen C, Mulvihill M. Long-Term Enteral Feeding of Aged Demented Nursing Home Patients. *J Am Geriatr Soc*. 1990;38(11):1195-8.
128. Banks MD, Graves N, Bauer JD, Ash S. The costs arising from pressure ulcers attributable to malnutrition. *Clin Nutr*. 2010;29(2):180-6.
129. Baulderstone L, Yaxley A, Luszcz M, Miller M. Diet liberalisation in older Australians decreases risk of frailty without increasing the risk of developing chronic disease. *JFA*. 2012;1(4):174-82.
130. Meijers J, Halfens R, Wilson L, Schols J. Estimating the costs associated with malnutrition in Dutch nursing homes. *Clin Nutr*. 2012;31(1):65-8.
131. White J, Guenter P, Jensen G, Malone A, Schofield M, Academy Malnutrition Work Group, et al. Consensus Statement of the Academy of Nutrition and Dietetics/American Society for Parenteral and Enteral Nutrition: Characteristics Recommended for the Identification and Documentation of Adult Malnutrition (Undernutrition). *J Acad Nutr Diet*. 2012;112(5):730-8.
132. Bottone F, Hawkins K, Musich S, Cheng Y, Ozminkowski R, Migliori R, et al. The relationship between body mass index and quality of life in community-living older adults living in the United States. *J Nutr Health Aging*. 2013;17(6):495-501.
133. Crogan NL, Pasvogel A. The Influence of Protein-Calorie Malnutrition on Quality of Life in Nursing Homes. *J Gerontol A Biol Sci Med Sci*. 2003; 58(2):M159-M64.
134. Webber A, Martin J, Harker J, Josephson K, Rubenstein L, Alessi C. Depression in Older Patients Admitted for Postacute Nursing Home Rehabilitation. *J Am Geriatr Soc*. 2005;53(6):1017-22.

Figure 1: Aetiology of malnutrition in the elderly (Adapted from 15, 33)




Table 1: Cause and consequences of malnutrition (Compiled from 32, 38, 52, 71-75, 79, 80, 87, 102, 132-134)

Cause	Consequence
Physiological	
Impaired immune response	Increased susceptibility to infection and impaired recovery from infection
Impaired wound healing	Surgical wound dehiscence, anastomotic breakdowns, post-surgical fistulae
Reduced muscle mass and strength	Limited physical function, increased disability, increased risk of falls, pressure ulcers, decreased activity leading to increased risk of thromboembolism
Reduced subcutaneous fat for cushioning	Increased risk of fractures in accidents, falls
Reduced respiratory function	Poor cough pressure, increased susceptibility to respiratory infection, fatigue, poor breathing control, pulmonary oedema
Water and electrolyte imbalances	Depletion of potassium, magnesium and phosphate stores combined with excess sodium and water leading to increased risk of re-feeding syndrome and iatrogenic sodium and water overload
Impaired thermoregulation	Hypothermia and increased risk of falls
Vitamin and mineral deficiencies	Specific deficiency states including scurvy and Wernicke-Korsakoff syndrome, iron-deficiency and pernicious anaemia or magnesium deficiency
Reduced hepatic function	Anaemia, fatigue, oedema
Reduced cardiac muscle	Decreased cardiac output

	Mortality
Psychosocial	
Lower quality of life	Apathy, depression, self-neglect, hypochondriasis, poor self-efficacy, poor body image, confusion, decreased interest in food, loss of libido, less engagement in social activities due to the overall physiological burden of malnutrition