

Malnutrition coding shortfalls in Australian and New Zealand hospitals

Agarwal, Ekta; Ferguson, Maree; Banks, Merrilyn; Bauer, Judith; Capra, Sandra; Isenring, Elisabeth

Published in:
Nutrition and Dietetics

DOI:
[10.1111/1747-0080.12116](https://doi.org/10.1111/1747-0080.12116)

Licence:
CC BY-NC-ND

[Link to output in Bond University research repository.](#)

Recommended citation(APA):
Agarwal, E., Ferguson, M., Banks, M., Bauer, J., Capra, S., & Isenring, E. (2015). Malnutrition coding shortfalls in Australian and New Zealand hospitals. *Nutrition and Dietetics*, 72(1), 69-73. <https://doi.org/10.1111/1747-0080.12116>

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

For more information, or if you believe that this document breaches copyright, please contact the Bond University research repository coordinator.

March, 2015

Malnutrition coding shortfalls in Australian and New Zealand hospitals

Ekta Agarwal, *The University of Queensland, Australia*

Maree Ferguson, *University of Queensland*

Merrilyn Banks, *The University of Queensland, Australia*

Judith Bauer, *The University of Queensland*

Sandra Capra, *The University of Queensland, et al.*



1 Authors:

- 2 • Ekta Agarwal (PhD, APD) Lecturer (Queensland University of Technology,
3 Kelvin Grove, QLD 4059), Australia; Adjunct Dietitian Princess Alexandra
4 Hospital, Ipswich Road, Woolloongabba, QLD 4102); and PhD candidate
5 (Centre for Dietetics Research (C-DIET-R), The University of Queensland,
6 Brisbane, St Lucia, Qld 4072);
- 7 • Maree Ferguson (PhD, MBA, AdvAPD), Director Nutrition and Dietetics
8 (Princess Alexandra Hospital, Ipswich Road, Woolloongabba, QLD 4102);
9 Adjunct Senior Lecturer (C-DIET-R, The University of Queensland, Brisbane,
10 St Lucia, Qld 4072, Australia);
- 11 • Merrilyn Banks (PhD, AdvAPD), Director Nutrition and Dietetics (Royal
12 Brisbane and Women's Hospital, Herston, QLD 4029); Adjunct Senior Lecturer
13 (C-DIET-R, The University of Queensland, Brisbane, St Lucia, Qld 4072,
14 Australia);
- 15 • Judith Bauer (PhD, FDAA, AdvAPD), Associate Professor (C-DIET-R, The
16 University of Queensland, Brisbane, St Lucia, Qld 4072, Australia);
- 17 • Sandra Capra (AM, PhD, FDAA, AdvAPD), Professor (C-DIET-R, The
18 University of Queensland, Brisbane, St Lucia, Qld 4072, Australia);
- 19 • Elisabeth Isenring (PhD, AdvAPD), Clinical Academic Fellow (Princess
20 Alexandra Hospital) and Conjoint Senior Lecturer (C-DIET-R, The University
21 of Queensland, Brisbane, St Lucia, Qld 4072, Australia).

22

23

24

25 Corresponding author:

26 Ekta Agarwal, Lecturer (Nutrition and Dietetics), School of Exercise and Nutrition
27 Sciences, Queensland University of Technology, Victoria Park Road, Kelvin Grove,
28 QLD 4059.

29 Tel: +61-7- 31387977

30 Fax: +61-7-31383980

31 Email: ekta.agarwal@qut.edu.au

32

1 **Conflict of interests**

2 The authors declare that there are no conflicts of interest.

3

4 **Funding**

5 EA is a recipient of the Australian Post-Graduate Award (APA) and the Small Research
6 Grant (2010) from the Australasian Society of Parenteral and Enteral Nutrition
7 (AuSPEN). The authors would like to acknowledge AuSPEN for organising webinars
8 for training dietitians involved with data collection; and Queensland Health for funding
9 Queensland-based hospitals to recruit additional dietitians to assist with data collection
10 in the ANCDS.

11

12 **Authorship**

13 This project was undertaken as part of the PhD study of EA and was supervised by EI,
14 MF and MB. The project was conceptualised, planned and designed by EA, EI, MF and
15 MB. Data collection was coordinated, acquired, analysed and interpreted by EA. The
16 original manuscript was written by EA, and all authors participated in editing and final
17 revisions. All authors have read and approved the final manuscript.

18

19 **Ethics Statement**

20 Ethics approval for the study was provided by the Medical and Research Ethics
21 Committee of The University of Queensland and local Human Research Ethics
22 Committees of participating hospitals.

23

24

25

26

27

28

29

30

1 **Title:** Malnutrition coding shortfalls in Australian and New Zealand hospitals

2

3 **ABSTRACT**

4 Aim: The International Classification of Diseases, Version 10, Australian modification
5 (ICD-10-AM) is used to classify diseases in hospital patients in Australia and New
6 Zealand. ICD-10-AM defines malnutrition as “BMI < 18.5 kg/m² or unintentional
7 weight loss of ≥ 5% with evidence of suboptimal intake resulting in subcutaneous fat
8 loss and/or muscle wasting.” The Australasian Nutrition Care Day Survey (ANCDS) is
9 the most comprehensive survey to evaluate malnutrition prevalence in acute care
10 patients from Australian and New Zealand hospitals. This study determined if
11 malnourished participants were assigned malnutrition-related codes according to ICD-
12 10-AM.

13 Methods: The ANCDS recruited acute care patients from 56 hospitals. Hospital-based
14 dietitians evaluated participants’ nutritional status using BMI and Subjective Global
15 Assessment (SGA). In keeping with the ICD-10-AM definition, malnutrition was
16 defined as BMI <18.5kg/m², SGA-B (moderately malnourished) or SGA-C (severely
17 malnourished). After three months, in this prospective cohort study, staff members from
18 each hospital’s health information/medical records department provided coding results
19 for malnourished participants.

20 Results: Malnutrition was prevalent in 30% (n=869) of the cohort (N=2976) and a
21 significantly small number of malnourished patients were coded for malnutrition (n=
22 162, 19%, $p<0.001$). In 21 hospitals, none of the malnourished participants were coded.

23 Conclusion: This is the largest study to provide a snapshot of malnutrition coding in
24 Australian and New Zealand hospitals. Findings highlight gaps in malnutrition

1 documentation and/or subsequent coding, which could potentially result in significant
2 loss of casemix-related revenue for hospitals. Dietitians must lead the way in
3 developing structured processes for malnutrition identification, documentation and
4 coding.

5 (246 words)

6 *This abstract will be presented at the 30th Dietitians Association of Australia National*
7 *Conference (23-25 May 2013, Canberra, Australia)*

8

9

10 **Keywords:** malnutrition, casemix, coding, International Classification of Diseases,
11 hospitals

12

13

14

15

16

17

18

19

20

21

22

23

24

1 INTRODUCTION

2 Malnutrition, a common problem in hospital patients, may occur as a result of
3 inadequate dietary intake and/or increased nutritional requirements, impaired nutrient
4 absorption, transport, and/or utilisation.¹ International studies report that malnutrition
5 affects 20 – 50% of hospital patients.^{2,3} The Australasian Nutrition Care Day Survey
6 (ANCDS) evaluated the nutritional status of 3122 adult patients admitted in 370 acute
7 care wards across 56 Australian and New Zealand hospitals, making this the largest
8 study thus far to report malnutrition point prevalence in Australian and New Zealand
9 hospitals.⁴ The survey found that one-in-three patients were malnourished⁴ and also
10 established an independent association between malnutrition and increased length of
11 stay (LOS) in hospital and higher incidence of in-hospital mortality.^{4,5}

12 A small number of cost-of-illness studies indicate higher hospital costs for
13 malnourished patients (attributable to longer hospital stay, higher incidence of
14 complications, increased costs of care and services) compared with well-nourished
15 patients.⁶⁻⁹ Despite the high prevalence and adverse consequences of malnutrition,
16 Australian literature continues to report that malnourished patients are often not
17 identified during hospitalisation.¹⁰⁻¹⁴ In 2009, Watterson et al published “Evidence
18 based practice guidelines for the nutritional management of malnutrition in adult
19 patients across the continuum of care”.¹⁵ These guidelines, endorsed by the Dietitians
20 Association of Australia (DAA) and Dietitians NZ, recommend the use of a number of
21 validated and reliable nutrition screening tools and assessment methods to identify
22 malnutrition.¹⁵ However, an evaluation of nutrition screening practices in 370 wards
23 that participated in the ANCDS revealed that nutrition risk screening was not routinely
24 performed in one-in-three wards.¹⁰

1 Identification of malnourished patients serves two purposes: it allows for the
2 (a) implementation of appropriate nutrition intervention/s to manage malnutrition¹⁶; and
3 (b) malnutrition coding using the diagnosis-related group (DRG) system.

4 Australian and New Zealand hospitals use the Australian Refined DRG (AR-DRG) to
5 classify the mix of cases (or ‘casemix’) according to their diagnoses.^{17,18} Medical coders
6 review documentation in medical charts and, based on the reason for hospitalisation and
7 coexisting comorbidities and complications, classify patients into casemix categories.¹⁸
8 The DRG-system is predicated on cases within a group being likely to utilise similar
9 levels of hospital resources thereby incurring similar costs.^{18,19}

10 The inclusion of malnutrition can potentially influence the DRG by resulting in a higher
11 classification and attracting greater financial reimbursement for the hospital.²⁰ However,
12 if a patient already has complex comorbidities, then the single effect of malnutrition
13 will not make a difference in the case severity, and therefore in reimbursement.²¹
14 Regardless, since medical coders depend on documentation to assign casemix
15 classification and coding, the quality of the clinician’s documentation of patients’
16 diagnoses in medical charts is important.²²

17

18 To the best of our knowledge, practices related to malnutrition coding in New Zealand
19 hospitals have never been reported. Since 1997, only four Australian studies have
20 reported malnutrition coding practices in hospitals and findings from all four studies
21 indicate that malnutrition was often not coded, therefore resulting in considerable loss
22 of casemix-related reimbursement for hospitals.^{11,19,23,24} Three of these studies reported
23 coding practices in individual hospitals and therefore their results cannot be generalised
24 across Australian and New Zealand hospitals.^{11,19,23} Rowell and Jackson (2011)

1 evaluated malnutrition coding for >250,000 inpatients admitted in 45 public hospitals in
2 Victoria and found that less than 2% of the patients had been coded for malnutrition.²⁴
3 This is perhaps the most comprehensive report on malnutrition coding practices in
4 Australian hospitals, however as the results are limited to hospitals in Victoria, they
5 cannot be generalised across Australia and New Zealand.
6 The present study aims to provide an insight into malnutrition coding for malnourished
7 patients from the ANCDS cohort.

8

9

10 **METHODS**

11 This was a prospective cohort study. Nutritional status of ANCDS participants was
12 assessed by dietitians from participating hospitals⁴ as follows: Dietitians screened
13 participants for nutrition risk using the validated Malnutrition Screening Tool (MST).²⁵
14 The MST is a two-question screening tool (appetite and recent unintentional weight
15 loss) and provides a score between
16 0–5.²⁵ Patients are considered at nutritional risk if they score ≥ 2 .²⁵ Participants who
17 were deemed “at risk” (MST score ≥ 2) underwent comprehensive nutrition assessment
18 using Subjective Global Assessment (SGA).²⁶ The SGA includes two components:
19 medical (changes in weight, dietary intake, gastrointestinal symptoms, nutrition-related
20 functional capacity) and physical (evidence of oedema, ascites, loss of subcutaneous fat
21 and muscle). Results from both components are combined to provide an overall rating:
22 well-nourished (SGA-A), moderately malnourished (SGA-B) or severely malnourished
23 (SGA-C). Dietitians also recorded participants’ weight and height, based on which BMI
24 was calculated.

1 The International Classification of Disease and Related Health Problems, 10th version,
2 Australian modification (ICD-10-AM) defines malnutrition in adults as “BMI <18.5
3 kg/m² or unintentional weight loss of at least 5% with evidence of sub-optimal intake
4 resulting in subcutaneous fat loss and/or muscle wasting”.²⁷ Therefore, in keeping with
5 this definition, patients with BMI <18.5 kg/m² or assessed as SGA-B or SGA-C were
6 included in the “malnourished” category.⁴ Patients with MST scores of <2 or an SGA
7 rating of well-nourished (SGA-A) were categorised as “well-nourished”.

8

9 Three months after evaluating ANCDS participants’ nutritional status, malnutrition
10 coding data were compiled by staff members of health information records departments
11 of participating hospitals. Admission-related information, including admission status
12 (emergency, elective or other), Australian Refined Diagnosis Related Group codes (AR-
13 DRG, reflecting participants’ clinical diagnosis), Patient Clinical Complexity Level
14 Scores (PCCL, reflecting participants’ disease severity) and type of admission (medical,
15 surgical, and other) were recorded. These have been reported elsewhere.⁵

16

17 For purpose of data cleaning, participants were excluded from analyses if post-three
18 months data were missing, and if LOS \leq 1 day (additional diagnoses for the episode of
19 admission was unlikely to be recorded if the LOS was \leq 1day.²⁸ Ethics approval for the
20 study was provided by [Removed for blind peer review].

21 **Statistical analyses:**

22 Data were analysed using IBM SPSS Statistics 21. Data are presented as frequency and
23 percentage. Chi-square tests were used to determine malnutrition coding according to
24 participants’ nutritional status.

1 **RESULTS**

2 Of the 3122 participants recruited in the ANCDs⁴, 146 were excluded during data
3 cleaning (Missing data: n=111 (3%); LOS \leq 1 day: n= 35 (1%)). Of the remaining 2976
4 participants (96% of the original cohort), 2067 participants were well-nourished (70%)
5 and 869 participants (30%) were malnourished.

6 Table 1 summarises malnutrition coding results for the cohort (n= 2976). A significantly
7 small number of malnourished participants were coded for malnutrition (n= 162, 19%;
8 p< 0.001). Three percent of the participants (n= 69) who were assessed as well-
9 nourished were coded for malnutrition.

10 When results were analysed for each of the 56 participating hospitals, malnutrition
11 codes were allocated to:

- 12 • all the malnourished patients in 1 hospital;
- 13 • none of the malnourished patients in 21 hospitals (malnutrition prevalence:
14 average 28% (range: 5 – 50%));
- 15 • 33% – 95% of the malnourished patients in the remaining 34 hospitals
16 (malnutrition prevalence: average 29% (range: 11 – 53%).

17

18

19 **DISCUSSION**

20 The objective of this paper was to report if malnutrition codes had been assigned to
21 patients who were identified as malnourished in the ANCDs cohort.⁴ According to the
22 findings, a significantly lower number of malnourished patients were coded. In fact,
23 malnourished participants from 21 hospitals were not coded for malnutrition. This may
24 have potentially translated to lower malnutrition-related reimbursements for the

1 hospitals. It is likely that the inclusion of malnutrition as a comorbidity may not have
2 increased the DRG-classification (and associated reimbursement) for patients who had
3 complex comorbidities already. However, correct assignment of malnutrition codes is
4 still warranted to reflect the burden of malnutrition in Australian and New Zealand
5 hospitals.

6

7 Our observation regarding poor malnutrition coding in hospitals is consistent with the
8 findings from the three previous Australian studies.^{11,19,23} Researchers in all three
9 studies retrospectively reviewed documentation and subsequent coding for malnutrition
10 (defined using SGA). When malnutrition coding was warranted but not included, the
11 codes were allocated hypothetically to derive an estimate of the financial shortfall
12 experienced by the hospitals. It appears that the inclusion of malnutrition changes the
13 DRG for approximately 20% of patients, which is equivalent to approximately
14 AU\$3500 of hypothetical reimbursement for each patient (Appendix I). Therefore, in
15 the ANCDs, the estimated collective average loss of reimbursement may have been
16 AU\$603780 (range: AU\$480240 – AU\$727320) (Appendix I).

17

18 When Ferguson et al (1997) and Lazarus and Hamlyn (2005) conducted their studies,
19 malnutrition was acceptable for DRG coding only if a medical practitioner documented
20 it in patients' medical charts.^{11,19} In 2008, the National Centre for Classification in
21 Health agreed that malnutrition may be coded when it is documented by a dietitian in
22 the medical record.^{29,30} This change presented dietitians with the opportunity to become
23 leaders in establishing concrete pathways for the identification and documentation of
24 malnutrition in hospital patients, and to collaborate with medical coders to ensure

1 malnutrition is correctly coded. Given the multicentre nature of this study, it was not
2 possible to identify whether malnutrition coding did not occur as a result of poor
3 documentation and/or due to the absence of a structured process for identification,
4 documentation, and coding for malnutrition. It has also been suggested that dietitians
5 may lack self-confidence in making a malnutrition diagnosis, which may result in
6 inadequate malnutrition coding.²³ There is emerging evidence to indicate that when
7 malnutrition is coded, the associated reimbursement also improves the profile of
8 dietitians amongst other healthcare members but can also result in increased funding for
9 employing more dietitians within the facility.^{31,32}

10

11 An interesting finding was that patients who had been previously identified as well-
12 nourished in the ANCDs were prospectively coded for malnutrition. A closer look at
13 the LOS data revealed that these patients had significantly longer ($p < 0.001$) median
14 LOS (23 days, range: 3 – 395 days) compared to patients who were identified as well-
15 nourished (10 days, range: 2 – 158 days) and malnourished (15 days, range: 2 – 119
16 days) in the ANCDs.⁵ Deterioration of nutritional status during hospitalisation has
17 previously been reported.^{2,33,34} Since these patients had a significantly extended LOS
18 compared with the rest of the cohort, it is possible that their nutritional status
19 deteriorated during hospitalisation, which reiterates the importance of regularly
20 reviewing patients' nutritional status during hospitalisation.

21

22 One limitation of this study is that malnutrition coding results may not be conclusive
23 due to the missing data. However, malnutrition coding data was missing for fewer than
24 five percent of the cohort and this is still the largest study to provide a snapshot of

1 malnutrition coding in Australian and New Zealand hospitals. It is also likely that the
2 screening process may have missed some malnourished patients. However, this reflects
3 real world practice. It was beyond the scope of this study to allocate malnutrition codes
4 hypothetically to estimate the potential financial shortfall experienced by participating
5 hospitals. Based on the three previous Australian studies,^{11,19,23} we have attempted to
6 provide a conservative estimate of the potential loss in reimbursement (Appendix I).

7

8 It is noteworthy that participating hospitals represent 20% of Australian acute care
9 hospitals³⁵ and 38% of acute care hospitals in New Zealand³⁶ (with >60 beds). Even
10 though malnutrition coding practices are not reflected for a majority of Australian and
11 New Zealand hospitals, this study provides insight into malnutrition coding practices in
12 public and private hospitals in this region.

13

14 In conclusion, the ANCDs has identified that malnutrition continues to be a common
15 problem in Australian and New Zealand acute care.⁴ This paper demonstrates gaps in
16 processes related to the documentation and subsequent coding for malnutrition, which
17 may result in financial loss of reimbursement for hospitals. This study highlights the
18 need for further research to identify barriers and enablers for malnutrition
19 documentation and coding. In the current stringent financial climate and rising
20 healthcare costs, a structured process for the identification, documentation and coding
21 for malnutrition will ensure appropriate casemix-related allocations for hospitals.³⁷ We
22 suggest that managers of dietetics departments, dietitians, and medical coders
23 collaborate to identify and address problems related to malnutrition documentation
24 and/or coding.

1 **ACKNOWLEDGEMENTS: [Removed for blind peer review]**

2

3 **Conflict of interest:** None to declare.

4

5 **REFERENCES**

- 6 1. White JV, Guenter P, Jensen G, et al. Consensus Statement: Academy of
7 Nutrition and Dietetics and American Society for Parenteral and Enteral
8 Nutrition. *JPEN JPparenter Enteral Nutr.* 2012 May 1, 2012;36:275-83.
- 9 2. Norman K, Pichard C, Lochs H, Pirlich M. Prognostic impact of disease-related
10 malnutrition. *Clin Nutr.* 2008;27:5-15.
- 11 3. Kubrak C, Jensen L. Malnutrition in acute care patients: A narrative review. *Int J*
12 *Nurs Stud.* 2007;44:1036-54.
- 13 4. Agarwal E, Ferguson M, Banks M, Bauer J, Capra S, Isenring E. Nutritional
14 status and dietary intake of acute care patients: Results from the Nutrition Care
15 Day Survey 2010. *Clin Nutr.* 2012;31:41-7. .
- 16 5. Agarwal E, Ferguson M, Banks M, et al. Malnutrition and decreased food intake
17 are associated with prolonged length of hospital stay, frequent hospital
18 readmissions, and greater in-hospital mortality: Results from the Nutrition Care
19 Day Survey 2010. *Clin Nutr.* 2013; 32: 737-745.
- 20 6. Chima CB, K, Dewitt M, Maeda M, Teran J, Mullen K. Relationship of
21 nutritional status to length of stay, hospital costs, and discharge status of patients
22 hospitalized in he medicine service. *J Am Diet Assoc.* 1997;97:975-8.
- 23 7. Correia M, Waitzberg D. The impact of malnutrition on morbidity, mortality,
24 length of hospital stay and costs evaluated through multivariate model analysis.
25 *Clin Nutr.* 2003;22:235-9. .
- 26 8. Lim S, Ong K, Chan Y, Loke W, Ferguson M, Daniels L. Malnutrition and its
27 impact on cost of hospitalisation, length of stay, readmission, and 3-year
28 mortality. *Clin Nutr.* 2012;31:345-50.
- 29 9. Marco J, Barba R, Zapatero A, et al. Prevalence of the notification of
30 malnutrition in the departments of internal medicine and its prognostic
31 implications. *Clin Nutr.* 2011;30:450-4.
- 32 10. Agarwal E, Ferguson M, Banks M, et al. Nutrition care practices in hospital
33 wards: Results from the Nutrition Care Day Survey 2010. *Clin Nutr.*
34 2012;31:995-1001.
- 35 11. Lazarus C, Hamlyn J. Prevalence and documentation of malnutrition in
36 hospitals: A case study in a large private hospital setting. *Nutr Diet.* 2005;62:41-
37 7.
- 38 12. Middleton MH, Nazarenko G, Nivison-Smith I, Smerdely P. Prevalence of
39 malnutrition and 12-month incidence of mortality in two Sydney teaching
40 hospitals. *Intern Med J.* 2001;31:455-61. .
- 41 13. Ferguson M, Banks M, Bauer J, Isenring E, Vivanti A, Capra S. Nutrition
42 screening practices in Australian healthcare facilities: a decade later. *Nutr Diet.*
43 2010;67:213-8.

- 1 14. Ferguson M, Capra S. Nutrition screening practices in Australian hospitals. *Nutr*
2 *Diet*. 1998;55:157-61.
- 3 15. Watterson C, Fraser A, Banks M, et al. Evidence based guidelines for nutritional
4 management of malnutrition in adult patients across the continuum of care. *Nutr*
5 *Diet*. 2009;66:s1-s34.
- 6 16. Writing Group of the Nutrition Care Process/Standardized Language
7 Committee. Nutrition Care Process and Model Part I: The 2008 Update. *J Am*
8 *Diet Assoc*. 2008;108:113-7.
- 9 17. Department of Health and Ageing: Australian Government. Australian Casemix
10 Glossary. AR-DRG Version 6.0. Accessed March 14, 2012 at
11 [http://www.health.gov.au/internet/main/publishing.nsf/Content/health-casemix-
13 glossary1.htm](http://www.health.gov.au/internet/main/publishing.nsf/Content/health-casemix-
12 glossary1.htm) .
- 13 18. Ravens J. A beginner's guide to casemix and its uses. *Nutr Diet*. 1997;54:100-1.
- 14 19. Ferguson M, Capra S. Coding for malnutrition enhances reimbursement under
15 casemix-based funding. *Nutr Diet*. 1997;54:102-7.
- 16 20. Victorian Government Department of Human Services. Victoria—public
17 hospitals and mental health services: Policy and funding guidelines 2008–09.
18 Accessed December 30, 2012 at <http://www.health.vic.gov.au/pfg0809>.
- 19 21. Ockenga J, Freudenreich M, Zakonsky R, Norman K, Pirlich M, Lochs H.
20 Nutritional assessment and management in hospitalised patients: Implication for
21 DRG-based reimbursement and health care quality. *Clin Nutr*. 2005;24:913-9.
- 22 22. Cheng P, Gilchrist A, Robinson K, Paul L. The risk and consequences of clinical
23 miscoding due to inadequate medical documentation: a case study of the impact
24 on health services funding. *HIM J*. 2009;38:35-46.
- 25 23. Gout B, Barker L, Crowe T. Malnutrition identification, diagnosis and dietetic
26 referrals: Are we doing a good enough job? *Nutr Diet*. 2009;66:206-11.
- 27 24. Rowell D, Jackson T. Additional costs of inpatient malnutrition, Victoria,
28 Australia, 2003-2004. *Eur J Health Econ*. 2011;12:353-61.
- 29 25. Ferguson M, Capra S, Bauer J, Banks M. Development of a Valid and Reliable
30 Malnutrition Screening Tool for Adult Acute Hospital Patients. *Nutrition*.
31 1999;15:458-64. .
- 32 26. Detsky A, McLaughlin J, Baker J. What is Subjective Global Assessment of
33 nutritional status. *JPEN J Parenter Enteral Nutr*. 1987;11:8-13.
- 34 27. National Centre for Classification in Health. The international statistical
35 classification of diseases and related health problems, 10th revision, Australian
36 modification (ICD-10-AM): New South Wales, Australia: National Centre for
37 Classification in Health, Faculty of Health Sciences, The University of Sydney;
38 2010.
- 39 28. Jackson T, Duckett S, Shepherd J, Baxter K. Measurement of adverse events
40 using ‘incidence flagged’ diagnosis codes. *J Health Serv Res Policy*
41 2006;11:21–6.
- 42 29. Department of Health and Ageing: Australian Government. AR-DRG Version
43 60 Accessed March 14, 2012 at
44 [http://www.health.gov.au/internet/main/publishing.nsf/Content/AR-DRG-
46 Version_60](http://www.health.gov.au/internet/main/publishing.nsf/Content/AR-DRG-
45 Version_60).
- 47 30. Cawood A, M E, Stratton R. Systematic review and meta-analysis of the effects
of high protein oral nutritional supplements. *Ageing Res Rev*. 2012;11:278-96.

- 1 31. Cobb B, Hardin E, Click A, Watson, L. Customising malnutrition
2 documentation accelerates hospital revenue and RD value. *J Am Diet Assoc.*
3 2011; 111: A77.
- 4 32. Hamilton M, Bullard S, Fugett K, Heath J, Jackson T, Jakobe D. Increasing the
5 value of the RD through Malnutrition DRG Coding. *J Am Diet Assoc.* 2010;
6 110: A83.
- 7 33. Kyle U, Genton L, Pichard C. Hospital length of stay and nutritional status. *Curr*
8 *Opin Clin Nutr Metab Care.* 2005;8:397-402.
- 9 34. Bauer JD, Hiscocks K, Fichera R, et al. Nutritional status of long-term patients
10 in the acute care setting. *Intern Med J.* 2012;42:1251-4.
- 11 35. Australian Institute of Health and Welfare. Australian hospital statistics 2009–
12 10. Health services series no. 40. Cat. no. HSE 107. Canberra: AIHW. Accessed
13 April 17, 2012 at <http://www.aihw.gov.au>. 2011.
- 14 36. New Zealand Ministry of Health – Manatū Hauora. Certified providers of
15 hospital and aged residential care services. Accessed April 17, 2012 at
16 <http://cert.moh.govt.nz/certification/review.nsf/default?OpenForm>.
- 17 37. Skipper A. Agreement on Defining Malnutrition. *JPEN J Parenter Enteral Nutr.*
18 2012 May 1, 2012;36:261-2.

1 **Appendix I: An estimate of the financial loss of reimbursement related to malnutrition coding**

Authors, Year	Number of malnourished patients	Number of patients for whom DRG was changed due to malnutrition coding	Percentage of patients for whom DRG changed due to malnutrition coding	Hypothetical reimbursement per patient
Lazarus et al, 2005¹¹	137	30	22	AU\$4180
Gout et al, 2009²³	53	10	19	AU\$2760
Average:			20%	AU\$3470

2 Notes: Since malnutrition coding for approximately 20% of the patients in the above studies led to an increase in financial reimbursement,

3 in the ANCDs (2010):

4 Number of malnourished patients: 869

5 20% of malnourished patients= 174

6 Estimated total average loss in reimbursement: 174 * \$3470= AU\$603780

7 Estimated total minimum loss in reimbursement: 174 * \$2760= AU\$480240

8 Estimated total maximum loss in reimbursement: 174 *\$4180= AU\$727320

1 **Table 1: Malnutrition coding results (n= 2976)**

	Well-nourished^a (n= 2067, 70%)		Malnourished^b (n= 869, 30%)		p-value^c
	Coding not required	Coded for malnutrition^c	Not coded for malnutrition	Coded for malnutrition	
Participants (n (%))	1998 (97%)	69 (3%)	707 (81%)	162 (19%)	<0.001

2 ^a Well-nourished participants: included those “not at risk” of malnutrition (according to the MST²⁵) and SGA-A²⁶.

3 ^b Malnourished participants: included those with BMI <18.5 kg/m² ²⁷, moderately malnourished (SGA-B)²⁶, or severely malnourished
4 (SGA-C)²⁶.

5 ^c Chi-square test