

Bond University  
Research Repository



## Malnutrition in the elderly

### A narrative review

Agarwal, E; Miller, M; Yaxley, A; Isenring, E

*Published in:*  
Maturitas

*DOI:*  
[10.1016/j.maturitas.2013.07.013](https://doi.org/10.1016/j.maturitas.2013.07.013)

*Licence:*  
CC BY-NC-ND

[Link to output in Bond University research repository.](#)

#### *Recommended citation(APA):*

Agarwal, E., Miller, M., Yaxley, A., & Isenring, E. (2013). Malnutrition in the elderly: A narrative review. *Maturitas*, 76(4), 296-302. <https://doi.org/10.1016/j.maturitas.2013.07.013>

#### **General rights**

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

For more information, or if you believe that this document breaches copyright, please contact the Bond University research repository coordinator.

## **Malnutrition in the elderly: A narrative review**

Agarwal E<sup>1</sup>, Miller M<sup>2</sup>, Yaxley A<sup>2</sup>, Isenring E<sup>3, 4</sup>

<sup>1</sup>School of Exercise and Nutrition Science, Queensland University of Technology, Victoria Park Road, Kelvin Grove, Queensland 4059, Australia

<sup>2</sup>Flinders University, Adelaide, Australia

<sup>3</sup>University of Queensland, Brisbane, Australia

<sup>4</sup>Princess Alexandra Hospital, Brisbane, Australia

### **Abstract**

The focus of nutrition is often on healthy diets and exercise to minimise the risk of developing lifestyle diseases such as cancer, diabetes and cardiovascular disease. However, during the shift into older years often the nutrition priorities change towards meeting increased nutrient needs with less energy requirements and minimising lean muscle loss. There are several causes of general malnutrition in the elderly that lead to depletion of muscle including starvation (protein-energy malnutrition), sarcopenia and cachexia. The prevalence of protein-energy malnutrition increases with age and the number of comorbidities. A range of simple and validated screening tools can be used to identify malnutrition in older adults e.g. MST, MNA-SF and 'MUST'. Older adults should be screened for nutritional issues at diagnosis, on admission to hospitals or care homes and during follow up at outpatient or General Practitioner clinics, at regular intervals depending on clinical status. Early identification and treatment of nutrition problems can lead to improved outcomes and better quality of life.

**Keywords:** protein-energy malnutrition, elderly, sarcopenia, cachexia, prevalence, nutritional management

### **Corresponding Author:**

Dr Elizabeth Isenring

Address

Email

Telephone

**Conflict of interest:** None to declare

## **1. Introduction**

The world's population is rapidly ageing with estimates that in the first five decades of the 21<sup>st</sup> century the proportion of the world's population over 60 years will double from 11% to 22%. The expected increase in the absolute number of older adults will triple from 605 million to 2 billion over this period [1]. As the number of older people continues to rise, provision of improved healthcare to the elderly- both in hospital and in the community- is imperative. Often, the focus of nutrition in older adults is a healthy diet and exercise to minimise the risk of developing lifestyle diseases (such as cardiovascular disease, Type 2 diabetes mellitus). However, there is a large body of evidence to indicate that protein-energy malnutrition (PEM) is a common problem in this age-group, including in the hospital, nursing home and community setting. Therefore, the purpose of this paper is to summarise the current literature regarding:

- the prevalence
- aetiology
- identification and
- effective nutritional management

of PEM in the elderly.

## **2. Causes of general malnutrition- starvation (PEM), Sarcopenia, and Cachexia**

While there is no universally accepted definition of malnutrition, one of the most commonly used definitions identifies malnutrition as “a state of nutrition in which a deficiency, or excess, of energy, protein and micronutrients causes measurable adverse effects on tissue/body form (body shape, size and composition) and function, and clinical outcome” [2]. However, in relation to under-nutrition, this definition does not take into account the aetiology of unintentional weight loss. Recent literature suggests that unintentional weight loss is comprised of three primary syndromes: starvation, sarcopenia and cachexia [3,4]. Furthermore, there is a level of complexity involved in that the unintentional weight loss may be a result of any two or three of those syndromes in combination [5]. The term malnutrition dominates the literature around unintentional weight loss and is likely to capture all unintentional weight loss as if it were one condition. Disentangling the primary aetiology is critical for implementation of appropriate nutrition support as responsiveness to dietary

modifications differ. Even now there is uncertainty and confusion amongst dietitians, which is likely to be reflected amongst all clinicians [6].

Starvation is generally accepted to occur purely as a result of protein-energy deficiency and is synonymous with PEM [4, 7]. The major factor that distinguishes starvation from other syndromes of unintentional weight loss is that it is reversed when adequate energy and protein intake is achieved (discussed further in Section 7) [7]. There are numerous nutrition screening tools to detect protein-energy PEM (discussed further in Section 5) and these are increasingly becoming mandatory across the continuum of care but primarily in the acute care setting, however none of these distinguishes starvation from any other syndrome of unintentional weight loss.

Results of recent attempts to provide, and agree upon, definitions and diagnostic criteria for sarcopenia indicate that it is a progressive loss of muscle mass that occurs with normal ageing although this area is still under investigation [8-10]. It is known to be associated with increased frailty, loss of strength, reduced physical function and diminished capacity for exercise, as a result of decreased muscle mass and alterations to muscle structure at the microscopic level which change the function of muscle in sarcopenia [11]. It is likely that effective interventions for the treatment of sarcopenia should be multi-disciplinary. Dietary management should provide adequate energy and protein intake however this alone would be unlikely to address weight loss as sarcopenia is thought to occur regardless of energy balance [12, 13]. Recent evidence indicates that the most effective intervention thus far is combination therapy comprising nutrition and resistance training exercises [14]. There are currently no screening tools for the detection of sarcopenia, and diagnosis is usually based on clinical judgement, although criteria have been proposed by at least one of the recent consensus papers [8]. Health professionals working with older adults with unintentional weight loss should be mindful that nutrition alone may not improve their condition.

Cachexia has long been associated with a number of chronic conditions such as cancers, HIV/AIDS heart failure and Chronic Obstructive Pulmonary Disease (COPD). In 2008 a group of prominent researchers in the field came together to develop a consensus definition for cachexia which indicates that “cachexia is a complex metabolic syndrome associated with underlying illness and characterised by loss of muscle with or without loss of fat mass [15]. Furthermore, an expert group used the Delphi technique to define diagnostic criterion for cancer cachexia as weight loss greater than 5%, or weight loss greater than 2% in individuals already showing depletion according to current bodyweight and height (body-mass index,

BMI;  $<20 \text{ kg/m}^2$ ) or skeletal muscle mass (sarcopenia) [16]. An agreement was made that the cachexia syndrome can develop progressively through various stages--precachexia to cachexia to refractory cachexia. Assessment for classification and clinical management should include: anorexia or reduced food intake, catabolic drive, muscle mass and strength, functional and psychosocial impairment [16]. Although there is little research into the condition there is evidence that geriatric cachexia also manifests in the elderly as a result of ageing [17,18]. Evans et al (2008) were also clear in identifying cachexia as a separate syndrome from starvation and sarcopenia [15]. There has been a considerable amount of research into interventions for the treatment of cachexia although the majority has been conducted in individuals with end-stage chronic disease therefore it may be prudent to treat results with caution. In addition to appropriate dietary intake, the most promising therapies appear to be appetite stimulants, combination pharmacological treatments and exercise interventions although studies in this area are prone to high levels of drop-out and side effects which may not be due to the interventions under investigation [19].

### **3. Prevalence of PEM in the elderly:**

Multicentre studies that have evaluated PEM prevalence in the acute care setting report that 23-60% of elderly patients are malnourished and an estimated 22-28% are at nutritional risk (Table 1).

In comparison to other healthcare settings, there is limited literature on the prevalence of PEM in community-dwelling older adults. However, the reported prevalence indicates a range of 5-30% [27-29]. In the residential aged care setting, the reported PEM prevalence ranges from 16-70% depending on the assessment tool used and the level of care required [30-33]. In general it is known that the prevalence of PEM increases as the level of care increases.

### **4. Risk factors of PEM in the elderly**

The aetiology of protein-energy PEM in the elderly is multifactorial and consists of physiological, social and economic parameters, often referred to as the “nine d’s” (dementia, dysgeusia, dysphagia, diarrhoea, depression, disease, poor dentition, dysfunction, and drugs) [34]. The physiological parameters also affect food consumption in the elderly (Figure 1); thereby further exacerbating the problem of PEM.

## 5. Identifying PEM:

Malnutrition screening is the recommended first step in the nutrition care process as it allows the early identification of nutritional concerns [35-37]. Given the multifactorial nature of PEM in the elderly, and in the absence of a single objective measure or “gold standard”, a number of nutrition screening tools specific to the older adult population have been developed. Each tool is unique as they include a range of different parameters [38]:

- Biochemical and clinical indices: Nutritional Risk Index (NRI) [39] and the Geriatric Nutritional Risk Index (GNRI) [40];
- Anthropometry, mobility, cognitive state, self-perceived health and nutrition: Mini Nutritional Assessment Screening Form (MNA-SF) [41], Malnutrition Universal Screening Tool (MUST) [42]; and unintentional weight loss and poor intake (MST);
- Medical history, clinical and subjective evaluation: Nutrition Risk Screening 2002 (NRS-2002) [25].

Tools such as the MNA-SF, MUST, and NRS-2002 incorporate body mass index (BMI) for detecting the risk of PEM. BMI is a simple index of weight-for-height (defined as weight in kilograms divide by the square of height in meters ( $\text{kg}/\text{m}^2$ ) [43]. Individuals are classified as underweight ( $\text{BMI} < 18.5 \text{ kg}/\text{m}^2$ ), normal weight ( $18.5\text{-}24.9 \text{ kg}/\text{m}^2$ ), overweight ( $25\text{-}29.9 \text{ kg}/\text{m}^2$ ) or obese ( $\geq 30 \text{ kg}/\text{m}^2$ ) [43]. However, nutrition screening tools such as the MNA-SF [41] support the use of higher cut-off points to identify PEM in older adults. The rationale for higher BMI cut-off points is that older adults are likely to [44]:

- have a smaller proportion of lean body mass than younger adults (as a result of aging, PEM, or inactivity);
- shorten with age due to the compression associated with an osteoporotic spine (kyphosis).

Research also indicates that moderately higher BMI is protective against mortality [45]. A recent study investigating the relationship between BMI and selected clinical outcomes found that after adjustment for age, sex and level of care,  $\text{BMI} < 22 \text{ kg}/\text{m}^2$  increased the risk of fracture by 38% and mortality by 52% in a sample of 1846 older adults in residential aged care [46].

The Subjective Global Assessment (SGA) [26] and Mini-Nutritional Assessment [47] are validated nutrition assessment methods that use a range of parameters that can be used to make a nutritional diagnosis and initiate nutritional management in elderly patients. In contrast to nutrition screening tools, nutritional assessment methods are much more

comprehensive and are used by trained professionals (such as dietitians, physicians, nurses or research assistants).

A recent study compared the accuracy of identifying malnutrition by seven validated nutrition screening tools against SGA and MNA in their accuracy of identifying malnutrition in an elderly cohort (N=134) [48]. With the exception of the Rapid Screen, all other tools performed with a high sensitivity and specificity when compared with the SGA [48]. The MST and NRS-2002 had the highest accuracy when compared with the SGA [48]. Although MNA-SF had a high sensitivity with MNA, it demonstrated poor specificity and positive predictive value when compared with the SGA [48]. The MNA-SF was developed to identify patients requiring comprehensive assessment using the MNA, and therefore the inherent differences between MNA and SGA could have explained this result [48].

Given that there are a range of validated nutrition screening tools for the elderly population, it is advised that tools are selected on the basis of who will perform the nutrition screening and the resources available within the healthcare setting. Some screening tools contain anthropometric measures and calculations and take more time to complete. Involuntary weight loss is easy to overlook in someone who is overweight and it is important to recognise that someone may be malnourished even though they appear healthy or overweight.

## **6. Consequences of PEM**

PEM can affect almost every function, organ and/or system of the human body [49] and therefore has been associated with a range of outcomes with implications for health and recovery from illness/surgery [50]. Although the onset of PEM depends on the body's nutritional reserves (independent of the disease state) [50], a connection between PEM and disease exists, whereby disease may influence PEM, or PEM may have a negative effect on disease [49]. Therefore, PEM is often referred to as “a cause and consequence of adverse outcomes” (Figure 2). Older patients are known to be especially vulnerable to PEM-related consequences [51] such as prolonged length of stay in hospital [29,52], increase risk of falls [29], admission to higher level care [52], decreased physical function [52], poorer quality of life [52], increased risk of life-threatening complications [53] and increased mortality [54].

The Australasian Nutrition Care Day Survey is the largest study in the Australian and New Zealand acute care setting to evaluate nutritional status and its association with health-related outcomes [20, 56]. The study recruited 3122 acute care patients from 56 hospitals across the two countries [20]. More than 50% of the cohort in this study (n=1650, 56%) were aged  $\geq 65$  years [20]. Participants were screened for nutritional risk using the PEM screening tool (MST) [57]. Patients identified with nutritional risk were then comprehensively assessed for PEM using Subjective Global Assessment (SGA) [26]. Health-related outcomes data (namely LOS, readmissions, and in-hospital mortality) were collated 90 days after nutritional assessment [56]. Results (unpublished) from the survey indicated that elderly malnourished patients ( $\geq 65$  years) had significantly longer LOS, readmissions, and 90-day in-hospital mortality in comparison to their well-nourished counterparts and those aged  $<65$  years (Table 2) [58].

A retrospective study aimed to evaluate if PEM at hospital admission predicted clinical outcomes (LOS, readmissions, mortality, change in level of care at discharge) at 18-months follow-up in 2076 geriatric patients (aged  $\geq 65$  years) recruited from two sub-acute hospitals in Australia [59]. Nutritional status was determined using the MNA within 72-hours of admission [59]. The study found that 30% of the patients were malnourished at admission and 53% were at risk of malnutrition [59]. Malnourished patients had significantly higher median LOS (34 days, range:21-58) in comparison to patients at nutritional risk (26 days, range: 15-41 days) or well-nourished (20 days, range: 14-26 days) ( $p<0.001$ ) [59]. Although number of readmissions was not significantly associated with PEM, malnourished patients demonstrated a significantly high rate of discharge to high level residential aged care facilities (33% versus 17% in patients with PEM risk and 5% in well-nourished patients,  $p\leq 0.001$ ) [59]. Survival analysis (controlling for age, gender, major disease classification, morbidity, and LOS at index admission) indicated that the hazard ratio of death in the malnourished group was 3.4 times (Confidence Interval (CI): 1.07-10.87,  $p<0.05$ ) the well-nourished group [59].

## **7. Management of PEM**

In addition to the pathophysiological, social and behavioural factors that play a role in the development of PEM amongst older adults, illness and hospitalisation have also been implicated in the development (or worsening of PEM) in elderly patients. A large number of barriers adversely affecting food intake in elderly patients have been identified in the



literature. These include self-limiting factors (loss of appetite, acute illness, oral issues, low mood, dysphagia, confusion, isolation), catering limitations (inflexible mealtimes, difficulty accessing food and beverage packaging, lack of menu variety, unappealing meals), and organisational barriers (interruptions during mealtimes, inadequate feeding assistance, unpleasant smells, disruptive behaviour from staff members and other patients) [60].

Weekes et al (2009) evaluated the efficacy of different nutritional care interventions (including nutrition screening and nutritional support, excluding artificial nutritional support like parenteral nutrition) aiming to improve nutritional or clinical outcomes or costs by reviewing 297 associated publications [61]. The review found a scarcity of unbiased, large randomised controlled trials in the published literature and highlighted that not enough research had been done to demonstrate the nutritional- and cost-effectiveness of interventions [61]. Given that oral nutritional supplements are commonly prescribed to alleviate PEM, another systematic review by Milne and colleagues assessed 55 randomised trials (n=9187) for the clinical and nutritional outcomes for older people offered supplements in different settings [62]. Although the review found that supplements could improve the nutritional status of older people, lead to a small gain in weight and muscle mass, the authors identified that elderly people may have difficulty accepting supplements as a result of reported gastrointestinal disturbance [62]. Milne et al also acknowledged that the results of their review were limited by inadequate studies, which used mainly poor methodologies, lacked sufficient statistical power and duration of follow-up to demonstrate any beneficial effects [62].

However, there is a growing body of evidence to demonstrate merit in offering nutritional support to patients following hospital discharge:

- A Cochrane review by Shepperd and colleagues indicates that structured discharge plans that are tailored for individual patients are associated with reductions in LOS and readmission rates in older people [63].
- In a randomised controlled trial, 259 elderly acute care patients who were at nutritional risk were randomised into two groups- the intervention group received individualised nutritional care from a dietitian and three post-discharge home visits; participants in the control group received either one dietitian visit during hospitalisation or standard care (nil dietitian review) [64]. Outcomes measured at baseline and at six months included nutritional status (MNA), mortality, health status,

biochemistry, cognitive, emotional and functional parameters [64]. At six months, researchers noticed lower mortality and moderate improvement in nutritional status in the intervention group [64].

- Two recent studies, involving over 200 participants each, have demonstrated that nutrition interventions (including enriched diets and/or oral nutritional supplements, home visits and/or telephone follow-ups) in elderly patients (aged 65 years and over) yielded improvements in nutritional status, functional status and mortality rates [65].
- A recent systematic review and meta-analysis of randomised controlled trials evaluating the use of ONS in medical and surgical patients (aged over 65 years) has also established positive associations with dietary intake and weight status [66].
- In an Australian prospective cohort pilot study an inter-disciplinary discharge team (consisting of a specialist discharge planning nurse and dietitian) provided 12 elderly participants (aged over 65 years) with nutrition support (education, advice, service coordination, home visits and telephone reviews) for six weeks following hospital discharge [67]. Participants found the proposed model of care acceptable and demonstrated improved nutritional status at 12 weeks post-discharge [67].
- In a Danish study 152 geriatric medical patients (aged over 65 years) who were at nutritional risk were randomised into two groups: one group was offered a 12-week individualised dietary counselling with a dietitian along with follow-up by a general practitioner; and the other group was offered a 12-week follow-up by the general practitioner only [68]. Outcome measures were functional status (hand-grip strength, chair stand, mobility, disability, tiredness in ADLs, rehabilitation capacity) and nutritional status (weight, BMI, energy and protein intake). Participants in the group receiving individualised dietary counselling had a positive effect on both outcomes measures [68]. This study is unique for demonstrating the effectiveness of dietary counselling in a general medical setting and perhaps the study design can be used to conduct larger randomised controlled trials [68].

These studies demonstrate positive outcomes associated with offering individualised nutritional support to malnourished elderly people- both during hospitalisation and post-discharge. These studies also highlight the importance of routine nutrition screening to monitor PEM (risk) in order to identify, diagnose and treat PEM in a timely manner.

The use of high energy and protein diets and oral nutritional supplements have demonstrated improved nutritional, clinical and functional outcomes and therefore should be provided to patients not meeting their dietary requirements during hospitalisation [35, 63, 69-77]. Pharmacologic treatment of some vitamin and/ mineral deficiencies may also be indicated clinically. In particular low levels of vitamin D, B<sub>12</sub> and/or Iron can occur in those with PEM. Low serum albumin levels are not necessarily indicative of PEM as this could reflect inflammation or disease state. Albumin and transferrin levels may be normal but glucose and cholesterol may be low in undernourished adults.

Food services are also increasingly using systems-level strategies designed to maximise nutritional intake, such as feeding assistance, red trays to identify those at risk of PEM and therefore in need of assistance, and protected mealtimes [78]. Thus far however, there is little evidence that these systems-level interventions result in improvements in intake or patient outcomes with a recent Australian review in the area identifying only seven studies investigating this type of strategy [78]. Only two of the seven included studies identified clinically meaningful improvements in energy intake and two studies demonstrating clinically meaningful improvements in protein intake [78]. Despite these increases in energy and protein intake there was no report of significant difference in weight, length of stay or mortality, between those in the intervention group and those in the control group in any included study [78].

## **8. Implications for practice**

Validated nutrition screening tools and monitoring weight (and body composition) are important to identify older adults at nutritional risk. Geriatricians should continue to identify and manage nutritional issues as part of a comprehensive geriatric assessment. General practitioners often have the primary role of monitoring the nutritional status of patients who are discharged from acute or rehabilitation settings, the residential aged care setting and in the community. Strategies to identify and manage patients at PEM risk will avoid further decline in their patients' health status. Where available a Dietitian can provide a comprehensive nutrition assessment, treatment and follow-up plan for patients. Future research is required to validate proposed definitions and criteria of sarcopenia and cachexia. In the meantime, clinicians should use their professional judgement in identifying and managing these conditions. Regardless of the underlying causes of muscle depletion,

nutritional therapy should still play a key role as part of the multimodal treatment that may also include exercise and pharmaceutical interventions.

## 9. Conclusion

Malnutrition risk increases with age and level of care. Muscle wasting related to aging (sarcopenia) is more evident in advanced years and can be masked by fat mass. Cachexia is mediated by proinflammatory cytokines and occurs with particular conditions like cancer but may also occur with aging. The prevalence of PEM in older adults due to reduced dietary intake remains unacceptably high. Several valid malnutrition screening tools e.g. MST, MNA-SF and MUST are available to identify nutritional risk but further work regarding tools to distinguish between PEM, sarcopenia and cachexia are required. PEM is independently associated with poorer outcomes and decreased quality of life. For elderly with PEM or at nutritional risk there is evidence that oral nutritional supplements and dietary counselling can help increase dietary intake and improve quality of life. Further research into multimodal (e.g. nutritional, pharmaceutical, exercise) treatment of sarcopenia and cachexia are required.

## 10. References

- [1] WHO. Interesting facts about ageing. 2012 [cited 13th June 2013]; Available from: [www.who.int/ageing/about\\_facts\\_en\\_index.html.pdf](http://www.who.int/ageing/about_facts_en_index.html.pdf).
- [2] Stratton R, Green C, Elia M. Scientific criteria for defining malnutrition. Disease related malnutrition: an Evidence-Based approach to treatment. UK: CABI Publishing; 2003. p. 1-34.
- [3] Chapman I. Weight loss in older persons. Medical Clinics of North America. 2011;95(3):579-93.
- [4] Thomas D. Loss of skeletal muscle mass in aging: examining the relationship of starvation, sarcopenia and cachexia. Clinical Nutrition. 2007;26(4):389-99.
- [5] Yaxley A, Miller M, Fraser R, Cobiac L, Crotty M. The complexity of treating wasting in ambulatory rehabilitation: is it starvation, sarcopenia, cachexia or a combination of these conditions? Asia Pacific Journal of Clinical Nutrition. 2012;21(3):386-93.
- [6] Yaxley A, Miller M. The Challenge of Appropriate Identification and Treatment of Starvation, Sarcopenia, and Cachexia: A Survey of Australian Dietitians. Journal of Nutrition and Metabolism. 2011;Vol 2011(Article ID 603161).
- [7] Thomas D. Distinguishing starvation from cachexia. Clinics in Geriatric Medicine. 2002;18:883-91.
- [8] Cruz-Jentoft A, Baeyens J, Bauer J, et al. Sarcopenia: European consensus on definition and diagnosis Report of the European Working Group on Sarcopenia in Older People. Age and Ageing. 2010;39(4):412-23.
- [9] Fielding R, Vellas B, Evans W, et al. Sarcopenia: An Undiagnosed Condition in Older Adults. Current Consensus Definition: Prevalence, Etiology, and Consequences. International Working Group on Sarcopenia. Journal of the American Medical Directors Association. 2011;12:249-56.
- [10] Muscaritoli M, Anker SD, Argilés J, et al. Consensus definition of sarcopenia, cachexia and pre-cachexia: Joint document elaborated by Special Interest Groups

- (SIG) "cachexia-anorexia in chronic wasting diseases" and "nutrition in geriatrics". *Clinical Nutrition*. 2010;29(2):154-9.
- [11] Lang T, Streeper T, Cawthon P, Baldwin K, Taaffe D, Harris T. Sarcopenia: etiology, clinical consequences, intervention and assessment. *Osteoporosis International*. 2010;21:543-59.
- [12] Messinger-Rapport B, Thomas D, Gammack J, Morley J. Clinical Update on Nursing Home Medicine: 2009. *Journal of the American Medical Directors Association*. 2009;10:530-53.
- [13] Rolland Y, Dupuy C, Abellan van Kan G, Gillette S, Vellas B. Treatment strategies for sarcopenia and frailty. *Medical Clinics of North America*. 2011;95:427-38.
- [14] Rolland Y, Onder G, Morley J, Gillette S, Abellan van Kan G, Vellas B. Current and Future Pharmacologic Treatment of Sarcopenia. *Clinics in Geriatric Medicine*. 2011;27:423-47.
- [15] Evans W, Morley J, Argilés J, et al. Cachexia: a new definition. *Clinical Nutrition*. 2008;27:793-9.
- [16] Fearon K, Strasser F, Anker SD, et al. Definition and classification of cancer cachexia: an international consensus. *The Lancet Oncology*. 2011;12(5):489-95.
- [17] Kotler D. Cachexia. *Annals of Internal Medicine*. 2000;133(8):622-34.
- [18] Morley JE, Thomas DR, M-MG W. Cachexia: pathophysiology and clinical relevance. *American Journal of Clinical Nutrition*. 2006;83:735-43.
- [19] Yaxley A, Miller M, Fraser R, Cobiac L. Pharmacological interventions for geriatric cachexia: A narrative review of the literature. *Journal of Nutrition, Health and Aging*. 2012;16:148-54.
- [20] Agarwal E, Ferguson M, Banks M, Bauer J, Capra S, Isenring E. Nutritional status and dietary intake of acute care patients: Results from the Nutrition Care Day Survey 2010. *Clin Nutr*. 2012;31:41-7. doi: 10.1016/j.clnu.2011.08.002.
- [21] Imoberforf R, Meier R, Krebs P, et al. Prevalence of undernutrition on admission to Swiss hospitals. *Clinical Nutrition*. 2009;29(1):38-41.
- [22] Pirlich M, Schütz T, Norman K, et al. The German Hospital Malnutrition Study. *Clinical Nutrition*. 2006;25:563-72. doi: 10.1016/j.clnu.2006.03.005.
- [23] Correia MITD, Campos ACL. Prevalence of hospital malnutrition in Latin America: The multicenter ELAN study. *Nutrition*. 2003;19(10):823-5.
- [24] Waitzberg DL, Caiaffa WT, Correia MITD. Hospital malnutrition: the Brazilian national survey (IBRANUTRI): a study of 4000 patients. *Nutrition*. 2001;17(7-8):573-80.
- [25] Kondrup J, Rasmussen H, Hamberg O, Stanga Z, An Ad Hoc Espen Working Group. Nutritional Risk Screening (NRS 2002): a new method based on analysis of controlled clinical trials. *Clinical Nutrition*. 2003;22(3):321-36.
- [26] Detsky A, McLaughlin J, Baker J. What is Subjective Global Assessment of nutritional status. *JPEN- Journal of Parenteral and Enteral Nutrition* 1987;11:8-13. doi: 0.1177/014860718701100108
- [27] Leggo M, Banks M, Isenring E, Stewart L, Tweeddale M. A quality improvement nutrition screening and intervention program available to Home and Community Care eligible clients. *Nutrition & Dietetics*. 2008;65:162-7.
- [28] Cuervo M, Garcia A, Ansorena D, Sanchez-Villegas A, Martinez-Gonzalez M, Astiasaran I. Nutritional assessment interpretation in 22,007 Spanish community-dwelling elders through the Mini Nutritional Assessment test. *Public Health Nutrition*. 2009;12:82-90.
- [29] Visvanathan R, Macintosh C, Callary M, Penhall R, Horowitz M, I. C. The nutritional status of 250 older Australian recipients of domiciliary care services and its association with outcomes at 12 months. *Journal of the American Geriatrics Society*. 2003;51:1007-11.
- [30] Banks M, Ash S, Bauer J, Gaskill D. Prevalence of malnutrition in adults in Queensland public hospitals and residential aged care facilities. *Nutrition & Dietetics*. 2007;64:172-8. doi: 10.1111/j.747-0080.2007.00179.x.

- [31] Valentini L, Schindler K, Schlawfer R, et al. The first NutritionDay in nursing homes: Participation may improve malnutrition awareness. *Clinical Nutrition*. 2009;28:109-16.
- [32] Meijers JMM, Halfens RJG, van Bokhorst-de van der Schueren MAE, Dassen T, Schols JMGA. Malnutrition in Dutch health care: Prevalence, prevention, treatment, and quality indicators. *Nutrition*. 2009;25(5):512-9.
- [33] Miller M, Loh W, Watterson C, Fraser A, C N. Nutrition screening in long term care facilities: an evaluation of the methodological quality of available instruments. *CAB Reviews: Perspectives in Agriculture, Veterinary Science, Nutrition and Natural Resources*. 2008;3:17.
- [34] Edington J, Boorman J, Durrant E. Prevalence of malnutrition on admission to four hospitals in England. The Malnutrition Prevalence Group. *Clinical Nutrition*. 2000;19:191-5.
- [35] Watterson C, Fraser A, Banks M, et al. Evidence based guidelines for nutritional management of malnutrition in adult patients across the continuum of care. *Nutrition and Dietetics*. 2009;66:s1-s34.
- [36] Kondrup J, Allison S, Elia M, Vellas B, Plauth M. ESPEN Guidelines for Nutrition Screening 2002. *Clinical Nutrition*. 2003;22(4):415-21.
- [37] Mueller C, Compher C, Ellen DM, the American Society for Parenteral Enteral Nutrition Board of Directors. A.S.P.E.N. Clinical Guidelines: Nutrition Screening, Assessment, and Intervention in Adults. *JPEN- Journal of Parenteral and Enteral Nutrition* 2011 January 1, 2011;35(1):16-24.
- [38] Poulia K, Yannakoulia M, Karageorgou D, et al. Evaluation of the efficacy of six nutritional screening tools to predict malnutrition in the elderly. *Clinical Nutrition*. 2012;31:378-85.
- [39] Wolinsky F, Coe R, McIntosh W, et al. Progress in the development of a nutritional risk index. *The Journal of Nutrition*. 1990;120(11):1549-53.
- [40] Cereda E, Pedrolli C. The geriatric nutritional risk index. *Current Opinion in Clinical Nutrition and Metabolic Care*. 2009;12:1-7.
- [41] Rubenstein L, Harker J, Salva A, Guigoz Y, Vellas B. Screening for undernutrition in geriatric practice: Developing the Short-Form Mini Nutritional Assessment (MNA-SF). *Journal of Gerontology*. 2001;56A:M366-M72.
- [42] Malnutrition Advisory Group (MAG). The 'MUST' Explanatory Booklet. A Guide to the 'Malnutrition Universal Screening Tool' ('MUST') for Adults: A Standing Committee of the British Association for Parenteral and Enteral Nutrition (BAPEN); 2003.
- [43] World Health Organization. Global database on body mass index. Accessed 13 September 2011 at <http://apps.who.int/bmi/index.jsp?introPage=intro3.html> 2010.
- [44] Elia M, Stratton R. An analytical appraisal of nutrition screening tools supported by original data with particular reference to age. *Nutrition*. 2012;28:477-94.
- [45] Donini L, Savina C, Gennaro E, et al. A systematic review of the literature concerning the relationship between obesity and mortality in the elderly. *The Journal of Nutrition, Health and Aging*. 2012;16(1):89-98.
- [46] Miller M, Thomas J, Cameron I, et al. Body mass index: a simple, rapid and clinically meaningful index of under-nutrition in the oldest-old? . *Br J Nutr*. 2009;101:1300-5.
- [47] Guigoz Y, Vellas B, Garry P. Assessing the Nutritional Status of the Elderly: The Mini Nutritional Assessment as part of the Geriatric Evaluation. *Nutrition Reviews*. 1996;54:S59-S65.
- [48] Young AM, Kidston S, Banks MD, Mudge AM, Isenring EA. Malnutrition screening tools: Comparison against two validated nutrition assessment methods in older medical inpatients. *Nutrition*. 2013 1//;29(1):101-6.
- [49] Correia M, Waitzberg D. The impact of malnutrition on morbidity, mortality, length of hospital stay and costs evaluated through multivariate model analysis. *Clinical Nutrition*. 2003;22:235-9. doi: 10.1016/s0261-5614(02)00215-7.
- [50] Corish CA, Kennedy NP. Protein-energy undernutrition in hospital in-patients. *Br J Nutr*. [Review]. 2000 Jun;83(6):575-91.

- [51] Kubrak C, Jensen L. Malnutrition in acute care patients: A narrative review. *International Journal of Nursing Studies*. 2007;44:1036-54.
- [52] Neumann S, Miller M, Daniels L, Crotty C. Nutritional status and clinical outcomes of older patients in rehabilitation. *Journal of Human Nutrition & Dietetics*. 2005;18(2):129-36.
- [53] Sullivan D, Bopp M, Roberson P. Protein-energy undernutrition and life-threatening complications among the hospitalised elderly. *Journal of General Internal Medicine*. 2002;17:923-32.
- [54] Sullivan DH, Walls R. Protein-Energy Undernutrition and the Risk of Mortality Within Six Years of Hospital Discharge. *Journal of the American College of Nutrition*. 1998;17(6):571-8.
- [55] Jeejeebhoy K. Nutritional Assessment. *Nutrition*. 2000;16(7/8):585-90.
- [56] Agarwal E, Ferguson M, Banks M, et al. Malnutrition and decreased food intake are associated with prolonged length of hospital stay, frequent hospital readmissions, and greater in-hospital mortality: Results from the Nutrition Care Day Survey 2010. *Clin Nutr*. In Press.
- [57] Ferguson M, Capra S, Bauer J, Banks M. Development of a Valid and Reliable Malnutrition Screening Tool for Adult Acute Hospital Patients. *Nutrition*. 1999;15(6):458-64. doi: 10.1016/S0899-9007(99)00084-2.
- [58] Agarwal E. Health-related outcomes of well-nourished and malnourished acute care patients (Australasian Nutrition Care Day Survey). Unpublished raw data.
- [59] Charlton K, Nichols C, Bowden S, et al. Poor nutritional status of older subacute patients predicts clinical outcomes and mortality at 18 months of follow-up. *European Journal of Clinical Nutrition*. 2012;66(11):1224-8.
- [60] Mudge AM, Ross LJ, Young AM, Isenring EA, Banks MD. Helping understand nutritional gaps in the elderly (HUNGER): A prospective study of patient factors associated with inadequate nutritional intake in older medical inpatients. *Clinical Nutrition*. 2011;30:320-5.
- [61] Weekes CE, Spiro A, Baldwin C, et al. A review of the evidence for the impact of improving nutritional care on nutritional and clinical outcomes and cost. *Journal of Human Nutrition & Dietetics*. 2009;22:324-35.
- [62] Milne AC, Avenell A, Potter J. Meta-Analysis: Protein and Energy Supplementation in Older People. *Annals of Internal Medicine*. [Article]. 2006;144(1):37-W14.
- [63] Shepperd S, McClaran J, Phillips CO, et al. Discharge planning from hospital to home. *Cochrane Database of Systematic Reviews*. 2010;Issue 1. Article Number CD000313.
- [64] Feldblum I, German L, Castel H, Harman-Boehm I, Shahar D. Individualised nutritional intervention during and after hospitalisation: The Nutrition Intervention Study Clinical Trial. *Journal of the American Geriatrics Society*. 2011;59:10-7.
- [65] Neelemaat F, Bosmans JE, Thijs A, Seidell JC, van Bokhorst-de van der Schueren MAE. Oral nutritional support in malnourished elderly decreases functional limitations with no extra costs. *Clinical Nutrition*. 2012;31(2):183-90.
- [66] Beck A, Holst M, Rasmussen H. Oral nutritional support of older (65 years+) medical and surgical patients after discharge from hospital: systematic review and meta-analysis of randomized controlled trials. *Clinical Rehabilitation*. 2013;27(19-27).
- [67] Mudge A, Young A, Ross L, et al. Hospital to home outreach for malnourished elders (HHOME) : a feasibility pilot. *Journal of Aging Research and Clinical Practice*. 2012;1(2):131-4.
- [68] Beck A, Kjær S, Hansen B, Storm R, Thal-Jantzen K, Bitz C. Follow-up home visits with registered dietitians have a positive effect on the functional and nutritional status of geriatric medical patients after discharge: a randomized controlled trial. *Clinical Rehabilitation*. 2013;27(6):483-93.
- [69] Cawood A, Elia M, Stratton R. Systematic review and meta-analysis of the effects of high protein oral nutritional supplements. *Ageing Research Reviews*. 2012;11:278-96.

- [70] Baldwin C, Parsons T. Dietary advice and nutritional supplements in the management of illness-related malnutrition: systematic review *Clinical Nutrition*. 2004;23:1267-79.
- [71] Baldwin C, Weekes C, Campbell K. Measuring the effectiveness of dietetic interventions in nutritional support. *Journal of Human Nutrition and Dietetics*. 2008;21(4):303-5.
- [72] Baldwin C, Weekes CE. Dietary advice with or without oral nutritional supplements for disease-related malnutrition in adults. *Cochrane Database of Systematic Reviews*. 2011; Issue 9. Art. No. CD002008.
- [73] Lochs H, Pichard C, Allison S. Evidence supports nutritional support. *Clinical Nutrition*. 2006;25(2):177-9.
- [74] Milne A, Potter J, Vivanti A, Avenell A. Protein and energy supplementation in elderly people at risk from malnutrition. *Cochrane Db Syst Rev*. 2009; Issue 2(Art.No.:CD003288).
- [75] Stratton R, Elia M. Who benefits from nutritional support: what is the evidence? *European Journal of Gastroenterology and Hepatology*. 2007;19:353-8.
- [76] Philipson T, Snider J, Lakdawalla D, Stryckman B, Goldman D. Impact of oral nutritional supplementation on hospital outcomes. *The American Journal of Managed Care*. 2013;19(2):121-8.
- [77] Somanchi M, Tao X, Mullin GE. The Facilitated Early Enteral and Dietary Management Effectiveness Trial in Hospitalized Patients With Malnutrition. *Journal of Parenteral and Enteral Nutrition*. 2011 March 1, 2011;35(2):209-16.
- [78] Wade K, Flett M. Which 'nutritional models-of-care' improve energy and protein intake, clinical outcomes and malnutrition in hospitalised patients? *Nutrition and Dietetics*. 2012;70:7-15.
- [79] Australian and New Zealand Society for Geriatric Medicine (Position Statement No. 6). Under-nutrition and the Older Person. *Australasian Journal on Ageing*. 2009;28(2):99-105.
- [80] Forster S, Gariballa S. Age as a determinant of nutritional status: A cross sectional study. *Nutrition Journal*. 2005;4:28.
- [81] Perera R, Ekanayake L. Relationship between nutritional status and tooth loss in an older population from Sri Lanka. *Gerodontology*. 2012;29(2):e566-e70.
- [82] Stanga Z. Basics in clinical nutrition: Nutrition in the elderly. *e-SPEN, the European e-Journal of Clinical Nutrition and Metabolism*. 2009;4(6):e289-e99.
- [83] Kuczmarski M, Weddle D. Position paper of the American Dietetic Association: Nutrition across the spectrum of aging. *Journal of the American Dietetic Association*. 2005;105(4):616-33.
- [84] Nieuwenhuizen W, Weenan H, Rigby P, Hetherington M. Older adults and patients in need of nutritional support: Review of current treatment options and factors influencing nutritional intake. *Clinical Nutrition*. 2010;29:160-9.
- [85] Patel M, Martin F. Why don't elderly hospital inpatients eat adequately. *Journal of Nutrition Health and Aging*. 2008;12(4):227-31.